

Exposure to Sub-Lethal Doses of Carbofuran Induces Hematological Alterations in Wistar Rats

¹Olusegun K. Afolabi, ²Adedoja D. Wusu, ¹John O. Fatoki and ³Oka S. Ayodeji

¹Department of Biochemistry, Faculty of Basic Medical Sciences, Ladoke Akintola University of Technology, Ogbomoso, Nigeria

²Department of Biochemistry, Faculty of Science, Lagos State University, Lagos, Nigeria

³Department of Medical Biochemistry, Kogi State University, Ayingba, Kogi, Nigeria

Abstract: Carbofuran is a widely used carbamate pesticide that has been detected in ground, surface and rain waters. To assess the toxic effect of carbofuran in this study, Wistar rats numbering twenty-eight, were orally exposed to sub-lethal doses of carbofuran at 1/40th, 1/20th and 1/10th of its LD₅₀. After 50 days of exposure certain hematological parameters were evaluated. A significant decrease ($p < 0.05$) in red blood cell, hemoglobin, hematocrit, mean corpuscular hemoglobin and mean corpuscular volume was observed in the carbofuran-exposed rats. Meanwhile, the exposure resulted in the elevation of the white blood cells, lymphocytes, eosinophils and neutrophils levels in the rats. However, a mixed trend in the thrombocytic indices (total platelet count, mean platelet volume, pro-calcitonin and platelet distribution width) was observed. The exposure to the pesticide, carbofuran produced microcytic anemia and leukocytosis in the rats, in a non-dose dependent manner.

Key words: Carbofuran • Hematology • Hematotoxicity • Thrombocytic Indices • Leucocyte Counts

INTRODUCTION

The past decades have seen an increase in the use of pesticides for the improvement of agricultural yield [1]. Pesticides have also find use in the protection of household and industrial items [2]. The application of these chemicals has toxicological implications for non-target organisms in the environment. The lower toxicities and lesser environmental persistence of some class of pesticides, such as carbamates have resulted in their higher application frequency than most other pesticides [1].

Carbofuran (CF) is a broad spectrum insecticide, acaricide and nematocide belonging to the carbamate group of pesticides and its presence has been reported in nontarget mammalian systems including humans [3]. It accumulates in fat depots and exerts adverse effects on organs such as brain, liver, skeletal muscles and heart [4, 5]. It has also been reported to be toxic to the mammalian nervous system with earlier studies

establishing that carbofuran could inhibit acetylcholine esterase activity [6]. Carbofuran exposure has been found to induce reactive oxygen and nitrogen species, resulting in oxidative stress through the enhancement of malondialdehyde generation and alteration of the activities of antioxidant enzymes such as superoxide dismutase, catalase and glutathione transferase [3]. These reactive species and free radicals are responsible for the peroxidation of membrane phospholipids which disrupt the normal function of lipid bilayer and leading to the development of several pathological conditions.

Detection of carbofuran in ground water and its ingestion through contaminated vegetables and fruits have spurred considerable interest and concern about its use. Since changes in hematological profile can indicate the toxicity of a toxic agent in the body, the present study therefore, investigated the effect of sub-lethal doses of carbofuran on some hematological parameters in Wistar rats.

MATERIALS AND METHODS

Chemicals: Commercial grade carbofuran marketed as Furadan® was obtained from Irorun Agbe Agrochemical Company (Ogbomoso, Nigeria).

Animals and Treatment: Twenty eight (28) adult male rats, weighing between 120 and 150 g were used in this study. The animals were housed at 25±2°C under 12h cycles of dark and light. The animals were allowed access to standard laboratory food and water *ad libitum*. After one-week acclimatization period, they were divided into four groups of seven rats each: a control group, CT 0, CT 1/40th, CT 1/20th and CT 1/10th. The animals were orally received 0, 1/40th, 1/20th and 1/10th LD50 of CF for rats, respectively. The oral LD50 for CF is 8 mg/kg [7]. The treatment was continued for 50 days. At the end of the experimental period, the rats were sacrificed by cervical dislocation after a light anesthesia with ether. Blood samples were collected through cardiac puncture in separate aliquots containing di-potassium salt of EDTA.

Hematological parameters including, red blood cell count (RBC), hemoglobin (Hb), hematocrit (HCT), red blood cell distribution width (RDW), mean corpuscular volume (MCV), mean corpuscular hemoglobin concentration (MCHC), mean corpuscular hemoglobin (MCH), red cell distribution width-standard deviation (RDW-SD), red cell distribution width-coefficient of variation (RDW-CV), white blood cell count (WBC) and differential leucocytic count, thrombocytic indices [Total platelet count (PLT), mean platelet volume (MPV), pro-calcitonin (PCT) and platelet distribution width (PDW)] were analyzed. The analyses were conducted with the use of an automated hematological assay analyzer (Medonic CA 620, Sweden).

Statistical Analysis: Data were analyzed by one-way analysis of variance, followed by Tukey's multiple comparisons test. Results were presented as mean±standard error of means (SEM) and values were considered statistically significant at $p < 0.05$. Data were analyzed using GraphPad Prism for Windows, version 6.01 (GraphPad Software, Inc., San Diego, CA, USA.).

RESULTS

Red Blood Cell Indices: Results showing the effect of sub-lethal doses of carbofuran on the erythrocyte indices of rats are presented in Table 1. After 50 days, RBC count was non-significantly reduced in the CF 1/40th group ($p > 0.05$) but significantly lowered in the CF 1/20th and CF 1/10th groups by 14 and 17% respectively ($p < 0.05$). Hb and HCT on the other hand, were significantly decreased by the exposure to carbofuran, although not in a dose-dependent manner ($p < 0.05$). MCV, MCH and MCHC were also determined in this study. In all the CF-treated groups, MCV and MCH were lowered by as much as, 17, 15 19 and 17, 7 17% respectively. The exposure however, increased the MCHC values, though not significant.

Leucocyte and Differential Leucocyte Counts: Table 2 depicts the effect of CF exposure on the differential leucocytes counts in the rats. Compared to the control, CF significantly caused an increase in WBC, neutrophils and lymphocytes counts in the animals ($p < 0.05$). WBC was increased by 61, 35 71% and the neutrophils by 209, 181 233% in the CF 1/40th, CF 1/20th and CF 1/10th groups, respectively. For the lymphocytes, only the CF 1/20th and 1/10th groups were increased by 32 and 39% respectively. Eosinophils were elevated by CF at the

Table 1: Effect of carbofuran on the hematological parameters of rats

	RBCs ($\times 10^{12}/L$)	Hb (g/dL)	HCT (%)	MCV (fL)	MCH (pg)	MCHC (g/dL)	RDW-CV	RDW-SD
Control	8.31±0.17 ^a	16.87±0.13 ^a	61.07±1.66 ^a	71.63±3.12 ^a	20.40±1.37 ^a	27.70±0.55 ^a	0.20±0.01 ^a	64.53±4.70 ^a
CF 1/40 th	8.58±0.23 ^a	14.57±0.48 ^b	50.80±1.57 ^b	59.17±0.42 ^b	16.97±1.61 ^b	28.67±0.22 ^a	0.17±0.00 ^{bc}	46.53±0.17 ^{bc}
CF 1/20 th	7.14±0.35 ^b	15.52±0.16 ^b	51.98±0.40 ^b	61.00±0.45 ^b	18.90±0.28 ^c	28.75±0.43 ^a	0.18±0.00 ^b	53.15±0.74 ^b
CF 1/10 th	6.90±0.17 ^c	15.57±0.25 ^b	46.90±0.75 ^c	58.23±1.60 ^b	16.97±0.29 ^b	29.20±0.29 ^a	0.16±0.00 ^c	41.97±1.06 ^c

Each value represents the mean±SEM of 7 rats. Values within a column with different alphabets are significantly different at $p < 0.05$.

Table 2: Effect of carbofuran treatment on the total and differential leucocytes counts of rats

	WBCs	Lymphocytes	Neutrophils	Monocytes	Eosinophils	Basophils
Control	6.44±0.12 ^a	4.61±0.18 ^a	0.75±0.14 ^a	1.29±0.10 ^a	0.09±0.01 ^a	0.06±0.01 ^a
CF 1/40 th	10.38±0.38 ^b	4.31±0.25 ^b	2.39±0.13 ^{bc}	1.73±0.04 ^b	0.13±0.01 ^a	0.06±0.01 ^a
CF 1/20 th	8.67±0.18 ^c	6.10±0.10 ^c	2.10±0.03 ^b	0.22±0.04 ^c	0.17±0.01 ^{ab}	0.08±0.00 ^a
CF 1/10 th	11.02±0.36 ^b	6.42±0.34 ^c	2.59±0.11 ^c	0.21±0.03 ^c	0.27±0.05 ^b	0.05±0.02 ^a

Each value represents the mean±SEM of 7 rats. Values within a column with different alphabets are significantly different at $p < 0.05$.

Table 3: Effect of carbofuran treatment on the thrombocytic indices of rats

	PLT ($10^9/L$)	MPV (fL)	PDW (μm)	PCT (%)
Control	405.70 \pm 41.99 ^a	8.60 \pm 0.07 ^a	15.78 \pm 0.09 ^a	3.49 \pm 0.36 ^a
CF 1/40 th	742.30 \pm 34.09 ^b	8.27 \pm 0.08 ^a	15.67 \pm 0.04 ^a	6.12 \pm 0.21 ^b
CF 1/20 th	596.70 \pm 37.00 ^c	7.35 \pm 0.16 ^b	15.75 \pm 0.07 ^a	4.23 \pm 0.30 ^a
CF 1/10 th	560.00 \pm 19.94 ^c	7.60 \pm 0.13 ^b	15.83 \pm 0.15 ^a	4.22 \pm 0.33 ^a

Each value represents the mean \pm SEM of 7 rats. Values within a column with different alphabets are significantly different at $p < 0.05$.

highest dose, 1/10th LD₅₀ by 200. However, with the monocytes, CF modulated the concentrations differently with the low dose, i.e. CF 1/40th causing an increase of 34%, while the other two doses, CF 1/20th and CF 1/10th, decreased the levels by 83 and 84% respectively. Basophils were unaffected by exposure to the pesticide.

Thrombocytic Indices: PLT, MPV and PCT values were all significantly altered in the CF-treated group compared to the control, while PDW was left unchanged (Table 3). PLT in the exposed rats was increased by 83, 47 and 38% in the CF 1/40th, CF 1/20th and CF 1/10th groups, respectively. On the other hand, CF showed no effect on MPV in the CF 1/40th group but depressed this index by 15 and 12% in the CF 1/20th and CF 1/10th groups, respectively. Similarly, the CF 1/40th group displayed no adverse effect of CF on PCT but reduced this index by more than 10% in the other two groups.

DISCUSSION

The widespread use of pesticides results in a number of adverse health effects in both humans and animals. Derangement in hematological parameters is indicative of the toxic effects of foreign compounds such as pesticides, as well as, changes in physiological and pathological status of an organism [8]. Several studies have utilized hematological changes as a biomarker of pesticide exposure and also for the evaluating the interaction between a toxic agent and biological system [9]. Generally, inducement of non-specific immunity by a pesticide leads to alterations in hematological parameters [10]. Carbofuran exposure in the present study resulted in remarkable hematological alterations.

Carbofuran caused a reduction in RBC, HCT and Hb in our study. These alterations may result from the disruptive action of the pesticide on the red cell membranes [11]. Furthermore, the lysing or shrinkage of red blood cells caused by the action of the pesticide on the erythropoietic tissue and the suppression of erythropoiesis may lead to a reduction in red blood cell,

hemoglobin and hematocrit [12]. The decrease in Hb and HCT values in our study indicates a non-specific immune response of the rats to the insecticide. It is also reflective of the rats' defensive reaction against pesticide stress [13]. The observed decrease in MCV and MCH in the treated groups suggested shrinkage of the RBCs. It may also be due to an increase in the level of immature RBCs in the circulation [12]. The MCH reflects Hb content and Hb rate to RBC which were reduced following carbofuran exposure [3]. The decrease indicates microcytic normochromic anemia in the exposed rats [14]. The observed increase in MCHC value, though non-significant, indicates an upsurge in Hb synthesis which could be a compensatory mechanism for the depleted Hb concentration in the carbofuran-exposed rats. The reduction in RDW-SD and RDW-CV, along with RBC, Hb and HCT support the suggestion of an up-regulation of the erythropoiesis in the bone marrow. This might have resulted in the production of impaired red cells or atrophied erythrocytes which would have been destroyed in circulation.

The deleterious effects of carbofuran on the total WBC, neutrophils, lymphocytes and monocytes were evident in this study. The WBC count is a biomarker of systemic inflammation, as the cells are involved in the regulation of immunological function [15]. A significant increase in total WBC, neutrophils and lymphocytes levels, along with a significant increase in the platelet count were observed in the carbofuran-exposed rats. This indicates either a protective mechanism against the pesticide or the activation of the rat immune system against the carbofuran-induced stress. An increased WBC count can be adduced to an increased antibody production in order to ameliorate pesticide-induced toxicity [16]. Leukocytosis in this study may occur as a result of increased leukocyte mobilization, reflective of the severity of causative stress condition [17]. Other pesticides have been reported to stimulate immune functions by increasing WBC levels, indicating the activation of defense and immune system where oedema and inflammation have been induced [18]. The increase in leucocytes was particularly evident in the case of lymphocytes which could have resulted from the stimulation of lymphopoietic process in the rats. Leukocytosis has been demonstrated in animals intoxicated with pesticides and has been suggested to be due to the mobilization of the immunological system and/or a shift in the leucocytic pool from the spleen to peripheral blood [19]. Non-specific irritation caused by the toxicant and/or its metabolites may induce the production

and release of inflammatory mediators like prostaglandins that produce neutrophilia. Neutrophilia can lead to enhanced release of mature neutrophils from the bone marrow and the subsequent demargination, contributing to the circulating pool [20]. A significant increase in PLT counts was found in all the CF-treated groups. Pesticides can induce histopathological damage [21] and such damage can increase the production of PLT, as well as WBC as seen in our study.

CONCLUSION

It can thus, be concluded that sub-lethal doses of carbofuran elicited hematological alteration, which was characterized by microcytic normochromic anemia and leukocytosis. Also, the modulated hematotoxic response induced by the pesticide in the rats was not dose-dependent. The data from this study further provide evidence of the toxic effect of carbofuran.

REFERENCES

1. Jaiswal, S.K., N.J. Siddiqi and B. Sharma, 2013. Carbofuran induced oxidative stress in rat heart: ameliorative effect of vitamin C. *ISRN Oxidative Medicine*, 2013: 1-10.
2. Milatovic, D., R.C. Gupta and M. Aschner, 2006. Anticholinesterase toxicity and oxidative stress. *The Scientific World Journal*, 6: 295-310.
3. Hossen, M.S., E. Tanvir, M.B. Prince, S. Paul, M. Saha, M.Y. Ali, S.H. Gan, M.I. Khalil and N. Karim. 2017. Protective mechanism of turmeric (*Curcuma longa*) on carbofuran-induced hematological and hepatic toxicities in a rat model. *Pharmaceutical Biology*, 55(1): 1937-1945.
4. de Siqueira, A., F.A. Salvagni, A.S. Yoshida, V. Gonçalves-Junior, A.S. Calefi, A.R. Fukushima, H. de Souza Spinosa and P.C. Maiorka. 2015. Poisoning of cats and dogs by the carbamate pesticides aldicarb and carbofuran. *Research in Veterinary Science*, 102: 142-149.
5. Ruiz-Suárez, N., L.D. Boada, L.A. Henríquez-Hernández, F. González-Moreo, A. Suárez-Pérez, M. Camacho, M. Zumbado, M. Almeida-González, M. del Mar Travieso-Aja and O.P. Luzardo, 2015. Continued implication of the banned pesticides carbofuran and aldicarb in the poisoning of domestic and wild animals of the Canary Islands (Spain). *Science of the Total Environment*, 505: 1093-1099.
6. Gupta, V.K., A. Pathak, N.J. Siddiqi and B. Sharma. 2016. Carbofuran modulating functions of acetylcholinesterase from rat brain *in vitro*. *Advances in Biology*, 2016: 1-7.
7. Rai D.K., R.K. Sharma, P.K. Rai, G. Watal, B. Sharma. 2011. Role of aqueous extract of *Cynodon dactylon* in prevention of carbofuran-induced oxidative stress and acetylcholinesterase inhibition in rat brain. *Cellular and Molecular Biology*, 57(1): 133-142.
8. Hu R., X. Huang, J. Huang, Y. Li, C. Zhang, Y. Yin, Z. Chen, Y. Jin, J. Cai, F. Cui. 2015. Long- and short-term health effects of pesticide exposure: a cohort study from China. *PLOS One*, 10(6): 1-11.
9. Da Cuna, R.H., G.R. Vazquez, M.N. Piol, N.V. Guerrero, M.C. Maggese and F.L.L. Nostro, 2011. Assessment of the acute toxicity of the organochlorine pesticide endosulfan in *Cichlasoma dimerus* (Teleostei, Perciformes). *Ecotoxicology and Environmental Safety*, 74(4): 1065-1073.
10. Aroonvilairat, S., W. Kespichayawattana, T. Sornprachum, P. Chaisuriya, T. Siwadune and K. Ratanabanangkoon, 2015. Effect of pesticide exposure on immunological, hematological and biochemical parameters in Thai orchid farmers—a cross-sectional study. *International Journal of Environmental Research and Public Health*, 12(6): 5846-5861.
11. Ramesh, M. and M. Saravanan, 2008. Haematological and biochemical responses in a freshwater fish *Cyprinus carpio* exposed to chlorpyrifos. *International Journal of Integrative Biology*, 3(1): 80-83.
12. Saravanan, M., K.P. Kumar and M. Ramesh, 2011. Haematological and biochemical responses of freshwater teleost fish *Cyprinus carpio* (Actinopterygii: Cypriniformes) during acute and chronic sublethal exposure to lindane. *Pesticide Biochemistry and Physiology*, 100(3): 206-211.
13. Narra, M.R., 2016. Single and cartel effect of pesticides on biochemical and haematological status of *Clarias batrachus*: a long-term monitoring. *Chemosphere*, 144: 966-974.
14. Adhikari, S., B. Sarkar, A. Chatterjee, C. Mahapatra and S. Ayyappan, 2004. Effects of cypermethrin and carbofuran on certain hematological parameters and prediction of their recovery in a freshwater teleost, *Labeo rohita* (Hamilton). *Ecotoxicology and Environmental Safety*, 58(2): 220-226.

15. Farhangi, M.A., S.A. Keshavarz, M. Eshraghian, A. Ostadrahimi A and A.A. Saboor-Yaraghi, 2013. White blood cell count in women: relation to inflammatory biomarkers, haematological profiles, visceral adiposity and other cardiovascular risk factors. *Journal of Health, Population and Nutrition*, 31(1): 58.
16. Joshp, P., M. Bose and D. Harish, 2002. Changes in certain haematological parameters in a siluroid cat fish *Clarias batrachus* (Linn) exposed to cadmium chloride. *Pollution Research*, 21(2): 129-131.
17. Celik, I., Z. Yilmaz and V. Turkoglu, 2009. Hematotoxic and hepatotoxic effects of dichlorvos at sublethal dosages in rats. *Environmental Toxicology: An International Journal*, 24(2): 128-132.
18. Yousef, M., F. El-Demerdash, K. Kamel and Al-Salhen, 2003. Changes in some hematological and biochemical indices of rabbits induced by isoflavones and cypermethrin. *Toxicology*, 189(3): 223-234.
19. Haratym-Maj, A., 2003. Hematological alternations after pyrethroids poisoning in mice. *Annals of Agricultural and Environmental Medicine*, 9(2): 199-206.
20. Khan, A.M., N. Dubey, R. Raina, G. Singh and S.A. Beigh, 2013. Toxic effects of deltamethrin and fluoride on hematological parameters in rats. *Fluoride*, 46(1): 34-38.
21. Afolabi, O.K., A.D. Wusu, R. Ugbaja and J.O. Fatoki, 2018. Aluminium phosphide-induced testicular toxicity through oxidative stress in Wistar rats: Ameliorative role of hesperidin. *Toxicology Research and Application*, 2: 1-11.