

Response of Yield and Quality of Sugar Beet to Different Tillage Methods

¹Majid Rashidi, ²Saeed Mohammadi, ³Ali Sahebi Kouzekanan and ²Mohammad Rahmani

¹Department of Agricultural Machinery, Takestan Branch, Islamic Azad University, Takestan, Iran

²Tehran Province Agricultural and Natural Resources Research Center, Varamin, Tehran, Iran

³Department of Agronomy, Roudehen Branch, Islamic Azad University, Roudehen, Tehran, Iran

Abstract: Field experiments were carried out at the Research Site of Hamedan Province, Iran to study the response of yield and quality of sugar beet to different tillage methods during 2008 and 2009 growing seasons. Tillage treatments were moldboard plow + two passes of disk harrow (MDD) as conventional tillage method; moldboard plow + one pass of rotavator (MR), chisel plow + one pass of rotavator (CR) and two passes of disk harrow (DD) as reduced tillage methods; one pass of rotavator (R) and one pass of tine cultivator (C) as minimum tillage methods and no-tillage (NT). Yield, yield components, i.e. root number per hectare (RNPH), sugar yield (SUGY), root dry matter (RODM), root length (ROTL), rim diameter (RIMD) and some selected quality characteristics of sugar beet such as sugar content (SUGC), potassium (POTA), sodium (SODI), alpha-amino nitrogen (ALAN) and molasses (MOLA) were measured for different tillage treatments. Different treatments significantly ($P \leq 0.05$) affected RNPH and POTA, but there was no significant difference in other studied traits. Although effect of different tillage treatments on yield, SUGY, RODM, ROTL, RIMD, SUGC, SODI, ALAN and MOLA was not significant, results of the study indicated that tillage operations were useful in improving the yield and quality of sugar beet. The highest values of yield (82.7 t ha^{-1}), RNPH ($135412 \text{ roots ha}^{-1}$), SUGY (11.4 t ha^{-1}), RODM (23.9%) and SUGC (17.0%) were recorded in the MR treatment, while the highest values of ROTL (20.5 cm), RIMD (1.5 cm), POTA (6.4 mmol/100 g), SODI (2.6 mmol/100 g), ALAN (2.5 mg/100 g) and MOLA (3.0%) were noted in the NT treatment. In contrast, the lowest values of yield (71.3 t ha^{-1}), RNPH ($115000 \text{ roots ha}^{-1}$), SUGY (9.15 t ha^{-1}), RODM (20.3%) and SUGC (15.2%) were recorded in the NT treatment, while the lowest values of ROTL (18.0 cm), RIMD (1.1 cm), POTA (4.5 mmol/100 g), SODI (1.5 mmol/100 g), ALAN (1.6 mg/100 g) and MOLA (2.2%) were noted in the MR treatment. Results also showed that the reduced tillage treatments MR and CR and the minimum tillage treatment R were considered as more beneficial and suitable tillage methods in improving the yield and quality of sugar beet.

Key words: Sugar beet • Tillage method • Yield • Quality • Hamedan • Iran

INTRODUCTION

Sugar beet (*Beta vulgaris*) is one of the most important crops in a wide variety of temperature climates [1-3]. It is a hardly biennial plant with large (1-2 kg) storage root and great amount (15-20%) of sucrose. Sugar beet accounts for 30% of the world's sugar production [4]. The European Union, the United States and Russia are the three biggest sugar beet producers in the world. The top ten sugar beet producer countries are France, Germany, United States, Russia, Ukraine, Turkey, Italy, Poland, United Kingdom and Spain with 29, 25, 25, 22, 16, 14, 12, 11, 8 and 7 million tons, respectively.

Also, the European Union and Ukraine are major exporters of sugar from beets. Besides, the United States harvested 406,500 hectares of sugar beets in 2008 alone [5]. On the other hand, the average cultivated area and national production of sugar beet in Iran for the last three years was about 178,000 hectares and 5.9 million tons, respectively [6]. Although the use of better varieties, mechanical planting, chemical fertilizers, herbicides application and mechanized harvesting have increased sugar beet production to a great extent, the complete potential of sugar beet production has not yet been attained as compared to the top ten sugar beet producers.

Tillage is one of the most essential crop production factors that influence soil properties [7, 8] and consequently crop yield [9-14]. Appropriate tillage operations can enhance soil properties, while excessive, inappropriate and unnecessary tillage operations may result in a range of undesirable processes [15-20]. Although for most situations, conventional tillage methods have been the main tillage methods for establishing sugar beet since the first part of the 20th century, they are now expensive operations in terms of work rate and fuel consumption [21]. The costs, as well as the environmental concerns have led farmers and researchers to adopt alternative tillage methods [22]. For these reasons, there is a considerable attention and significant emphasis on moving towards the conservation tillage methods, i.e. reduced tillage, minimum tillage and no-tillage methods [7, 8, 10-15, 20, 23-27]. Conservation tillage methods may be used for sugar beet [28-31]. However, the results of these methods may be contrary [20]. Conservation tillage operations may reduce yield of sugar beet [4]. Conversely, decrease of soil tillage practices may have no significant effect on the yield of other crops [25-27, 32, 33]. Conservation tillage methods may also lead to raised diversity of weed species and population [33, 34] and have a harmful effect on crop yield [35]. But, other studies have confirmed the opposite [36].

In Iran, most of the cultivated area is under conventional tillage methods and conservation tillage methods have not been studied enough. Therefore, this study was conducted to study the response of yield and quality of sugar beet to different tillage methods.

MATERIALS AND METHODS

Research Site: This study was conducted at the Research Site of Hamedan Province, Iran for two successive growing seasons (2008 & 2009). The research site is located at latitude of 34° 52' N, longitude of 48° 21' E and altitude of 1730 m in semi-arid climate (298 mm rainfall annually) in the west of Iran. Mean temperature and monthly rainfall of the experimental site from sowing to harvest during study years (2008 & 2009) are indicated in Fig. 1.

Soil Sampling and Analysis: A composite soil sample (from 21 points) was collected from 0-30 cm depth during the study years and was analyzed in the laboratory for pH, EC, OC, N, P, K, Fe, Zn, Cu, Mn, B and particle size distribution. Details of soil physical and chemical properties of the research site during both years (2008 & 2009) are given in Table 1.




Fig. 1: Mean temperature and monthly rainfall during crop growth (mean of 2008 & 2009)

Field Methods: The experiments were laid out in a RCBD with four replications. Tillage treatments were moldboard plow + two passes of disk harrow (MDD) as conventional tillage method; moldboard plow + one pass of rotavator (MR), chisel plow + one pass of rotavator (CR) and two passes of disk harrow (DD) as reduced tillage methods; one pass of rotavator (R) and one pass of tine cultivator (C) as minimum tillage methods and no-tillage (NT). During the study years, tillage treatments were carried out on the same plots. The size of each plot was 20.0 m long and 6.0 m wide. There were 12 rows of sugar beet in each plot with 50-cm row spacing. In both years of study, one of the commercial varieties of sugar beet cv. Zarghan was planted on April 3, 2008 and April 5, 2009 using a 6-row sugar beet drill. Recommended levels of urea (300 kg ha⁻¹) in both years and triple super phosphate (50 kg ha⁻¹) only in the first year of study were used. For all treatments, irrigation scheduling was based on the basis of evaporation from A-class pan installed close to the experimental plots. Also, pest and weed control operations were performed based on common local practices and commendations. All other essential operations were kept identical for all the treatments.

Observation and Data Collection: At harvest, plants from an area of 12.0 m² per each plot were harvested to determine yield and yield components, i.e. root number per hectare (RNPH), sugar yield (SUGY), root dry matter (RODM), root length (ROTL) and rim diameter (RIMD) for all treatments. Moreover, a sample of 20 kg of sugar beet roots were taken at random and sent to the Sugar Beet Laboratory at Hamedan Sugar Factory to determine some quality characteristics, i.e. sugar content (SUGC), potassium (POTA), sodium (SODI), alpha-amino nitrogen (ALAN) and molasses (MOLA) for all treatments.

Table 1: Soil physical and chemical properties of the experimental site (0-30 cm depth), 2008 & 2009

Date	pH	EC (dS m ⁻¹)	OC (%)	N (%)	P (ppm)	K (ppm)	Fe (ppm)	Zn (ppm)	Cu (ppm)	Mn (ppm)	B (ppm)	Soil texture
2008	7.9	0.72	0.92	0.09	10.5	280	6.2	0.8	2.3	16.2	0.7	Loam
2009	8.3	0.55	0.36	0.04	25.6	310	6.4	1.0	2.4	14.4	0.7	Loam

SUGC (sucrose content) was measured in fresh root samples by using Saccharometer as described by AOAC [37]. POTA, SODI, ALAN and MOLA were measured using an auto analyzer.

Statistical Analysis: All data were subjected to the Analysis of Variance (ANOVA) following Gomez and Gomez [38] using SAS statistical computer software. Moreover, means of the different treatments were separated by Duncan's Multiple Range Test (DMRT) at $P \leq 0.05$.

RESULTS

Results of ANOVA and means comparison for yield and yield components of sugar beet under different methods of tillage during the study years (mean of 2008 & 2009) are presented in Tables 2 and 3, respectively. Results showed that different tillage methods significantly ($P \leq 0.05$) influenced RNPH, but there was no significant difference in other studied traits (Table 2). Moreover, results of ANOVA and means comparison for the selected quality characteristics of sugar beet under different tillage methods during the years of study (mean of 2008 & 2009) are presented in Tables 4 and 5, respectively. Results also showed that different methods of tillage significantly ($P \leq 0.05$) influenced POTA, but there was no significant difference in other studied traits (Table 4).

DISCUSSION

In this study, yield, yield components (RNPH, SUGY, RODM, ROTL and RIMD) and some selected quality characteristics (SUGC, POTA, SODI, ALAN and MOLA) of sugar beet were investigated to study the response of yield and quality of sugar beet to different tillage methods.

Yield and Yield Components: The highest value of RNPH (135412 roots ha⁻¹) was recorded in the MR treatment, while the lowest value of RNPH (115000 roots ha⁻¹) was recorded in the NT treatment (Table 3). Although there was no significant difference in yield, SUGY, RODM, ROTL and RIMD during the study years, results indicated that tillage operations were useful in increasing the yield of sugar beet. The highest values of yield (82.7 t ha⁻¹),

SUGY (11.4 t ha⁻¹) and RODM (23.9%) were recorded in the MR treatment, while the highest values of ROTL (20.5 cm) and RIMD (1.5 cm) were noted in the NT treatment (Table 3). Based on the results, tillage method affected the yield of sugar beet (yield, SUGY and RODM) in the order of MR > CR > R > MDD > DD > C > NT. These results are in line with the results reported by Khurshid *et al.* [9], Rashidi & Keshavarzpour [10], Rashidi *et al.* [11], Rashidi & Khabbaz [12] and Iqbal *et al.* [20] that tillage practices can be associated with improved soil physical and mechanical properties (increased pore space, decreased bulk density, increased moisture preservation and decreased penetration resistance), enhanced soil structure, better seed-soil/root-soil contact and superior weed control which positively influence RNPH and consequently yield and SUGY of sugar beet. Similar results were also obtained by Romanekas *et al.* [28, 30], Adamaviciene *et al.* [29] and Jabro *et al.* [31]. They concluded that intensive tillage methods enhanced soil quality and had no significant effect on yield and most yield components of sugar beet. In contrast, the lowest values of yield (71.3 t ha⁻¹), SUGY (9.15 t ha⁻¹) and RODM (20.3%) were recorded in the NT treatment, while the lowest values of ROTL (18.0 cm) and RIMD (1.1 cm) were noted in the MR treatment (Table 3). These results are in agreement with those of Hill [15], Horne *et al.* [16], Ozpinar [33], Carter & Ivany [34], Borresen [35] and Bauder *et al.* [39] who concluded that conservation tillage methods may be associated with worse soil physical and mechanical properties (decreased pore space, increased bulk density, decreased moisture preservation and increased penetration resistance), inferior seed/root-soil contact and raised diversity of weed species and population which negatively influence RNPH and as a results yield and SUGY of sugar beet.

Selected Quality Characteristics: The highest value of POTA (6.4 mmol/100 g) was recorded in the NT treatment, while the lowest value of POTA (4.5 mmol/100 g) was noted in the MR treatment (Table 5). Although there was no significant difference in SUGC, SODI, ALAN and MOLA during the years of study, results again indicated that tillage operations were useful in enhancing the quality of sugar beet. The highest value of SUGC (17.0%) was recorded in the MR treatment, while the highest values of SODI (2.6 mmol/100 g), ALAN (2.5 mg/100 g)

Table 2: Analysis of variance for yield and yield components of sugar beet under different tillage methods (mean of 2008 & 2009)

Source of Variation	Df	Mean square					
		Yield	RNPH	SUGY	RODM	ROTL	RIMD
Replication	3	257.9 ^{NS}	127777616 ^{NS}	6.10 ^{NS}	7.21 ^{NS}	1.32 ^{NS}	0.10 ^{NS}
Treatment	6	72.36 ^{NS}	184223872*	3.27 ^{NS}	5.96 ^{NS}	3.40 ^{NS}	0.08 ^{NS}
Error	18	390.7	62268312	10.5	3.17	5.12	0.04
C.V. (%)	---	25.4	6.20	31.3	8.04	11.7	15.2

NS = Non-significant

* = Significant at 0.05 probability level

(RNPH: root number per hectare; SUGY: sugar yield; RODM: root dry matter; ROTL: root length; RIMD: rim diameter)

Table 3: Means comparison for yield and yield components of sugar beet between different tillage methods (mean of 2008 & 2009)

Treatment	Yield (t ha ⁻¹)	RNPH	SUGY (t ha ⁻¹)	RODM (%)	ROTL (cm)	RIMD (cm)
MDD	78.5 a	130000 a	10.5 a	22.0 a	19.6 a	1.2 a
MR	82.7 a	135412 a	11.4 a	23.9 a	18.0 a	1.1 a
CR	81.0 a	133333 a	11.2 a	23.4 a	18.6 a	1.1 a
DD	76.5 a	127500 a	9.97 a	21.8 a	19.6 a	1.3 a
R	80.9 a	130833 a	10.8 a	22.4 a	18.9 a	1.2 a
C	73.4 a	124583 ab	9.27 a	21.3 a	20.4 a	1.3 a
NT	71.3 a	115000 b	9.15 a	20.3 a	20.5 a	1.5 a

Means in the same column with different letters differ significantly at 0.05 probability level according to DMRT.

(RNPH: root number per hectare; SUGY: sugar yield; RODM: root dry matter; ROTL: root length; RIMD: rim diameter)

Table 4: Analysis of variance for some selected quality characteristics of sugar beet under different tillage methods (mean of 2008 & 2009)

Source of variation	Df	Mean square				
		SUGC	POTA	SODI	ALAN	MOLA
Replication	3	8.78 ^{NS}	0.22 ^{NS}	0.33 ^{NS}	0.78 ^{NS}	0.12 ^{NS}
Treatment	6	3.03 ^{NS}	0.56*	0.60 ^{NS}	0.54 ^{NS}	0.27 ^{NS}
Error	18	13.4	0.15	0.68	0.65	0.11
C.V. (%)	---	28.9	7.04	43.0	40.5	13.3

NS = Non-significant

* = Significant at 0.05 probability level

(SUGC: sugar content; POTA: potassium; SODI: sodium; ALAN: alpha-amino nitrogen; MOLA: molasses)

Table 5: Means comparison for some selected quality characteristics of sugar beet between different tillage methods (mean of 2008 & 2009)

Treatment	SUGC (%)	POTA (mmol/100 g)	SODI (mmol/100 g)	ALAN (mg/100 g)	MOLA (%)
MDD	16.8 a	5.4 b	1.9 a	1.9 a	2.4 a
MR	17.0 a	4.5 b	1.5 a	1.6 a	2.2 a
CR	17.0 a	5.3 b	1.6 a	1.7 a	2.3 a
DD	15.6 a	5.5 b	2.0 a	2.1 a	2.5 a
R	16.9 a	5.4 b	1.6 a	1.7 a	2.4 a
C	15.2 a	5.7 b	2.2 a	2.5 a	2.5 a
NT	15.2 a	6.4 a	2.6 a	2.5 a	3.0 a

Means in the same column with different letters differ significantly at 0.05 probability level according to DMRT.

(SUGC: sugar content; POTA: potassium; SODI: sodium; ALAN: alpha-amino nitrogen; MOLA: molasses)

and MOLA (3.0%) were noted in the NT treatment. In contrast, the lowest value of SUGC (15.2%) was recorded in the NT treatment, while the lowest values of SODI (1.5 mmol/100 g), ALAN (1.6 mg/100 g) and MOLA (2.2%)

were noted in the MR treatment (Table 5). Again, a similar trend was obtained for the selected quality characteristics and tillage method affected sugar beet quality in the order of MR > CR > R > MDD > DD > C > NT (Table 5). Similar

results were also obtained by Romaneckas *et al.* [28, 30], Adamaviciene *et al.* [29] and Jabro *et al.* [31]. They reported that different methods of tillage had no significant effect on most quality characteristics of sugar beet.

CONCLUSION

Different tillage methods significantly ($P \leq 0.05$) affected RNPH and POTA, but there was no significant difference in yield, SUGY, RODM, ROTL, RIMD, SUGC, SODI, ALAN and MOLA. Although there was no significant difference in most studied traits, tillage operations were useful in improving the yield and quality of sugar beet. Also, the reduced tillage treatments MR and CR and the minimum tillage treatment R were considered as more beneficial and suitable tillage methods in improving the yield and quality of sugar beet.

REFERENCES

1. Jafari, A., S.S. Mohtasebi, H.E. Jahromi and M. Omid, 2006. Weed detection in sugar beet fields using machine vision. *Int. J. Agri. Biol.*, 8: 602-605.
2. Sohrabi, Y. and G. Heidari, 2008. Influence of withholding irrigation and harvest times on yield and quality of sugar beet (*Beta vulgaris*). *Int. J. Agri. Biol.*, 10: 427-431.
3. Abdel-Motagally, F.M.F. and K.K. Attia, 2009. Response of sugar beet plants to nitrogen and potassium fertilization in sandy calcareous soil. *Int. J. Agric. Biol.*, 11: 695-700.
4. Draycott, A.P., 2006. Sugar Beet. Blackwell Publishing Ltd., Oxford, UK.
5. Food and Agriculture Organization of the United Nations; FAO Statistical Yearbook, 2009.
6. Iranian Ministry of Agriculture, Statistical Yearbook, 2009.
7. Keshavarzpour, F. and M. Rashidi, 2008. Effect of different tillage methods on soil physical properties and crop yield of watermelon (*Citrullus vulgaris*). *World Appl. Sci. J.*, 3: 359-364.
8. Rashidi, M. and F. Keshavarzpour, 2008. Effect of different tillage methods on soil physical properties and crop yield of melon (*Cucumis melo*). *American-Eurasian J. Agric. and Environ. Sci.*, 3: 31-36.
9. Khurshid, K., M. Iqbal, M.S. Arif and A. Nawaz, 2006. Effect of tillage and mulch on soil physical properties and growth of maize. *Int. J. Agri. Biol.*, 8: 593-596.
10. Rashidi, M. and F. Keshavarzpour, 2007. Effect of different tillage methods on grain yield and yield components of maize (*Zea mays* L.). *Int. J. Agri. Biol.*, 9: 274-277.
11. Rashidi, M., F. Keshavarzpour and M. Gholami, 2008. Effect of different tillage methods on yield and yield components of forage corn. *American-Eurasian J. Agric. and Environ. Sci.*, 3: 347-351.
12. Rashidi, M. and B.G. Khabbaz, 2009. Response of crop yield and yield components of tomato to different tillage methods in the arid lands of Varamin, Iran. In: Proceedings of Biennial Conference of the Australian Society for Engineering in Agriculture (SEAg). September 13-16, Brisbane, QLD, Australia.
13. Rashidi, M., M. Gholami and B.G. Khabbaz, 2009a. Response of yield and yield components of tomato (*Lycopersicon esculentum*) to different tillage methods. *Int. J. Agri. Biol.*, 11: 626-630.
14. Rashidi, M., S. Abbassi and M. Gholami, 2009b. Interactive effects of plastic mulch and tillage method on yield and yield components of tomato (*Lycopersicon esculentum*). *American-Eurasian J. Agric. and Environ. Sci.*, 5: 420-427.
15. Hill, R.L., 1990. Long-term conventional and no-tillage effects on selected soil physical properties. *Soil Sci. Soc. Amer. J.*, 54: 161-166.
16. Horne, D.J., C.W. Ross and K.A. Hughes, 1992. Ten years of maize/oats rotation under three tillage systems on a silt-loam soil in New Zealand. 1. A comparison of some soil properties. *Soil Till. Res.*, 22: 131-143.
17. Lal, R., 1993. Tillage effects on soil degradation, soil resilience, soil quality and sustainability. *Soil Till. Res.*, 51: 61-70.
18. Khan, F.U.H., A.R. Tahir and I.J. Yule, 1999. Impact of different tillage practices and temporal factor on soil moisture content and soil bulk density. *Int. J. Agri. Biol.*, 1: 163-166.
19. Khan, F.U.H., A.R. Tahir and I.J. Yule, 2001. Intrinsic implication of different tillage practices on soil penetration resistance and crop growth. *Int. J. Agri. Biol.*, 3: 23-26.
20. Iqbal, M., A.U. Hassan, A. Ali and M. Rizwanullah, 2005. Residual effect of tillage and farm manure on some soil physical properties and growth of wheat (*Triticum aestivum* L.). *Int. J. Agri. Biol.*, 7: 54-57.
21. Ecclestone, P., 2004. To plough or not to plough? *British Sugar Beet Review*, 72: 7-11.
22. Ecclestone, P., 2001. Minimum tillage, options for economic sugar beet production. *British Sugar Beet Review*, 69: 24-29.

23. Cannel, R.Q., 1985. Reduced tillage in north-west Europe - a review. *Soil Till. Res.*, 5: 129-177.
24. Chaudhary, A.D., M. Javed, M.A. Rana, A. Sarwar and Q. Zaman, 1992. Comparative performance of direct drilling and conventional tillage practices under rice-wheat rotation system. *Pakistan J. Agric. Sci.*, 29: 5-8.
25. Ekeberg, E., 1993. Minimum tillage for potatoes. *Soil Tillage and Environment. The 228th Seminar of NJF. Finland.*
26. Hao, X., C. Chang, R.L. Conner and P. Bergen, 2001. Effect of minimum tillage and crop sequence on crop yield and quality under irrigation in Southern Alberta clay loam soil. *Soil Till. Res.*, 59: 45-55.
27. Glab, T. and B. Kulib, 2008. Effect of mulch and tillage system on soil porosity under wheat (*Triticum aestivum*). *Soil Till. Res.*, 99: 169-178.
28. Romaneckas, K., R. Romaneckiene and E. Sarauskis, 2006. The effect of primary soil tillage methods on sugar beet growth on a light loam luvisol. *Zemdirbyste/Agriculture*, 93: 81-87.
29. Adamaviciene, A., K. Romaneckas, E. Sarauskis and V. Pilipavicius, 2009. Non-chemical weed control in sugar beet crop under an intensive and conservation soil tillage pattern: II. Crop productivity, *Agronomy Res.*, 7: 143-148.
30. Romaneckas, K., R. Romaneckiene and V. Pilipavicius, 2009. Non-chemical weed control in sugar beet crop under intensive and conservation soil tillage: I. Crop weediness, *Agronomy Res.*, 7: 457-464.
31. Jabro, J.D., W.B. Stevens, W.M. Iversen and R.G. Evans, 2010. Tillage depth effects on soil physical properties, sugar beet yield and sugar beet quality. *Comm. Soil Sci. Plant Anal.*, 41: 908-916.
32. Hakansson, I., M. Stenberg and T. Rydberg, 1998. Long-term experiments with different depth of mouldboard ploughing in Sweden. *Soil Till. Res.*, 46: 209-223.
33. Ozpinar, S., 2006. Effects of tillage systems on weed population and economic for winter wheat production under the Mediterranean dry land conditions. *Soil Till. Res.*, 87: 1-8.
34. Carter, M.R. and J.A. Ivany, 2006. Weed seed bank composition under three long-term tillage regimes on a fine sandy loam in Atlantic Canada. *Soil Till. Res.*, 90: 29-38.
35. Borresen, T., 1993. The long-term effect of tillage practice on soil properties and crop yields. *Soil Tillage and Environment. The 228th Seminar of NJF. Finland.*
36. Campbell, C.A., A.G. Thomas, V.O. Biederbeck, B.G. McConkey, F. Selles, D. Spurr and R.P. Zentner, 1998. Converting from no tillage to pre-seeding tillage: Influence on weeds, spring wheat grain yields and soil quality. *Soil Till. Res.*, 46: 175-185.
37. AOAC, 1995. Association of Official Analytical Chemists. Official methods of analysis, 16th edition, AOAC International, Washington, DC.
38. Gomez, K.A. and A.A. Gomez, 1984. *Statistical Procedures for Agriculture Research.* A Wiley-Inter Science Publication, John Wiley and Sons Inc., New York, USA.
39. Bauder, J.W., G.W. Randall and J.B. Swan, 1981. Effects of four continue tillage systems on mechanical impedance of a clay-loam soil. *Soil Sci. Soc. Amer. J.*, 45: 802-806.