

Assessment of Public Knowledge, Attitude and Practices towards Anthrax in Dessie City, Ethiopia

Tigist Awraris Dejene

Amahara Regional State, Dessie City Administration,
Agricultural Office, Animal Health Case Team, Ethiopia

Abstract: A cross-sectional study was conducted from June 2022 to September 2022 to assess the knowledge, attitudes and practices (KAP) on Anthrax and associated risk factors among the community of Dessie City, Amhara regional state, Ethiopia. A simple random sampling procedure was employed to select kebeles. From the list of kebeles, five were randomly selected using the lottery method. Then, 40 households were selected and interviewed from each kebele using a systematic random sampling method. A structured questionnaire was used to collect the data through face-to-face interviews among 190 respondents. Then, the data were analyzed using Excel spreadsheets. Out of the 190 respondents interviewed, 65.2 % of them were males and 34.7 % were female. The majority of the respondents 63.6 % were Muslim followed by Orthodox 36.3%. Of the total of respondents, (55.2%) had awareness about Anthrax. The result also established that (28.4%) of them know that eating raw meat transmit diseases from animal to humans. Sudden death (46.8%) followed by stop eating and drinking (19.5%) were described as a major clinical sign of Anthrax in animals by the majority of the respondents. Of the participants, (51.1%) knew that the disease could not easily treatable. From the total of respondents, (31.5%) respondents agree on Grazing pasture is source of infection for animal. Of the participants, (54.1%) were aware of the fact that Anthrax is a very serious disease in humans and animals. Regarding safety measures (79.4%) respondents take any safety measure when they are caring anthrax-suspected animals. The majority of the respondents (87.8%) were aware of animal vaccination as a means of anthrax prevention. More than (65%) of the respondents vaccinate their animals while (34.7%) didn't vaccinate their animals. Educating the community about the health risks of anthrax and the ways of prevention should be given priority by health extension workers, veterinary professionals and the government at large. Medical and veterinary personnel should be collaborating in a multidisciplinary approach for the prevention and control of anthrax.

Key words: Anthrax • Attitude • Dessie City • Ethiopia • Knowledge • Practice

INTRODUCTION

Anthrax is a zoonotic bacterial disease caused by *Bacillus anthracis*, a gram positive rod-shaped and capsulated bacillus [1]. It is the most common bacterial disease in sub-Saharan countries [2].

The name of the bacterium is derived from “anthracis”, the Greek word for coal, because anthrax in humans causes black, coal-like lesions on the skin at the site of inoculation [3]. The bacterium form spores when exposed to oxygen and allows it to remain viable in the environment for many years before coming into contact

with a susceptible host and when exposed to a nutrient-rich environment, such as the tissues or blood of an animal or human host [4].

The disease in animals is characterized by septicemia and sudden death with exudation of tarry blood from the natural orifices of the cadaver. Failure of the blood to clot, the absence of rigor mortis and the presence of splenomegaly are the most important necropsy findings of the disease. Prior to the development of the anthrax vaccine and antibiotics it was the foremost cause of uncontrolled mortality in different species of animals worldwide [5].

Animal anthrax is an endemic disease and seasonal in Ethiopia which occurs in May and June every year in different localities of the country. Several districts of the country are reporting suspected cases of anthrax outbreaks in animals and few of which are confirmed by laboratories [6]. A number of factors such as changing rainfall patterns, soil disturbance, increased animal and human populations, and poor grazing systems and human behavior have been reported to be associated with outbreaks of anthrax. Interaction of wildlife with livestock and humans has also been reported as a key predisposing factor of anthrax among humans and livestock. The Disease usually re-occurs in areas where there has been a previous outbreak, making vaccination the recommended form of control [7].

The objectives of this study were;

- To assess the level of knowledge, attitudes and practices of communities in Dessie city, Ethiopia.
- To identify factors associated with community knowledge, attitudes and practices about anthrax in the study area.

MATERIALS AND METHODS

Study Area: The study was conducted from June 2022 to September 2022 to assess the level of knowledge, attitudes and practices towards Anthrax in Dessie city, Amhara regional state, Ethiopia. Dessie city has a total of 8 kebeles which are located 410 km away from Addis Ababa, the capital city of Ethiopia. The City is found at a longitude of 38°E and latitude of 11-40°N with an elevation or altitude of 2,470 to 2,550 m above sea level

(m.a.s.l). The area receives an annual rainfall of 1000-1400mm. The average temperature is 15°C-17.5°C [8]. Based on projections from the 2007 Ethiopian national census, the 2018/19 total population of the city was estimated to be 200,000 with an area of 161,828square kilometers. The City livestock population is estimated to be 23,750 cattle, 27,096 sheep, 23,406 goats, 8,231 equines and 125,252 poultry [9].

Study Design and Study Period: A cross-sectional study design was employed to assess the knowledge, attitudes and practices (KAP) on Anthrax and associated risk factors among the community of Dessie City from June 2022 to September 2022.

Study Populations: The study population was animal owners who have lived in randomly selected kebeles of Dessie City. (Gerado, Tita, Kurkur, Boru and Boruselase) as permanent residents for more than six months. Both males and females in the age group above 15 years of age were included in this study. Questions were answered by a single adult from a given household.

Sample Size Determination and Sampling Techniques: In order to generate sufficient information on the knowledge, attitudes and practices regarding anthrax among community members in Dessie City, the sample size was determined by using Slovin Formula for qualitative data of small sample size with a margin error of (e=8%) with an average household size of 1250 livestock in one kebele and a total of 10,000 livestock population size in the study population in the total kebeles [10]. Generally, the sample size of the study can be calculated by using the general formula for small-size qualitative data:




Fig. 1: Map of the study area of Dessie City in the Amhara regional stet

$$n = \frac{N}{(1 + N * e^2)}$$

where

n=sample size

N= the size of the population

e = the margin of error (margin of error=8%)

Therefore the sample size of the study would be

$$n = \frac{10000}{(1 + 10000 * (0.08)^2)} = 154 - 200$$

As a result, 190 respondents were selected as the study population by adding a 10% non-response rate; thus, the total sample size was 209 subjects. From the entire primary sampling unit that is, 8 kebeles (lowest administrative structure); Five were randomly selected using the lottery method. Then, 40 households were selected and interviewed from each kebele using the systematic random sampling method, as there was no significant difference in the number of households. Whenever the selected household was found locked, the next household (on the right side) was substituted automatically for interview. A pretested structured questionnaire consisting of closed-ended questions was used for this study. The data were collected via face-to-face interviews. The questionnaire was first developed in English and then translated into Amharic language (native language) for appropriateness and easiness in approaching the study participants.

Data Management and Analysis: After collecting, the data were cleaned and checked for its completeness. Those incomplete and inconsistent were corrected when possible and removed otherwise. After a complete check-up, the data were coded and entered into a Microsoft Excel sheet and analyzed by using Excel. The frequency distribution of both dependent and independent variables was worked out by using descriptive statics techniques.

Ethical Clearance: The study protocol was reviewed and approved by the Ministry of Agriculture in collaboration with the Food and Agriculture Organization of the United Nations (FAO); Oral informed consents were obtained from each participant after informing them about the purpose of the study as well as the risks, benefits and rights of the study participants. Only voluntary participants were involved in the study. All the information obtained from the study participants was kept confidential.

Data Quality Assurance: Before the beginning of the full study, the pre-test was performed on some participants to see the applicability of the questionnaire. Each questionnaire was checked for incompleteness, missed values and unlikely responses and then manually cleaned upon such indications. The data was cross-checked for consistency and accuracy.

RESULTS

Socio-Demographic Characteristics: Two hundred nine (209) heads of household were interviewed during the study period. Of these, the data collected from nine tin respondents were found to be incomplete and excluded from the analysis. Only data from 190 households were considered for the analysis. The majorities of the respondents were male 124 (65.3%) and were above 15 years of age, of which 28% and 37.9% were between 15 to 30 and 31 to 46 years old, respectively. The majority of the respondents 121(63.6%) were Muslim followed by orthodox 69 (36.3%). Concerning educational status, 58(33.5%) of the participants were illiterate (cannot read and write). Of the participants, 49 (25.7%) and 56 (29.4%) were in primary school and secondary school, respectively. In addition, these 27(14.2%) were in higher education. The majority of the respondents were farmers 97 (51%) (Table 1).

Knowledge of Participants Related to the Cause and Host Range of Anthrax: Of the total of respondents, 105 (55.2%) had awareness of Anthrax. Thirty-nine (20.5%) of respondents knew that Bacteria is the cause of Anthrax. The majority of the participants 137 (72.1%) responded that they did not know the causative agent, however, fewer numbers of respondents 14 (7.3%) were found to have a miss-perception about a causative agent which was the virus.

Table 1: Socio-Demographic information of the study participants.

Socio-Demographic Variables	Frequency	Percentage %
Sex		
Male	124	65.2
Female	66	34.7
Religion		
Muslim	121	63.6
Orthodox	69	36.3
Educational status		
Illiterate(cannot read and write)	58	33.5
Primary school	49	25.7
Secondary school	56	29.4
Higher education	27	14.2
Age		
15-30	64	33.7
31-46	73	38.4
>46	53	27.8




Fig. 2: Knowledge of participants related to cause and host range on Anthrax




Fig. 3: Source of information about Anthrax.

Table 2: Knowledge of participants related to cause and host range on Anthrax

Awareness on anthrax	Frequency	Percent
Yes	105	55.2
No	85	44.7
Cause of Anthrax		
Bacteria	39	20.5
Virus	14	7.3
I don't know	137	72.1

(Table 2). The majority of the respondents 132 (69.4%) knew that Cattle can be affected by the disease (Figure 2).

Out of 190 respondents, 84 (44.2%) were not have the knowledge about Anthrax and 67(35.2%) got the knowledge from veterinary clinic, 9 (4.7%) and 3(1.6%) of the respondents had the awareness from Books\magazines and from social media respectively. However, 14 (7.4%) and 13 (6.8%) of the respondents had the awareness through formal ways (such as radio and television) and informal (such as traditional healer's, neighbors, friends and relatives respectively, as shown in Figure 3.

Knowledge of Participants Related to Modes of Transmission, Clinical Signs and Symptoms and Treatment of Anthrax: As shown in Figure 4, sixty-four (33.7%) respondents didn't know the means of transmission from animal to humans. While 54 (28.4%) and 22(11.5%) of them know that eating raw meat and slaughtering of infected animals transmit the diseases from animal to humans respectively. Sudden death 89 (46.8%) followed by stop eating and drinking 37(19.5%) were described as a major clinical sign of Anthrax in animals by the majority of the respondents, described in Figure 5. Ninety seven (51.1%) of the respondents knew that the disease could not easily treatable. Of the participants, 101 (54.1%) were aware of the fact that Anthrax is a very serious disease in humans and animals (Table 3).

Knowledge of Participants about Human Anthrax: Majority of the respondents 178(93.6%) were not seen human with Anthrax while 12(6.3%) were seen human with Anthrax. The greater number of the respondents 180 (94.7%) were didn't know clinical signs observed in




Fig. 4: Frequencies and percentages of modes of transmission on Anthrax




Fig. 5: Knowledge on the severity of anthrax




Fig. 6: Percentage of clinical sign of anthrax in animals

Table 3: Knowledge of participants related to severity and treatment of Anthrax

Knowledge Related Variables	Frequency	Percentage
Have you ever get training		
Yes	27	14.2
No	163	85.7
How serious a disease in humans and animals		
very serious	101	54.1
somewhat serious	41	21.5
Not very serious	17	8.9
I don't know	31	16.3
Easily treatable after onset of clinical signs		
Yes	93	48.9
No	97	51.1

infected human. While 10(5.2%) know that skin ulcer with black center were described as a major clinical sign of human Anthrax. 167(87.8%) were aware of animal's vaccination as a means of Anthrax prevention. The greater numbers of the respondents 121 (63.6%) were willing to use Anthrax vaccine for their animals (Table 4).

Attitudes of Participants Towards Anthrax: Eighty-eight respondents (46.3%) strongly agree that the consumption of raw meat of an animal is the source of infection for human anthrax. 38(20%) respondents agree that the consumption of raw meat of an animal is the source of infection for human anthrax. Fewer of the respondents




Fig. 7: Attitudes of participants towards Anthrax.

Table 4: Knowledge of participants about Human Anthrax

Knowledge Related Variables	Frequency	Percent
Have you ever seen human with Anthrax		
Yes	12	6.3
No	178	93.7
What kind of signs could be observed in infected human		
Skin ulcer with black center	10	5.2
Don't know	180	94.7
How can a person prevent him from getting Anthrax?		
avoid anthrax infected animal	17	8.9
Burn all suspected anthrax animal carcasses	15	7.9
Bury all suspected anthrax animal carcasses	66	34.7
Vaccinate animals annually	23	12.1
I do not know	101	53.1

Table 5: Attitudes of participants towards Anthrax.

Attitude Related Variables	Frequency	Percent
Grazing pasture is source of infection for animal		
Strongly agree	24	12.6
Agree	60	31.5
Uncertain	99	52.1
Disagree	7	3.7
Anthrax can be prevented through vaccination of animal		
Strongly agree	123	64.7
Agree	52	27.3
Uncertain	10	5.2
Disagree	5	2.6
Anthrax can be controlled through burying of dead animal		
Strongly agree	116	61.1
Agree	66	34.7
Uncertain	6	3.1
Disagree	2	1.1

Table 6. Practices of participants towards Anthrax

Practice Related Variables	Frequency	Percent
Do you use indoor management for your animals		
Yes	121	63.6
No	69	36.3
Have you ever vaccinated your animals?		
Yes	128	67.3
No	66	34.7
In your opinion dose vaccination animals help to prevent anthrax		
Yes	167	87.8
No	23	12.1
Any safety measure taken caring anthrax suspected animal patient?		
Yes	151	79.5
No	39	20.5

20(10.5%) disagree that the consumption of raw meat of animals is the source of infection for human anthrax. 60 (31.5%) respondents agree that grazing pasture is a source of infection for animals. Majority of the respondents 99(52.1%) uncertain that grazing pasture is the source of infection for animals. One hundred twenty-three respondents (64.7%) strongly agree that anthrax can be prevented through the vaccination of animals. Fewer respondents 10(5.2%) were uncertain that vaccination of animals can prevent anthrax. 116(61.1%) respondents strongly agree that anthrax can be controlled through burying of dead animals and 66(34.7%) agree that anthrax can be controlled through burying of dead animal.

Practices of Participants Towards Anthrax: Out of 190 respondents one hundred twenty-one (63.6%) use indoor management for their animals while 69(36.3%) didn't use indoor management for their animals 128 (67.3%) respondents vaccinate their Animals while 66 (34.7%) didn't vaccinate their Animals. Majority of the respondents 167 (87.8%) have knowledge as vaccination of animals help to prevent anthrax. Regarding safety measure 151(79.4%) respondents take any safety measure when they are caring anthrax suspected animal while 39(20.5%) respondents didn't take any safety measure when they are caring anthrax suspected animal. (Table 6).

DISCUSSION

The study was aimed at assessing public knowledge, attitudes and practices towards anthrax in Dessie city. In this study, the total of the respondents (55.2%) had knowledge of Anthrax. This finding is lower than the findings of different studies done by Mesfen *et al.* [11] in South Gonder, Romha *et al.* [12] in Northern Ethiopia and Chacha [13] in Maragua, Kenya, which reported knowledge rates of 71%, 62% and 96.3%, respectively. These differences could be associated with the awareness level of the community, educational status and information access. In this study, (33.7%) and (28.4%) of respondents didn't know the means of transmission of anthrax from animals to humans and eating raw meat transmitted the diseases from animal to humans which is 43.16% and 47% respectively with findings of Dutta *et al.* [14] in selected rural areas of Bangladesh and Few participants mentioned that the disease was caused by microbes/germs [12]. In this study (46.3%) of respondents strongly agree that the consumption of raw meat of an animal is the source of infection for human

anthrax. (20%) of respondents agree that the consumption of raw meat of an animal is the source of infection for human anthrax, this result was consistent with Chacha [13]. About 75.2% of the participants reported that they would not consume meat from cattle found dead, because they were discouraged by veterinary authorities but there were high cases of consumption of meat from an anthrax-related carcass [15].

More than 35% of respondents in this study obtained information from the veterinary clinic (4.7%) and (1.6%) of the respondents had the awareness from Books/magazines and from social media respectively. This finding is higher than the findings of Chacha [13] 21.0% from veterinarians and veterinary Paraprofessionals and 15.1% from the radio.

Sudden death was mentioned as a major clinical sign by the majority of the respondents which is in line with Dutta *et al.* [14]. Stop eating and drinking was described as a second major clinical sign in this study next to sudden death. Of the total (72.1%) of participants responded that they did not know the causative agent while (20.5%) of respondents knew that bacteria was the cause of anthrax. However, fewer numbers of respondents (7.3%) were found to have a miss-perception about a causative agent which was a virus. Consistent with this study, Opere *et al.* [16] showed that most respondents do not know the causes of anthrax but recognize the signs of the disease. Moreover, in the questionnaire survey, the number of respondents who knew the clinical signs was higher than that of respondents who knew the cause of the disease.

Since anthrax is zoonotic and its main transmission to humans is from animals. Animal keeping becomes one of the major risky practice towards anthrax in animals and humans [13]. this study (54%) of the respondents thought that anthrax was a very serious disease; (21.5%) of them thought that anthrax is somewhat serious while only (8.9%) thought that anthrax is not a serious disease, this concurs with a study by Munyua *et al.* [17].

In livestock, anthrax can be prevented largely by vaccination of all grazing animals in the endemic area and implementation of control measures during epizootics vaccination should be done 2-4 weeks before the season when outbreaks may be expected [18].

In this study, More than half of the respondents vaccinate their animals and have knowledge that vaccination of animals help to prevent anthrax. Similar study from Maragua, Kenya and Tigray, vaccination of animals helps to prevent their animals against anthrax;

almost all (98.0%) participants agreed that indeed vaccination helps a few (2.0%) said no, while giving reasons for vaccine failure [13].

Proper and early diagnosis is one of the important components for treatment, prevention and control of anthrax. However, diagnostic facilities were insufficient in the endemic districts, which is similar to the observations from other studies in Asia and Africa [19, 20]. It was revealed that timely diagnosis can control the outbreak of anthrax [21]. It is important to respond quickly to eliminate anthrax-confirmed or suspected carcasses by immediate incineration since spores are rapidly formed and spread by flies and scavengers, especially vultures, that may transmit anthrax over long distances [22]. It is necessary to ensure increased public awareness of vaccination of the livestock population along with sufficient coverage of the anthrax vaccine that will make a large contribution to the control of anthrax outbreaks [14].

CONCLUSION AND RECOMMENDATIONS

This study revealed that there were some gaps in the community concerning knowledge of anthrax in the study area by community members on cause, transmission, signs and control. Due to this, people continue to consume raw and un-inspected meat and fail to present their animals for vaccination. In addition, there is also a lack of knowledge about human with anthrax and clinical signs observed in infected humans. The knowledge among the community members has been enhanced over time by awareness created by veterinarians in the area were the major risk factors of consumption of anthrax-related meat, failure to vaccinate their livestock and poor disposal of carcasses contributed to anthrax transmission. There was a gap between medical and veterinary personnel collaboration in terms of anthrax control given the fact that this disease needs a multidisciplinary approach especially the two professionals for effective control. Educating the community about the health risks of anthrax and the ways of prevention should be given priority by health extension workers, veterinary professionals and the government at large.

Based on the above conclusion the following recommendations are forwarded:

- ✓ Creating awareness for the community on causes, transmission, signs and control anthrax disease.

- ✓ Creating awareness for the city health office to enhance the surveillance system so that anthrax cases could be identified earlier and increase livestock vaccination.
- ✓ With the availability of resources, additional studies should be extended to other regions of Ethiopia so as to compare findings and wholesome intervention measures of the disease.
- ✓ The capacity of veterinary and medical workers should be strengthened in the diagnosis of zoonotic diseases for early outbreak detection and subsequent interventions.
- ✓ Medical and veterinary personnel should be collaborating in a multidisciplinary approach for the prevention and control of anthrax.

REFERENCES

1. Hugh-Jones, M. and J. Blackburn, 2009. The ecology of *Bacillus anthracis*, *Mol. Aspect. Med.*, 30: 356-367.
2. Mohamed, A., G.P. Baaro and S.J. Gelle, 2019. Assessment of knowledge, attitude and practices (KAPS) of anthrax among Pastoralists in Wajir, Isiolo and Marsabit Counties, Kenya, *J. Agric. Sci. Technol.*, 9: 56-63.
3. Inglesby, T.V., D.A. Henderson, J.G. Bartlett, M.S. Ascher, E. Eitzen and A.M. Friedlander, 1999. Anthrax as a biological weapon-medical and public health management. *JAMA*, 281: 1735-1745.
4. Quinn, P.J., 1995. Department, Animal Health and Quarantine Department Annual Report, *Canad Vet. J.*, 1995.
5. WHO (World Health Organization), 2005. Guidelines for the Surveillance and Control of anthrax in humans and animals, *Emerging and Communicable Disease Surveillance, Control*. <http://www.who.int/emc>
6. Shiferaw, G., 2004 Anthrax in Wabessa village in Dessie Zuria District of Ethiopia. *Rev. Sci. Tech. Int. Epiz.*, 23(3): 952-956.
7. Sitali, D.C., M.C. Twambo, M. Chisoni, M.J. Bwalya and M. Munyeme, 2012. Lay perceptions, beliefs and practices linked to the persistence of anthrax outbreaks in cattle in the Western Province of Zambia, *Onderstepoort Journal of Veterinary Research*, 30: 1-8, 2015. Ethiopia, 2012.
8. CSA (Central Statistical Authority), 2007. Population and Housing Census of Ethiopia, 2012.
9. DAOR (Dessie agricultural office report), 2013. Dessie City Agricultural and livestock development office.

10. Adam Anokye, M., 2020. Sample Size Determination in Survey Research. *Journal of Scientific Research and Reports* (June), pp: 90-97.
11. Mesfin, D., D. Mulatu, K. Birara, A. Sibabaw, T. Brhanu and D. Yalew, 2021. Anthrax prevention practice and associated factors among farmers in farta district, South Gonder, Amhara region, North Ethiopia, pp: 2-6.
12. Romha, G., 2020. Knowledge, attitude and Practice towards anthrax in northern Ethiopia a Mixed Approach Study, pp: 6-17.
13. Chacha, N., 2017. Knowledge Attitude Practice Regarding Anthrax among Kommunity members, Health and veterinary Workers in maragua, Kenya.
14. Dutta, P., H. Biswas, J. Ahmed and J. Anita, 2021. Knowledge, attitude and practices (KAP) towards Anthrax among Livestock farmers in selected rural areas of Bangladesh.
15. Chikerema, S., G. Matope and D. Pfukenyi, 2013. Awareness and attitude toward zoonoses with particular reference to anthrax among cattle owners in selected rural communities of Zimbabwe. *Vector-Borne and Zoonotic*.
16. Opare, C., A. Nsiire, B. Awumbilla and B. Akanmori, 2000. Human behavioural factors implicated in outbreaks of human anthrax in the tamale municipality of northern Ghana. *Acta Trop.*, 76(1): 49-52.
17. Munyua, P., A. Bitek, E. Osoro, E. Pieracci, J. Muema, A. Mwatondo, M. Kungu, M. Nanyingi, R. Gharpure, N. Kariuki and M. Thumbi, 2016. Prioritization of Zoonotic Diseases in Kenya. *PLoS ONE*, 11(8): e0161576.
18. Radostits, O.M., C.C. Gay, K.W. Hinchcliff and P.D. Constable, 2007. A textbook of the diseases of cattle, horses, sheep, pigs and goats. *Veterinary Medicine*, 10: 2045-2050.
19. Chakraborty, A., S.U. Khan, M.A. Hasnat, S. Parveen, M.S. Islam, A. Mikolon, R.K. Chakraborty, B.N. Ahmed, K. Ara, N. Haider, S.R. Zaki, A.R. Hoffmaster, M. Rahman, S.P. Luby and M.J. Hossain, 2012. Anthrax outbreaks in Bangladesh, 2009-2010. *The American Journal of Tropical Medicine and Hygiene*, 86(4): 703-710.
20. Vaismoradi, M., H. Turunen and T. Bondas, 2013. Content analysis and thematic analysis: Implications for conducting a qualitative descriptive study. *Nursing and Health Sciences*, 15(3): 398-405.
21. Sahoo, K.C., S. Negi, D. Barla, G. Badaik, S. Sahoo, M. Bal, A.K. Padhi, S. Pati and D. Bhattacharya, 2020. The Landscape of anthrax prevention and control: Stakeholders' perceptive in Odisha, India. *International Journal of Environmental Research and Public Health*, 17(9): 3094.
22. Turnbull, P., 2001. Guidelines for the surveillance and control of anthrax in human and animals 3rd Ed Geneva, World Health Organization.