

## Extrinsic Factors Influencing Antibacterial Activities of *Tapinanthus bangwensis* Against Diarrheal Causing Organisms

<sup>1</sup>B.T. Thomas, <sup>2</sup>H.I. Effedua, <sup>5</sup>G.C. Agu, <sup>3</sup>P.A. Akinduti,  
<sup>4</sup>O. Ejilude, <sup>5</sup>M.O. Efuntoye, <sup>6</sup>A.E. Ayodele and <sup>1</sup>A. Oluwadun

<sup>1</sup>Department of Medical Microbiology and Parasitology,  
Olabisi Onabanjo University, Ago-Iwoye, Ogun State, Nigeria

<sup>2</sup>Department of Medical Laboratory Science, School of Public Health, Babcock,  
University, Ilishan-Remo, Ogun State, Nigeria

<sup>3</sup>Department of Veterinary Microbiology and Parasitology,  
University of Agriculture, Abeokuta, Ogun State, Nigeria

<sup>4</sup>Department of Microbiology and Parasitology,  
Sacred Heart Hospital, Lantoro, Abeokuta, Ogun State, Nigeria

<sup>5</sup>Department of Microbiology, Olabisi Onabanjo University, Ago-Iwoye, Ogun State, Nigeria

<sup>6</sup>Department of Microbiology and Botany, University of Ibadan, Oyo State, Nigeria

**Abstract:** *Tapinanthus bangwensis* is a parasitic plant with wide distribution and documented antimicrobial efficacies. Inconsistence in its activities has however, been suggestively linked with some factors. Hence, this study was carried out to establish the effect of some extrinsic factors on the antibacterial activities of *T. bangwensis* against some diarrheal causing bacteria. Antibacterial activity of *T. bangwensis*, collected from some parts of Nigeria was determined by standard agar-diffusion method. Results from this study showed a significantly higher zone of bacterial inhibition with chloroform extract when compared with ethanol and aqueous extracts ( $P < 0.05$ ). Higher antibacterial activities was also observed with extracts obtained from air dried plants than those obtained from sun dried and oven dried plants ( $P < 0.05$ ). Steaming method of extraction produced a significantly higher zone of bacterial inhibition than cold and hot methods of extraction ( $P < 0.05$ ). Percentage weight yield of active crude compounds of *T. bangwensis* was highest in chloroform ( $4.63 \pm 1.99\%$ ) than methanol ( $2.83 \pm 2.06\%$ ) and water ( $2.28 \pm 1.90\%$ ) ( $P < 0.05$ ). A significant positive correlation ( $r = +0.91$ ,  $P < 0.05$ ) was observed between percentage weight yield and zone of antibacterial inhibition, exhibited by *T. bangwensis*. Weight yield accounted for 70% of antibacterial activities with a linear relationship of  $y = 3x + 5.6$ . Antibacterial activities of *T. bangwensis* were however, not affected by its host plants and varying concentrations of its crude extracts ( $P > 0.05$ ). Conclusion from this study has shown that solvents and method of extraction (as mode of plant drying and means of concentrating extracts) are important and influential extrinsic factors that determines the antibacterial activities of *T. bangwensis*.

**Key words:** *Tapinanthus bangwensis* • Extrinsic factors • Diarrheal organisms

### INTRODUCTION

*Tapinanthus bangwensis* (order Santalales) belong to the family of Loranthaceae and constitute the largest group of parasitic plants which has about 950 species distributed in 77 genera [1]. Loranthaceae:- including *Tapinanthus* constitute a great deal of pestilence in the natural forests, plantations, cultivated fruit trees and

ornamental plants; causing damages to the host plants [2, 3]. *Tapinanthus bangwensis* is widely distributed in Nigeria. The plant exists as xerophytes on many host trees. Its leaves and young twigs have in the past been used in folklore treatment of diseases and sterility in cow. Documentations have been made about the antimicrobial efficacy of *T. bangwensis* [4] but its activity tends to vary among investigators. Also, there

have been controversies over the hypothesis that the antimicrobial efficacies of *T. bangwensis* is host plant dependent. Hence, we have decided to investigate the impact of host plant factor and other extrinsic factors on the antibacterial activities of *T. bangwensis* against some bacterial pathogens causing diarrhea.

## MATERIALS AND METHODS

**Collection of Plants:** Fresh leaves and twigs of *Tapinanthus bangwensis* were collected from Iseyin, Southern part of Oyo State and Chagas village in Abuja, Nigeria. The plants were authenticated by Dr A.E. Ayodele at the department of Botany and Microbiology, University of Ibadan, Nigeria and were designated T1 – T4 on the basis of the host plants from which they were collected. T1 was from *Triclisia gillettii* (De wild) stainer, T2 from *Parkia biglobosa* (Jacq.) Benth, T3 from *Citrus aurantifolia* (Christm.) swingle and T4 from *phyllanthus muellerianus* (Oktze).

**Test Microorganisms:** Authentic pure cultures of human pathogenic bacteria like *Escherichia coli*, *Proteus mirabilis*, *Salmonella typhi*, *Shigella sonnei*, *Shigella dysenteriae*, *Shigella flexneri* *Staphylococcus aureus*, *Bacillus cereus*, *Pseudomonas aeruginosa* and *Yersinia enterocolitica*, were obtained from Nigeria institute of Medical Research (NIMR) Lagos, Nigeria. These isolates were further confirmed using standard microbiological techniques [4].

**Antibacterial Activity of Plant Extracts:** Susceptibility of diarrheal causing bacterial isolates to the extracts was determined by standard disc diffusion assay [14]. Well dried Mueller Hinton Agar plates were seeded with 24 hour old culture of bacterial strains. The inoculum size was adjusted to achieve a final concentration of  $10^5$ cfu/ml after dilution from  $10^8$ cfu/ml (equivalent of 0.5 McFarland standards). The sterile Whatman filter paper discs (5mm in diameter) impregnated with plants extract (50,100,150 and 200mg/ml) were placed on the surface of the culture plates and incubated at 37°C for 24 hrs and diameter of zone of inhibition were measured in mm. Discs with chloroform, methanol and sterile distilled water were used as control.

**Preparation of Plant Extracts:** Fresh Leaves were divided into three groups. Group one were air-dried, Group two were sun-dried and Group three were oven-dried. After drying, the leaves were shredded and separately

preserved in air-tight cellophane bags. The shredded leaves were milled into powder. Cold extraction of the plant was made by soaking 100g of powdered plant into 400ml each of methanol, chloroform and water respectively in flasks. The flasks were manually agitated at intervals for 5 days. All extracts were then filtered with whatman no.2 [11]. The filtrates were later concentrated to dryness with the aid of a rotary evaporator. The steam extraction was carried out as described by Adeolu and Oladimeji [5]. The hot extraction involves placing the leaves and twigs of *T. bangwensis* in a pot and boiling for 30 minutes. This was then allowed to cool and the leaves were then squeezed to obtain the extract that was later concentrated by rotary evaporator [5]. The yield of concentrates from the various extracts was then calculated using the following formula [12]:

$$\text{Percentage weight yield (\%)} = W_2 / W_1 \times 100.$$

Where

$W_1$  = Weight of herbal powder before extraction and;

$W_2$  = Weight of concentrate after extraction.

**Statistical Analysis:** All data were analyzed by SPSS package version 15. Comparison of mean zones of inhibition were determined by ANOVA (Analysis of Variance) while regression analysis was used to determine the linearity between percentage weight yields (%) of plant and their proportional antibacterial activities.

## RESULTS

The effect of host plants on the antibacterial activities of *T. bangwensis* was determined in Table 1. Although, the zones of inhibition (mm) varied with different host plants, but the difference was insignificant ( $F=1.06$ ,  $P>0.05$ ). In Table 2, when the mean of zones of inhibition of *T. bangwensis* concentrates from different solvents were compared, a significant difference was observed ( $F=10.13$ ,  $P<0.05$ ); with highest antibacterial activity being recorded with chloroform. Table 3 displays various means of zones of inhibition of *T. bangwensis* extracts processed by different methods of drying. Analyzed data showed that extracts prepared from air-dried plants of *T. bangwensis* had the highest zone of inhibition of  $13.50 \pm 3.50$ mm ( $F= 6.47$ ,  $P<0.05$ ). Comparison between antibacterial activities among concentrates of *T. bangwensis* yielded by different extraction techniques was made in Table 4.

Table 1: Effect of host plant on the antibacterial activity of *Tapinanthus bangwensis*

Host plants	Zones of inhibition (mm)	
	No	Mean±SEM
<i>Triclisia gillettii</i>	10	8.43±1.65
<i>Parkia biglobosa</i>	10	6.45±1.94
<i>Citrus aurantifolia</i>	10	8.34±2.63
<i>Phyllanthus muellerianus</i>	10	6.42±2.03

F = 1.06, P > 0.05.

Table 2: Effect of solvent of extraction on the antibacterial activity of *Tapinanthus bangwensis*

Solvents of extraction	Zones of inhibition(mm)	
	No	Mean±SEM
Methanol	10	15.64±1.20
Chloroform	10	19.00±1.50
Water	10	11.20±0.85

F = 10.13, P < 0.05.

Table 3: Effect of drying method on the antibacterial activity of *Tapinanthus bangwensis*

Drying methods	Zones of inhibition(mm)	
	No	Mean±SEM
Air drying	10	13.50±3.50
Sun drying	10	8.88±2.03
Oven drying	10	9.00±1.63

F = 6.47, P < 0.05.

Table 4: Effect of methods of extraction on the antibacterial activity of *Tapinanthus bangwensis*

Methods of extraction	Zones of inhibition(mm)	
	No	Mean±SEM
Steam extraction	10	13.13±1.59
Cold extraction	10	9.58±0.68
Hot extraction	10	7.42±0.85

F = 6.66, P < 0.05.

Table 5: Effect of concentrations of chloroform extracts on the antibacterial activity of *Tapinanthus bangwensis*

Concentrations	Zones of inhibition(mm)	
	No	Mean±SEM
100mg/ml	10	12.05±2.06
150mg/ml	10	14.30±1.99
200mg/ml	10	16.00±1.90
50mg/ml	10	19.50±2.45

F = 2.23, P > 0.05.

Table 6: Effect of type of solvents on weight yield of the active ingredients of *Tapinanthus bangwensis*

Solvents	Weight yield (%)	
	No	Mean±SEM
Methanol	3	2.83±2.06
Chloroform	3	4.63±1.99
Water	3	2.28±1.90

F = 448.17, P < 0.05.




Fig. 1: Relationship between inhibitory activities and percentage weight yield of active chloroform extract of *T. bangwensis*.

The significantly highest zone of inhibition was observed with steam (13.13±1.59mm) when compared with cold (9.58±0.68) and hot (7.42±0.85) methods of extraction (F=6.66, P<0.05). When means of zone of inhibition of *T. bangwensis* was compared at different concentrations that ranged from 50–200mg/ml, no significant difference was observed (F=2.23, P>0.05) in Table 5. Comparison of percentage weight yields (%) of crude *T. bangwensis* concentrates regarding the type of solvents of extraction demonstrated that highest yield of 4.63±1.99% was with chloroform (F = 448.17, P<0.05) (Table 6). A significant correlation (r = +0.91, P<0.05) was observed between the percentage weight yield of *T. bangwensis* and its antibacterial activities, with a linear equation:  $y = 3x + 5.60$ . Regression determinant showed that percentage weight yield was accountable for 70% of the antibacterial activities (Figure 1).

## DISCUSSION

Results of this study have shown that the antibacterial activities of *T. bangwensis* were independent of the host plant factor. This observation oppose

the hypothesis which says that the host plant influence the antibacterial activities of *T. bangwensis* [4]. The significant highest percentage yield of crude concentrate of *T. bangwensis* by chloroform in comparison with other solvents demonstrated high extraction strength with chloroform ( $P < 0.05$ ). Also, the highest antibacterial activity was found in the crude chloroform extract which strongly suggests that the active antibacterial compound might be organic and less polar in origin [11].

After processing of *T. bangwensis* leaves by different methods of drying, the highest antibacterial activity was observed in air dried extract while less activity was observed with oven-dried extract and least activity was observed with sun dried extract ( $P < 0.05$ ). Lesser antibacterial activity in sun dried extract may be due to photochemical degradation of the active compounds which in turn may result in structural modification of functional groups required for active antimicrobial activity [6]. Also, lesser antibacterial activity in oven-dried extract indicated that the active antibacterial compound of *T. bangwensis* might be heat-labile. This observation was similar to those of Olaniyi *et al.* [7], Harnischfeger [8] and Niggermann and Gruber [9]. Furthermore, the lesser antimicrobial activities in oven-dried extract might also be linked with loss of volatile contents of *T. bangwensis* such as phenols and essential oils [10]. Steam method of extraction of *T. bangwensis* has been shown to produce bioactive compound with better antibacterial activities than cold and hot methods of extraction ( $P < 0.05$ ). The weakest antibacterial activity in concentrate of *T. bangwensis* derived from hot method of extraction further re-emphasize the volatile nature of active antibacterial constituents of *T. bangwensis* while lower antibacterial activity in cold method of extraction is a reflection of the decreased solubility of active plant constituent at lower temperature [11]. Comparison of the antibacterial activity of crude chloroform extract of *T. bangwensis* at concentrations such as 50, 100, 150 and 200mg/ml showed no significant difference ( $P > 0.05$ ). This indicated that *T. bangwensis* exhibits antibacterial activities almost at equal level at concentration range of 50-200mg/ml. Equilibrium of antibacterial activities between the lowest and highest concentrations of *T. bangwensis* clearly showed that the antibacterial quality of the active constituent of the plant is largely dependent on its molecular weight and diffusion rates through agar rather than its concentration [13]. Since higher zones of inhibition in agar diffusion

antibiotic susceptibility tests is an attribute of faster rates of drug diffusion and low molecular weight, it can then be inferred that the active antibacterial constituents of *T. bangwensis* might be among compounds with lower molecular weight.

In conclusion, the outcome of this study has demonstrated that antibacterial activities of concentrates from *T. bangwensis* could be best enhanced by methods that include air-drying and chloroform extraction of the plant via steaming.

## REFERENCES

1. Engone, O.N.L. and G. Salle, 2006. Faut il eradiquer hramanthera capitata, parasite des heveas en Afrique? C.R. Biologies, 329: 185-195.
2. Sonke, B., D. Kenfaek and M. Tindo, 2000. Parasitisme de lavocatier (*Persea Americana*, lauraceae) par les loranthaceae dans la region de Yaounde (Cameroun). Fruits, 55: 325-331.
3. Boussim, I.J., S. Guinko, C. Tuquet and G. Salle, 2004. Mistletoes of the parklands of Burkina-Faso. Agro-forest. Syst., 60: 39-49.
4. Efuntoye, M.O., A.E. Ayodele, B.T. Thomas and T.O. Ajayi, 2010. Does host plant affect the antibacterial activity of *Tapinanthus bangwensis* (Engl. and Krause) Danser (Loranthaceae)? J. Medicinal Plants Res., 4(13): 1281-1284.
5. Adebolu, T.T. and S.A. Oladimeji, 2005. Antimicrobial activity of leaf extracts of *Ocimum gratissimum* on selected diarrhea causing bacteria in South Western Nigeria. Afr. J. Biotech., 4(7): 682-684.
6. Adegbohun, O.M. and O.O. Olukemi, 2010. Effect of light irradiation on the antimicrobial activity of *Zanthoxyloides* (Iam) methanolic extract. Afr. J. Pharm. Pharmacol., 4(4): 145-150.
7. Olaniyi, A.A., J.S.K. Ayun, A.O. Ogundaini and T.A. Olugade, 2000. Essential inorganic and organic pharmaceutical chemistry. Shaneson C.I. Ltd., Ibadan, pp: 487-512.
8. Harnischfeger, G., 2005. Development, quality assurance and regulatory aspects of herbal medicinal products. UNIDO-ICS Training course, Panama.
9. Niggermann, B. and C. Gruber, 2003. Side effects complementary and alternative medicine. Allergy, 58: 707-716.
10. Rajendran, A., V. Narayanan and I. Gnanavel, 2007. Photochemical and Electrochemical stabilities of *Aloevera* sap. J. App. Sci. Res., 3(12): 1871-1878.

11. Marjorie, M.C., 1999. Plant products as antimicrobial agents. *Clin. Microbiol. Rev.*, 12(4): 564-582.
12. Oyi, A.R., M.A. Ibrahim and Y.J. Oyeniya, 2011. Physicochemical Properties of Ethyl Cellulose Derived From Sawdust. *Nig. J. Pharm. Sci.*, 10(1): 81-90.
13. Agbolahor, D.E., K.C. Uzoma, K.A.U. Nwobu, S.O. Adedeji and R.A. Adegbola, 1996. Questions and Models Answers in Medical Bacteriology. Ayo Abiodun Nig. Enterprises, Ibadan, Nigeria, pp: 95.
14. Riess, J.L., M.C. Recio and A. Vilar, 1998. Screening methods for natural products with antimicrobial activity: a review of the literature. *J. Ethnopharmacol.*, 23: 127-149.