

## Studies on Microbial Population Density and Soil Respiration in Cropland Soil and Middens of Earthworm *Linnogaster pusillus*, Stephenson

Suruchi Kumari and M.P. Sinha

Department of Zoology, Ranchi University, Ranchi-834 008, Jharkhand, India

**Abstract:** The present paper dealt with the relationship of the microbial population and the CO<sub>2</sub> evolution in soil and midden of earthworm *Linnogaster pusillus*, Stephenson. A trend of decrease in bacterial population and CO<sub>2</sub> evolution was recorded up to 42 days of observation in both the samples. The initial bacterial population (number/g soil) was found to be  $36.3 \pm 1.504 \times 10^9$  and  $51.1 \pm 1.350 \times 10^9$  in soil and midden respectively and thereafter a declining pattern was noticed. Maximum value of CO<sub>2</sub> evolution (mg CO<sub>2</sub>/kg soil/h) was  $6.23 \pm 0.98$  and  $8.20 \pm 1.72$  observed in soil and midden respectively. A positive significant correlation was observed between the two in both soil and midden of earthworms with  $r^2=0.9524$  ( $p \leq 0.001$ ) and  $0.9856$  ( $p \leq 0.01$ ) respectively. The CO<sub>2</sub> release and bacterial population was comparatively higher in midden of earthworm than in cropland soil.

**Key words:** *Linnogaster pusillus* • Midden • CO<sub>2</sub> Evolution • Microbial Activity

### INTRODUCTION

Soil biotas are thought to harbor a large part of the world biodiversity that include organisms from microscopic microflora to micro fauna. The microbes are the important element of soil environment as they participate in the degradation of organic matter and make the nutrient available to other soil organisms. Soil microbes are responsible for sequestration of green house gases especially methane. They oxidize atmospheric methane and are responsible for an estimated 5-10% of its total removal from atmosphere [1]. In agricultural soils the earthworms are recognized as soil ecosystem engineers. They form micro-habitats where a large number of physical, chemical and microbial changes occur because the structures are the outcome of an intestinal transit in earthworm cast and midden [2]. Protection of the soil habitat is the first step towards sustainable management of its biological properties that determine long-term quality and productivity.

Soil metabolism refers to the overall activities of the soil organisms involving biochemical process of their metabolic activity which is computed by the CO<sub>2</sub> evolution and enzyme activity of the soil [3]. Microorganisms are the main source of enzymes in soil [4]. The best index of overall metabolic activity of soil

microbial population is CO<sub>2</sub> evolution or soil respiration, which can be rapidly determined. The measurement of soil CO<sub>2</sub> respiration is a means to gauge biological soil fertility [5]. Soil respiration is an important aspect of soil quality and an indication of soil fertility [6]. Therefore, soil metabolism may be depicted by microbial activity and CO<sub>2</sub> evolution. Earthworms are the natural soil engineers, which serve as bio catalytic agent to enhance the soil fertility through physical, chemical and biological process. Structure known as earthworm midden formed at the soil surface by the feeding and casting activities of earthworm, contributes significantly to soil heterogeneity. Midden acts as a focus for soil micro fauna and encourages microbial activity [7]. In this study, experiments were carried out to relate indices of bacterial population and CO<sub>2</sub> evolution in cropland soil and midden of earthworm *Linnogaster pusillus*, Stephenson in order to assess the utilization of earthworms in agricultural prospect.

### MATERIALS AND METHODS

**Soil Collection and Bacterial Isolation:** Soil was collected from agro-ecosystem of Ranchi, Jharkhand and was kept in two separate plastic containers under moist condition. One container was normal soil without earthworm and

other was culture of the earthworms *Linnogaster pusillus*, Stephenson in plastic trays. The middens were collected from the plastic trays and used for microbial study. Bacterial cultures were done from both soil and midden samples by dilution plate method [8]. The isolation of bacteria from soil samples was initiated by taking 1g of soil from both groups and was diluted with 9 mL of sterilized deionized water till  $10^{-7}$  dilution. One mL inoculums of the primary suspension was taken for bacterial culture in a Petri plate (diameter = 100mm) containing Czapek Dox agar media, (peptone - 10g/L, beef extract - 10g/L, agar - 15g/L NaCl- 5g/L, pH- 7.2) and inoculated at 37°C for 48 h [9]. After that colony counts were continued at every interval of 7 days till 42<sup>nd</sup> day.

**Estimation of CO<sub>2</sub> Evolution:** CO<sub>2</sub> evolution was measured by alkali absorption technique [10]. CO<sub>2</sub> evolution was expressed as mg CO<sub>2</sub>/kg soil/h.

**Statistical Analysis:** The data were subjected to the analysis of variance (ANOVA). Correlation and linear regression analysis were also done to assess the strength or weakness of the relationship between the microbial population and CO<sub>2</sub> evolved or soil respiration in the cropland soil and midden of earthworm *Linnogaster pusillus*.

## RESULTS AND DISCUSSION

Bacterial population in soil in the beginning was  $36.3 \pm 1.504 \times 10^9$  and in midden was  $51.1 \pm 1.35 \times 10^9$ . Thereafter a sharp decline in bacterial population was observed. The change in population was found to be significant ( $p \leq 0.001$ ). In soil, bacterial population gradually decreased to  $32.8 \pm 2.80 \times 10^9$ ,  $24.5 \pm 2.44 \times 10^9$ ,  $21.4 \pm 2.55 \times 10^9$ ,  $18.9 \pm 1.66 \times 10^9$ ,  $13.3 \pm 1.171 \times 10^9$  and  $7.9 \pm 0.907 \times 10^9$  on 7<sup>th</sup>, 14<sup>th</sup>, 21<sup>st</sup>, 28<sup>th</sup>, 35<sup>th</sup> and 42<sup>nd</sup> day respectively. On 7<sup>th</sup> day of observation bacterial population in midden was  $44.9 \pm 2.41 \times 10^9$  which was decreased to  $12.7 \pm 0.750 \times 10^9$  (Fig. 1). The percentage

decrease in bacterial population over initial population in cropland soil was recorded as 9.64, 32.50, 41.04, 47.93, 63.36 and 78.23% on 7<sup>th</sup>, 14<sup>th</sup>, 21<sup>st</sup>, 28<sup>th</sup>, 35<sup>th</sup> and 42<sup>nd</sup> day of observation. While the initial percentage decrease in bacterial population of midden was 12.13%, which was more pronounced on last day of observation i.e. 75.14% over the initial population (Table 1). Percentage change in bacterial population in between soil and midden was 40.77% which has been gradually increased till up to the 60.78% on the 42<sup>nd</sup> day that was more pronounced difference in bacterial population in soil and midden in which bacterial population was always higher in midden. Bacterial population has been reported higher in midden compared to the standardized soils ingested by the earthworm [11].

Soil respiration and bacterial population were gradually decreased. Soil respiration or CO<sub>2</sub> evolution (mg CO<sub>2</sub>/kg soil/h) in soil and midden were maximum at 1<sup>st</sup> day  $6.23 \pm 0.98$  and  $8.20 \pm 1.72$  respectively. CO<sub>2</sub> evolution rate in midden gradually decreased from  $7.92 \pm 0.99$  to  $3.62 \pm 0.31$  on 42<sup>nd</sup> day of experiment, due to the lower moisture and nutrient content. Bacterial population showed a highly significant correlation ( $r^2 = 0.9524$ ,  $p \leq 0.001$ ) with CO<sub>2</sub> evolution in soil with incremented value from  $6.23 \pm 0.98$  to  $3.02 \pm 0.22$  mg CO<sub>2</sub>/kg soil/h (Fig. 2). The two parameters were highly correlated in midden (Fig. 3), with  $r^2 = 0.9856$ . The effective activity was observed with higher bacterial population and CO<sub>2</sub> evolution in midden of earthworm.

In the present investigation a significant positive correlation ( $r^2 = 0.9856$ ,  $p < 0.01$ ) of bacterial population and CO<sub>2</sub> evolution was found. The CO<sub>2</sub> evolution showed significant dependency on the population count as represented by the equation  $y = 0.67826 + 0.164166x$  (Fig. 2). One way ANOVA showed that bacterial population plays a significant role in CO<sub>2</sub> evolution in the midden of earthworm ( $F = 9.92$ ,  $df = 6$ ,  $p \leq 0.001$ ). Respiration is probably the process that is most closely associated with life [12]. Soil respiration is attributed to a wide range of microorganisms. Soil fauna also makes a

Table 1: Percentage change between the bacterial populations of cropland soil and midden of earthworm *Linnogaster pusillus*

Days of observation	Non ingested soil	Earthworm midden	% change
0	$36.3 \pm 1.504 \times 10^9$	$51.1 \pm 1.350 \times 10^9$	+ 40.77 %
7	$32.8 \pm 2.809 \times 10^9$ (-9.64)	$44.9 \pm 2.417 \times 10^9$ (-12.13)	+36.89%
14	$24.5 \pm 2.441 \times 10^9$ (-32.50)	$27.8 \pm 1.02 \times 10^9$ (-36.39)	+33.19%
21	$21.4 \pm 2.553 \times 10^9$ (-41.04)	$31.1 \pm 0.650 \times 10^9$ (-45.59)	+29.90%
28	$18.9 \pm 1.665 \times 10^9$ (-47.93)	$26.3 \pm 0.80 \times 10^9$ (-47.94)	+40.7%
35	$13.3 \pm 1.171 \times 10^9$ (-63.36)	$24.3 \pm 0.750 \times 10^9$ (-61.64)	+47.36%
42	$7.9 \pm 0.907 \times 10^9$ (-78.23)	$12.7 \pm 0.750 \times 10^9$ (-75.14)	+60.78%

Values in parenthesis are percentage increase (+) or decrease (-) over initial value




Fig. 1: Bacterial population in soil and midden over a period of 42 days

\* represent the significant difference at 1% level; n=3




Fig. 2: Linear regression between bacterial population and CO<sub>2</sub> evolution in cropland soil




Fig. 3: Linear regression between bacterial population and CO<sub>2</sub> evolution in midden of earthworm *Linnogaster pusillus*

significant contribution (about 10%). But plant roots also contribute between 12 and 30% to the total release of CO<sub>2</sub> through respiration in field [12]. Hence, field based methods give the total respiration of all organisms, whereas laboratory methods give only the microbial respiration. Measurement of CO<sub>2</sub> evolution was taken as

an index of the metabolic activities of soil organisms and it helps in assessing the organic input into the system. The energy flow and rate of mineralization comparatively the parameters of soil metabolism were incremented by the incorporation of earthworm than in their absence [13-15].

The enhancement in the population with earthworm is due to the ameliorating effect of the earthworm. Earthworm activity may raise N<sub>2</sub>O emissions from agro ecosystems. Rather than emitting N<sub>2</sub>O themselves, earthworms are thought to enhance soil microbial activity (nitrification, denitrification) by changing physiochemical properties, excreting mucus and increasing carbon available by enhancing microbial activities earthworm's aid in nutrient cycling processes and in soil structure development [16]. They have been considered beneficial animals which can significantly influence soil structure and soil fertility [17].

CO<sub>2</sub> evolution was higher in midden than soils where there were significant variations in CO<sub>2</sub> evolution due to temperature, moisture and nutrients. There was a difference in response of microbial respiration between soil and midden. Specific respiration in midden was significantly decreased with age, which is in agreement with the finding of Tiunov and Scheu [18]. In this experiments, significant positive correlation ( $r^2 = 0.9856$ ) for bacterial population and CO<sub>2</sub> evolution was found. And also significant positive correlation between bacterial population and CO<sub>2</sub> evolution has been reported by many workers [19, 20].

The microbial community of soil may be profoundly affected by the presence of earthworms. Studies of various earthworm species have generally shown an increase in microbial activity [11, 21-28]. In contrast other studies have found earthworm reduced the microbial activity were of the opinion that selective feeding of earthworms on materials with higher microbial counts accounted for the population increase in midden produced [29-36]. Earthworm strongly enhanced microbial activity measured as CO<sub>2</sub> production in midden, which indicates that there are hot spots for soil microbial dynamics and increasing habitat heterogeneity for soil microorganism and increase soil fertility which is important for sustainable development [37]. The earthworms can specifically affect soil fertility that may be of great importance to increase sustainable land use in naturally degraded ecosystems as well as agro ecosystems. Proper earthworm management may sustain crop yields whilst fertilizer inputs could be reduced. Moreover, our data strongly support the fact that the impact of this earthworm species *Linnogaster pusillus*, Stephenson, in the soil is restricted to their middens and increasing soil heterogeneity.

## ACKNOWLEDGEMENT

The first author (S. Kumari) thankfully acknowledge DST Govt. of India for financial support as INSPIRE fellowship and HOD Zoology for providing facilities.

## REFERENCES

1. Cicerone, R.I. and R.S. Oremland, 1988. Biogeochemical aspects of atmospheric methane. *Global Biogeochemical Cycles*, 2: 299-327.
2. Jones, C.G., J.H. Lawton and M. Shachak, 1994. *Organisms as Ecosystem Engineers*. *Oikos*, 69(3): 373-386.
3. Serdar, B., 2010. Effect of cement dust pollution on microbial properties and enzyme activities in cultivated and no-till. *Soils African J. Microbiology Research*, 4(22): 2418-2425.
4. Wong, M.H. and W.C. Wong, 1986. Effect of fly ash on soil microbial activity. *Environ. Pollut. Ser. A.*, 40: 127-144.
5. Hanley, R.L., W.H. Brinton. and E. Evans, 2008. Estimating soil carbon, nitrogen and phosphorus mineralization from short-term carbon dioxide respiration. *Communications in Soil Science and Plant Analysis.*, 39: 2706-2720.
6. Staben, M.L., D.F. Bezdicsek, J.L. Smith and M.F. Fauci, 1997. Assessment of soil quality in conservation reserve program and wheat-fallow soils. *Soil Sci. Soc. Am. J.*, 61: 124-130.
7. Hale, C., L. E. Frelich, P.B. Reich and J. Pastor, 2005. Effects of European earthworm invasion on soil characteristics in northern hardwood forests of Minnesota, USA. *Ecosystem.*, 8: 911-927.
8. Parkinson, D., T.R.G. Gray and S.T. Williams, 1971. *Methods to study ecology of soil microorganisms*. IBP Handbook No. 19, Blackwell Scientific Publ. Oxford, pp: 116.
9. Thom, C. and K.B. Raper, 1945. *A Manual of the Aspergilli*. Baltimore, MD, Williams and Wilkins Co., pp: 846.
10. Witkamp, M., 1966. Rate of carbon dioxide evolution from the forest floor. *Ecology*, 47: 492-494.
11. Daniel, O. and J.M. Anderson, 1992. Microbial biomass and activity in contrasting soil materials after passage through the gut of the earthworm *Lumbricus rubellus* Hoffmeister. *Soil Biology and Biochemistry*, 24(5): 465-470.

12. Bloem, J., D.W. Hopkins and A. Benedetti, 2006. Microbiological methods for assessing soil quality. Wallingford, UK: CABI Publishing, pp: 307.
13. Macfadyen, A., 1970. Soil metabolism in relation to ecosystem energy flow and to primary and secondary production. In: Methods of study in soil ecology. Ed., Phillipson, J. Publisher, pp: 167-172.
14. Wanner, H., 1970. Soil respiration, litter fall and productivity of tropical Rain Forest. J. Ecol., 58: 543-547.
15. Witkamp, M., 1971. Soils as components of ecosystem. Annual review of Ecology and Systematics, 2: 85-109.
16. Lubbers, I.M., Lucas Nebert, Georgios Giannopoulos, Lijbert Brussaard and Jan Willem Van Groeningen, 2010. Interactions between earthworm ecological strategy and microbial activity affect N<sub>2</sub>O dynamics, COS, Wageningen University, pp: 93-5.
17. Lee, K.E., 1985. Earthworms. Their ecology and relationships with soils and land use. Academic press. Australia, pp: 411.
18. Tiunov, A.V. and S. Sechu, 2000. Microbial biomass, biovolume and respiration in *Lumbricus terrestris* L. cast material of different age. Soil Biol. Biochem., 28(4): 265-275.
19. Dkhar, M.S. and R.R. Mishra, 1987. Microbial population, fungal biomass and CO<sub>2</sub> evolution in maize (*Zea mays* L.) field soils, Plant and Soil, 99(2-3): 277-283.
20. Pandey, R.R., G. Sharma, T.B. Singh and S.K. Tripathi, 2010. Factors influencing soil CO<sub>2</sub> efflux in northeastern Indian oak forest and plantation. African J. Plant Sci., 4(8): 280-289.
21. Parle, J.N., 1963. A microbiological study of earthworm casts. J. Gen., 31: 13-22.
22. Burtelow, A., P.J. Bohlenand and P.M. Groffman, 1998. Influence of exotic earthworm in vasion on soil organic matter, microbial biomass and denitrification potential in forest soils of the northeastern United States. Appl. Soil Ecol., 9: 197-202.
23. Bohlen, P.J., R.W. Parmelee, M.F. Allen and Q.M. Ketterings, 1999. Differential effects of earthworm on nitrogen cycling from various nitrogen 15-labeled substrates. Soil Sci. Soc. Am. J., 63: 882-890.
24. Kumari, S., S. Jabeen, B.S. Raipat and M.P. Sinha, 2009. Impact of aging of earthworm middens on density, dynamism and biomass of microbial population. The Bioscan., 4(3): 535-538.
25. Kumari, S., N. Yadav, S. Kumari, P. Saha, S. Jabeen, B.S. Raipat and M.P. Sinha, 2010. Ecobiological and bioinformatic study of soil bacterium from agro-ecosystem. The Bioscan: Special Issue, 1: 95-104.
26. Bohlen, P.J. and C.A. Edwards, 1995. Earthworm effects on N dynamics and soil respiration in microcosms receiving organic and inorganic nutrients. Soil Biology and Biochemistry, 27(3): 341-348.
27. Saha, P., S. Kumari, B.S. Raipat and M.P. Sinha, 2011. Phylogenetic analysis of tolerant bacteria from *Parthenium hysterophorous* (L.) amended soil by bootstrap approach. Inter. J. Microbiol. Res., 2(2): 176-183.
28. Kumari, S., P. Saha and M.P. Sinha, 2012. 16 S rDNA based genomic analysis of a bacterium isolated from earthworm (*Linogater pusillus*, Stephenson) midden. World Applied Sci. J., 16(3): 427-432.
29. Devliegher, W. and W. Verstraete, 1995. *Lumbricus terrestris* in asoil core experiment: nutrient enrichment process (NEP) and gut associated processes (GAP) and their effect on microbial biomass and activity. Soil Biol. Biochem., 12: 1573-1580.
30. Zhang, Q.L. and P.F. Hendrix, 1995. Earthworm (*Lumbricus rubellus* and *Aporrectodea caliginosa*) effects on carbon flux in soil. Soil Sci. Soc. Am. J., 59: 816-823.
31. Gorres, J.H., M.C. Savin and J.A. Amador, 1997. Dynamics of carbon and nitrogen mineralization, microbial biomass and nematode abundance within and outside the burrow walls of anecic earthworms (*Lumbricus terrestris*). Soil Sci., 162: 666-671.
32. Callaham, M.A. and P.F. Hendrix, 1998. Impact of earthworms (Diplocardia: megascolecidae) on cycling and uptack of nitrogen in coastal plain forest soils from northwest florida. USA. Appl. Soil Ecol., 9: 233-239.
33. Zhang, B.G., G.T. Li, T.S. Shen, J.K. Wang and Z. Sun, 2000. Changes in microbial biomass C, N, P and enzyme activities in soil incubated with the earthworms *Metaphire guillelmi* or *Eisenia fetida*. Soil Biol. Biochem., 32: 2055-2062.
34. Lachnicht, S.L. and P.F. Hendrix, 2001. Interaction of the earthworm *Diplocardia mississippiensis* (Megascolecidae) with microbial and nutrient dynamics in a subtropical spodosol. Soil Biol. Biochem., 33: 1411-1417.

35. Brown, G.G., 1995. How do earthworms affect microflora and fauna community diversity? *Plant Soil*, 170: 209-231.
36. Winding, A., R. Ronn and N.B. Hendrisken, 1997. Bacteria and protozoa in soil microhabitat as affected by earthworms. *Biol. Fertil. Soils*, 24: 133-140.
37. Aira, M., N. Mc Namara, T. Pearce and J. Dominguez, 2009. Microbial communities of *Lumbricus terrestris* L. middens, Structure, activity and changes through time in relation to earthworm presence. *J. Soil and Sediments*, 9(1): 54-61.