

## Detection of the *Coa* Gene in *Staphylococcus aureus* from Different Sources by Polymerase Chain Reaction

Ahlam A. Gharib, M.A. Adel Attia and M.M. Bendary

Department of Bacteriology, Mycology and Immunology,  
Faculty of Veterinary Medicine, Zagazig University, Zagazig, Egypt

**Abstract:** One hundred and ninety eight samples of human and animal sources from different areas in El-Sharkia Governorate were used in this study to isolate *Staphylococcus* species and to determine the presence of coagulase ( *coa* ) gene, a virulence gene detected in *S. aureus* (CoPS), by PCR. Primary isolation on mannitol salt agar,  $\alpha$ -haemolysis, tube coagulase test (TCT) and other biochemical characterization revealed that 32.8% (65/198) of the total samples were infected with staphylococci, only 35 isolates were biotyped as coagulase-positive staphylococci (CoPS) and represented as 8 (8.8%), 10 (22.2%) and 17 (26.9%) in milk, meat and human samples, respectively. Meanwhile, Coagulase-negative Staphylococci (CoNS) were obtained only from cows milk. Fifteen CoPS (5 milk samples, 4 meat products and 6 human samples) were subjected to analysis by PCR for detection of *coa* gene. The resulted amplicons were 648, 723, 812 and 913 bp. According to *coa* gene polymorphism, milk and meat isolates were grouped into 5 groups, while human isolates were grouped into 4 groups. It was noted that two of the strains, classified as coagulase negative by tube coagulase test were found to be positive with PCR amplification of the gene which clearly emphasizes the use of molecular methods in detecting *S. aureus*. In conclusion, the *coa* amplification has been considered as a simple and accurate method for typing of *S. aureus*.

**Key words:** Mastitis % Coagulase Polymorphism % CoPS % CoNS % Food Products

### INTRODUCTION

*Staphylococcus aureus* is considered as a significant pathogen of animal and human. Economic importance to dairy industry due to staphylococcal infection results from subclinical mastitis accompanied with reduction in the milk quality and a loss in its production. In spite of increasing frequency of isolation of coagulase negative staphylococci (CoNS) from the bovine mammary glands, CoPS is recognized worldwide as a major pathogen causing subclinical intramammary infection in dairy cows [1]. *S. aureus* is the most common bacteria, followed by *E.coli* and *S. agalactiae* in subclinical mastitis cases, while *E.coli* is the most common bacteria detected in clinical mastitis cases, followed by *S.aureus* and *S.agalactiae* [2].

*S. aureus* is a predominant colonizer of the skin. It is responsible for many different types of human infections. The predominant type of infection usually relates to skin abscesses and can occur in the form of

furuncles or carbuncles, but it rarely causes infection without a predisposing route of entry. Impetigo is a well known staphylococcal related infection which is characterized by honey-crusted lesions of the skin [3]. *S. aureus* has a large repertoire of virulence factors, including structural and secreted products that play a role in pathogenesis [4]. Staphylocoagulase is an extracellular protein that has traditionally been used to differentiate *S.aureus* from the less virulent staphylococci (CoNS). *S. aureus* secretes two clotting factors, coagulase (*Coa*) and von Willebrand factor binding protein (vWbp), the *Coa* and vWbp together are required for the formation of abscesses and promote the non proteolytic activation of prothrombin and cleavage of fibrinogen, reactions that are inhibited with specific antibody against each of these molecules. *Coa* and vWbp specific antibodies confer protection against abscess formation and *S. aureus* lethal bacteraemia, suggesting that coagulases function as protective antigens for a staphylococcal vaccine, so,

coagulases may be used as vaccine antigens to elicit antibodies that protect humans against *S. aureus* infections [5]. Moreover, abscesses formation, bacterial persistence in host tissues, blood clotting of coagulase producing *S. aureus* enable the pathogen to cause lethal sepsis, disseminate as thromboembolic lesion and resist opsonophagocytic clearance by host immune cells. Preclinical evidence suggests that inactivation or neutralization of coagulase may prevent the pathogenesis of staphylococcal infection [3, 6, 7].

Rapid and accurate typing of *Staphylococcus aureus* is crucial to understand the transmission of this infectious organism. The traditional phenotypic methods have several drawbacks [8]. Molecular typing can shorten or prevent an epidemic and reduce the number and cost of nosocomial infections. This typing can also clarify whether the strains from the environment, instruments, staff, or food are responsible for causing infection. In this way this helps to trace the source of infection or an outbreak, tracking the spread of infections and helps to take specific infection control measure [9]. The *coa* gene amplification has been considered a simple and accurate method for typing of *S. aureus* isolated from distinct sources, the coagulase protein is an important virulence factor of *S. aureus*. Like *spa*, *coa* has a polymorphic repeat region that can be used for differentiating *S. aureus* isolates. The variable region of *coa* is comprised of 81bp tandem short sequence repeats (SSRs) [10]. The objective of this study was to detect the presence of *coa* gene in *S. aureus* from different sources by PCR amplification.

## MATERIALS AND METHODS

**Samples:** One hundred and ninety eight samples of human and animal sources were collected from different areas in Sharkia Governorate in the period from January to May 2011. Samples of Human source (63) as blood (10), urine (12), C.S.F. (8), sputum (8), pericardial fluid (7), pus (10) and peritoneal fluid (8) were collected from different clinical cases of different ages administered to ultra laboratory and El Gamah hospital. Samples of animal source (135) were mastitic milk (90) collected from 4 farms (Abo-Mandor, Elsalhia, El hosania and Italian company farms) and random meat products (45) as minced meat (20), burger (15) and sausage (10) were collected from different sale markets. All samples were sent as soon as possible to laboratory of bacteriology, Faculty of Veterinary Medicine, Zagazig University and examined for staphylococci.

### Isolation and Identification of *Staphylococcus* Species:

All samples from different sources were inoculated onto mannitol salt agar (Difco,USA.) and incubated for up to one week at 37°C and all developed colonies with grapes like shape under light microscope were inoculated into soft agar (Difco,USA.) for culture preservation. The distribution of the isolates according to growth onto MSA, OF test and bacitracin susceptibility was detected [11]. The developed colonies onto MSA were subjected to tube coagulase test (TCT) [12] and for  $\beta$ -haemolysis onto blood agar to differentiate between the CoPS and CoNS. These isolates were further identified as *S. aureus* by API20S kits, Commercial identification system [13] and kept frozen at -20°C in nutrient broth with 3% glycerol (Difco,USA.) until molecular tests were carried.

**DNA Extraction:** The bacterial genomic DNA was extracted from only 15 *S. aureus* isolates (CoPS), using DNA extraction kits (Biofermentus).

**DNA Amplification:** The PCR for amplification of *coa* gene was performed in a total reaction volume of 25  $\mu$ l for one sample according to Himabindu *et al.* [14] using PCR master mix (DreamTaq™ Green Master Mix (2X), Fermentas, Catalogue No. #K1081., Lot: 00055548). It is a ready to use solution containing DreamTaq™ DNA polymerase, optimized DreamTaq™ Green buffer (2X), 4mM MgCl<sub>2</sub> and dNTPs (dATP, dCTP, dGTP and dTTP, 0.4 mM each). The sequence of primer used for amplification of *coa* gene [14] was forward: 5'CGAGACCAAGATTCAACAAG 3' and Reverse: 5'AAAGAAAACCACTCACATCA 3' The PCR cycling protocol was applied as following: initial denaturation at 94°C for 5 min, followed by 30 cycles of denaturation at 95°C for 30 second, annealing at 55°C for 45 second and extension at 72°C for 2 min, followed by a final extension at 72°C for 7 min.

### Agarose Gel Electrophoresis and Visualization of PCR Products:

5  $\mu$ l of each amplicon was electrophoresed in 1.5 % agarose gel (Sigma -USA) and visualized under U.V. transilluminator (Spectrolite Model TC- 312 A, USA) [15], then compared with the marker DNA ladder (100 bp, Gene Ruler™, Fermentas).

## RESULTS

**Culture Findings:** Staphylococci were isolated and identified by traditional phenotypic methods from

32.8% (65/198) of the samples represented 38, 10 and 17 isolates from milk, meat products and human samples, respectively.

Only one isolate of micrococcus was obtained from milk samples. On MSA agar, yellow colonies showing mannitol fermentation were selected and divided into 2 groups based on the coagulase test and  $\beta$ -hemolysis.

CoPS were confirmed as golden yellow pigmented colonies, coagulase positive and  $\beta$ -haemolytic 53.8%(35/65) represented 8 ( 8.8%), 10 (22.2%) and 17 (26.9%) from milk, meat and human samples, respectively.

Meat staphylococci isolates (10) recovered from meat products were distributed as following: (1 from minced meat, 4 from sausages and 5 from burger). Meanwhile, human staphylococci (17) were 3 from urine, 5 from pus, 3 from sputum, 2 from CSF, 1from blood, 2 from pericardial fluid and 1 from peritoneal fluid).

CoPS were predominated in isolates of meat products and human origin (the percentage of coagulase positive was equal to the percentage of  $\beta$ -haemolytic organisms 100%), meanwhile, in staphylococcal isolates of milk samples, the percentage of coagulase positive (21%) (8/38) was less than that of  $\beta$ -haemolytic (26%) 10/38, as two isolates were  $\beta$ -haemolytic but coagulase negative.

CoNS 46.2% (30/65) were isolated only from milk samples and could be confirmed by API 20-S. Biotyping of the isolates by analytical profile index (API20-S) revealed that all coagulase positive and  $\beta$ -haemolytic isolates were *S. aureus* in addition to the two isolates of milk samples that were coagulase negative and  $\beta$ -haemolytic.

**PCR Findings:** Most of isolates 80% (12/15) produced a single band, with molecular sizes ranging from 648-913 bp, whereas 2(13.3%) human isolates yielded 2 amplification products (double amplicons with molecular sizes of 723 + 812bp and 812+913 bp) and only 1 (6.6%) isolate of milk samples yielded 3 amplification products (triple amplicons with molecular sizes of 723+812 + 913 bp) (Photos 1&2, Figure 1).

The products 812 bp was the most frequent and accounted for 5/15 (33.3%) of the isolates, followed by 648 bp (26.6%), 723 bp (13.3%) and 913 bp (6.6%). According to *coa* gene polymorphism, the samples were grouped into 5 groups in (nine isolates of the milk and meat product) and 4 groups in ( the six isolates of human subjects) (Figures 2& 3). The two isolates from milk that were  $\beta$ -haemolytic and negative tube coagulase test, gave specific amplicon by *coa* gene amplification.




Fig. 1: Genotypic classes of *S. aureus* by *coa* gene polymorphism




Fig. 2: Genotypic classes of *S. aureus* of human origins




Fig. 3: Genotypic classes of *S. aureus* of animal origins




Photo 1: Coa gene polymorphism for *S. aureus* isolates of milk and meat products.

Lan 1 : +ve control obtained from central diagnostic and research lab. Kafrel -Sheikh university, Lan 2 (15M): with amplicon 648 bp, Lan 3 (18M): with amplicon 812 bp, Lan 4 (28M): with amplicon 648 bp, Lan 5 (139M): with 3 amplicons 723 bp , 812 bp and 913 bp, Lan 6 (180M): with amplicon 723 bp, Lan 7 (242Br) : with amplicon 723 bp, Lan 8 (243Br): with amplicon 913 bp, Lan 9 (217Mm) with amplicon 812 bp, Lan10 (233Sg): with amplicon 648 bp, Lan 11 (12M) and Lan 12 (140M): -ve control.




Photo 2: Coa gene polymorphism for *S. aureus* isolates of human subjects.

Lan 1: +ve control obtained from central diagnostic and research lab. Kafrel Sheikh university, Lan 2 (206Pu) : with two amplicon 812 pb and 913, Lan 3 (212Pu): with two amplicon 723 pb and 812 pb, Lan 4 (197Pu): with amplicons 812 pb, Lan 5 (210U): with amplicon 812 pb, Lan 6 (207Sp): with amplicon 812 pb, Lane7 (208Cs): with amplicon 648 pb, Lan 8(150M) : *S. Lentus* was used as -ve control.

## DISCUSSION

Molecular typing of microorganisms is now recognized as an essential component of infection control program. These molecular techniques are effective in tracking the spread of nosocomial infections and planning the activities of the infection control program. Production of coagulase is an important phenotypic feature used worldwide for the identification of *Staphylococcus aureus* [16]. The *coa* gene amplification has been considered as a simple and accurate method for typing of *Staphylococcus aureus*.

In this study, *S. aureus* was recovered from different human and animal sources, phenotypically and genotypically identified. 7 *coa* PCR types and 9 groups patterns were detected, which indicated considerable heterogeneity in the *coa* gene of *S. aureus* in the studied samples. Until recently, single and also double banded *coa* PCR products were reported in *S. aureus* strains derived from bovine mastitis in Brazil [17]. Double-banded amplification product was detected only in one *coa* positive isolate [16], which explained the presence of double-banded amplification products with different allelic forms of the *coa* gene.

Our amplification of DNA of *S. aureus* isolates obtained from human and animal sources revealed four amplicons (723, 812, 648 and 913 bp), These amplicons could classify the isolates into 4 groups (human) or

5groups (animal). Similar results were obtained by Himabindu *et al.* [14] who showed that the sizes of PCR products obtained after amplification of *S. aureus* of human subjects range from 650-1000 bps. Ishino *et al.* [18] classified 678 *S. aureus* isolates of human subjects into 8 classes and the sizes of the PCR products of *coa* gene ranged from 350 to 917 bp in increments of 81 bp. In addition, Schlegelova *et al.* [19] reported the size of *coa* gene PCR product of *S. aureus* isolates from dairy cow and human 650 -1050 bp, this result gave 4 classes at 650,730,810 and 1050 bp ;class 730 bp was the most common class between the isolates. Also Hookey *et al.* [20] detected the size of coagulase PCR product of *S. aureus* isolates from human subjects that was either 660, 603 or 547 bp. Furthermore, Reinoso *et al.* [21] detected that PCR amplification of the *coa* gene of *S. aureus* isolated from human, bovine subclinical mastitis and food samples which-yielded seven different *coa* types from 45 *S. aureus* strains with amplicon sizes ranging from 400 to 1000 bp. Seven (42%) human infection strains had *coa* polymorphic regions with an amplicon of 700 bp, indicating the presence of five repeats, six (40%) of the bovine strains showed an amplicon of 900 bp, indicating the presence of seven repeats and four (50%) strains from food samples showed four repeats.

Also, Da Silva and Da Silva [17] showed that the amplification of the *coa* gene from the 64 *S. aureus* isolated from cow with mastitis produces 27 different PCR- products; which ranged from approximately 579 to approximately 1442 bp. Sizes 790, 759, 725 and 579 bp were the most frequent. Two CoNS isolates showed similarity to *S. aureus* and were reclassified as CoPS species by API20s and *coa* gene detection with 648 and 812 bp amplicons. The same results were obtained by Himabindu *et al.* [14] and De Moura *et al.* [22] who noted that two of the strains that were classified as coagulase negative by tube coagulase test were found to be positive with PCR amplification of the gene . So the correct amplification of all isolates by PCR not only confirm the results of biochemical tests but is more accurate . Coagulase production is the principal criterion used by the clinical microbiology laboratory for the identification of *Staphylococcus aureus*. Numerous allelic forms of *S. aureus* coagulase exist, with each isolate producing one or more of these enzyme variants [23].

In conclusion, this study has shown that infection in the studied samples was caused by *S. aureus* strains harboring more than one *coa* genotype and that only one genotype predominated. However, further studies using a RFLP technique and nucleotid nucleotide

sequencing methods on a large collection of strains could be conducted to determine the common characteristics of the predominant strains.

## REFERENCES

1. Turutoglu, H., F. Tasci and S. Ercelik, 2005. Detection of staphylococcus aureus in milk by tube coagulase test. Bulletin of the Veterinary Institute in Pulawy, 49: 419-422.
2. Amin, A.S., R.H. Hamouda and A.A. Abdel-All, 2011. PCR Assays for Detecting Major Pathogens of Mastitis in Milk Samples World. Journal of Dairy and Food Sciences, 6(2): 199-206.
3. Friedman, D.A. and R.C. Ratard, 2007. The louisiana antibiogram. In vitro antibiotic sensitivity patterns 2003-2004: Louisiana State University School of Public Health and Louisiana Department of Health.
4. Defres, S., C. Marwick and D. Nathwani, 2009. MRSA as a cause of lung infection including airway infection, community acquired pneumonia and hospital-acquired pneumonia. European Respiratory Journal, 34(6): 1470-1476.
5. Cheng, A.G., M. Mcadow, H.K. Kim, T. Bae, D.M. Missiakas and O. Schneewind, 2010. Contribution of coagulases towards *Staphylococcus aureus* disease and protective immunity. Plos pathogen, 6(8): 1-17.
6. Chandrakanth, K., D.B.M. Virupakshaiah, C.C. Gavimath, M. Udaykumar and V.A. Kangralkar, 2010. Comparative genomics of *staphylococcus aureus* coagulase gene. Journal of Advanced Bioinformatics Applications and Research, 1(1): 31-36
7. Mcadow, M., D.M. Missiakas and O. Schneewind, 2012. *S.aureus* secretes coagulase and von willebrand factor binding protein to modify the coagulation cascade and establish host infections. Journal of Innate immune, 4(2): 141-148.
8. Tenover, F.C., R. Arbeit, G. Archer, J. Biddle, S. Byrne, R. Goering, G. Hancock, G.A. Hebert, B. Hill and R. Hollis, 1994. Comparison of traditional and molecular methods of typing isolates of *Staphylococcus aureus*. Journal of Clinical Microbiology, 32(2): 407-415.
9. Hacek, D.M., T. Suriano, G.A. Noskin, J. Kruszynski, B. Reisberg and L.R. Peterson, 1999. Medical and economic benefit of a comprehensive infection control program that includes routine determination of microbial clonality. American Journal of Clinical Pathology, 111: 647-654.
10. Van Belkum, A., S. Scherer, L. Van-Alphen and H. Verbrugh, 1998. Short-sequence DNA repeats in prokaryotic genomes. Microbiology and Molecular Biology Reviews, 62: 275-293.
11. Mackie, T.J. and J.E. McCartney, 1996. Practical Medical Microbiology, 14<sup>th</sup> ed., Chapter 11. U.S.A., Churchill Livingstone.
12. Gillespie, E.B., 1943. The routine used of the coagulase test for staphylococci. Monthly Bulletin of the Emergency zPublic Health Laboratory Service, 2: 19-22.
13. BioMerieux, S.A., 1992. Analytical profile index. BioMerieux Sa au capital de 45068 400 F/ imprime en France/ RCS Lyon B 673620399.
14. Himabindu, M., S.D. Muthamilselvan, D.K. Bishi and R.S. Verma, 2009. Molecular analysis of coagulase gene polymorphism in clinical isolates of methicillin resistant *Staphylococcus aureus* by restriction fragment length Polymorphism based genotyping. American Journal of Infectious Diseases, 5(2): 170-176.
15. Sambrook, J., E.F. Fritsch and T. Maniatis, 1989. Molecular cloning: A laboratory manual, 2<sup>nd</sup> Ed. Cold Spring Harbor Laboratory Press, New York, pp: 2001-2344.
16. Goh, H.S., K.S. Byrne, J.L. Zhang and A.W. Chow, 1992. Molecular typing of *Staphylococcus aureus* on the basis of coagulase gene Polymorphisms, Journal of Clinical Microbiology, 30(7): 1642-1645.
17. Da Silva, E.R. and N. Da Silva, 2005. Coagulase gene typing of *Staphylococcus aureus* isolated from cows with mastitis in southeastern Brazil. Canadian Journal of Veterinary Research, 69: 260-264
18. Ishino, K., N. Tsuchizaki, J. Ishikawa and K. Hotta, 2007. Usefulness of PCR-restriction fragment length polymorphism typing of the coagulase gene to discriminate arbekacin-resistant methicillin-resistant *Staphylococcus aureus* Strains. Journal Clinical Microbiology, 45(2): 607-609.
19. Schlegelova, J., M. Dendis, J. Benedik, V. Babak and D. Rysanek, 2003. *Staphylococcus aureus* isolates from dairy cow and human on a farm differ in coagulase genotype. Veterinary Microbiology Journal, 92: 327-334
20. Hookey, J.V., V. Edwards, B.D. Cookson and J.F. Richardson, 1999. PCR-RFLP analysis of the coagulase gene of *Staphylococcus aureus*: Application to the differentiation of epidemic and sporadic methicillin-resistant strains. Journal of Hospital Infection, 42: 205-212.

21. Reinoso, B.E., A. El-Sayed, C. Lämmner, C. Bognia and M. Zschöck, 2008. Genotyping of *Staphylococcus aureus* isolated from humans, bovine subclinical mastitis and food samples in Argentina. *Microbiological Research*, 163: 314-322.
22. De Moura, T.M., F.S. Campos, P.A. Azevedo, S.T. Van Der Sand, A.C. Franco, J. Frazzon and A.P.G. Frazzon, 2012. Prevalence of enterotoxin-encoding genes and antimicrobial resistance in coagulase-negative and coagulase-positive *Staphylococcus* isolates from black pudding] *Revista da Sociedade Brasileira de Medicina Tropical*, 45(5): 579-585.
23. Landolo, J.J., 1990. The genetics of staphylococcal toxins and virulence factors. In I.C. Gunsalus (ed), Academic Press, New York, pp: 399-426.