

## Callus Induction and Plant Regeneration in Finger Millet (*Eleusine coracana* L.)

E. Anjaneyulu, S. Hemalatha, S. Bharath Raj and M. Balaji

Department of Biochemistry, Sri Venkateswara University, Tirupati, A.P., India

**Abstract:** Shoot tips of finger millet (*Eleusine coracana*) were inoculated on to Murashige and Skoog (MS) medium supplemented with different concentrations and combinations of auxins [(2,4-D, NAA), cytokinins (KN, BAP) and other growth promoters (Proline, Tryptophan, casein enzyme hydrozylate (CEH)]. The higher degree of embryogenic callus formation observed with 2,4-D was at 2.5 mg/LG<sup>1</sup> in the presence of 0.5 mg/LG<sup>1</sup> of BAP. With NAA, the higher degree of embryogenic callus formation observed was at 3.0 mg/LG<sup>1</sup> in the presence of 1.0 mg/LG<sup>1</sup> of BAP. However, the addition of growth regulators like Tryptophan, Proline and CEH to either 2,4-D or NAA increased greatly the degree of callus formation. Among all the combinations employed the highest degree of callus formation was observed at 2.5 mg/LG<sup>1</sup> of 2,4-D, 0.5 mg/LG<sup>1</sup> of BAP, Proline (500 mg/LG<sup>1</sup>), Tryptophan (250 mg/LG<sup>1</sup>) and CEH (300 mg/LG<sup>1</sup>). A maximum of 97 % of shoot tip explants were successful in inducing callus at this combination of growth regulators. Highest mean number of shoot buds (67%) and good rooting was observed when embryogenic calli was subcultured on MS medium supplemented with 1 mg/LG<sup>1</sup> Kinetin and 0.5 mg/LG<sup>1</sup> NAA. The regenerated plantlets could be transferred successfully to the field with 85% survival.

**Key words:** MS medium % Callus formation % 2,4-Dichlorophenoxyacetic acid (2,4-D) % Alpha-Naphthalene acetic acid (NAA) % Benzyl amino purine (BAP) % Kinetin (KN)

### INTRODUCTION

Finger millet (*Eleusine coracana*) is an annual cereal crop widely grown in arid and semiarid areas of Africa and Asia [1]. The grain is a good source of protein, carbohydrate but contains less amount of fat. It is rich in calcium, phosphorus, iron, cysteine, tyrosine, tryptophan and methionine [2, 3]. Finger millet can be ground and cooked into cakes, puddings or porridge. The grain may also be malted and a flour of the malted grain could be used as a nourishing food for infants [4]. Production of transgenic plants with desired qualities is possible by genetic transformation of the desired genes in to the selected plants through the methodology of tissue culture. Efficient callus formation and regeneration is an important requisite to perform Agrobacterium mediated transformations for producing transgenics. Our present study is aimed to develop heavy metal resistance in finger millet by transferring phytochelatin synthase gene in to it. To achieve our aim we have developed an efficient protocol for callus induction and regeneration in finger millet.

Somatic embryogenesis is the most common method of plant regeneration in all the major species of cereals

and grasses [5]. Callus formation and regeneration in finger millet was reported first from mesocotyl explants [6]. [7] Thiru, A.N reported callus formation but no regeneration from different seedling explants. Mohanty [8] reported plant regeneration through shoot bud formation. [9] in a detailed morphogenic and histological study reported formation of apical dome like structures which, upon sub-culture produced multiple shoots. The first in vitro study in finger millet has been reported in which immature inflorescences and shoot apical meristem were induced to form somatic embryos on Murashige and Skoog [10] basal medium containing 2,4-Dichlorophenoxyacetic acid (2,4-D) and Kinetin (KN) [11, 12] and also from medium supplemented with picloram and kinetin [13]. The present paper describes efficient embryogenic callus formation and regeneration of finger millet.

### MATERIALS AND METHODS

Seeds of *Eleusine coracana* were obtained from Regional Agricultural Research Station, Tirupati. The seeds were sterilized and grown on filterpapers in petriplates at room temperature. From one week old

seedlings, shoot tips were collected, washed and inoculated on to MS basal medium supplemented with various growth regulators as outlined in the result section [10]. Sucrose at 3% was used as the sole carbon source. The cultures were then incubated in biotron with 16 hr of light cycle in a 24 hr cycle and the temperature regulated at  $25 \pm 1^\circ\text{C}$ . The culture media consisted of a) callus induction medium and b) regeneration medium.

## RESULTS AND DISCUSSION

**Callus Induction:** Different concentrations and combinations of various growth regulators were added to MS medium for effective callus induction. When compared to NAA, 2,4-D was found to be better for callus induction. The degree of callus formation was improved when 2,4-D (2.5 mg/L) was supplemented with 0.5 mg/L

Table 1: Callus induction from shoot tips of *Finger millet* in the presence of different growth regulators

Plant growth regulators (mg/L)	% of explants induced callus formation	Degree of callus formation
2,4-D		
0.5	12%	+
1.0	25%	+
1.5	35%	+
2.0	60%	+++
2.5	82%	+++
3.0	70%	+++
4.0	64%	++
NAA		
0.5	20%	+
1.0	35%	+
1.5	56%	+++
2.0	68%	+++
2.5	72%	+++
3.0	69%	++++
4	56	++
2, 4-D + KN		
0.5 + 0.25	26%	+
1.0 + 0.5	35%	+
1.5 + 0.75	55%	++
2.5 + 0.5	85%	++++
3.0 + 1.0	90%	++++
4.0 + 1.0	60%	++
2, 4-D + BAP		
0.5 + 0.25	20%	+
1.0 + 0.5	45%	+
1.5 + 0.75	64%	++
2.5 + 0.5	92%	+++
3.0 + 1.0	85%	++++
4.0 + 1.0	55%	++
NAA+ KN		
0.5 + 0.25	15%	+
1.0 + 0.5	25%	+
1.5 + 0.75	35%	+
2.5 + 0.5	75%	+++
3.0 + 1.0	88%	++++
4.0 + 1.0	54%	++
NAA+ BAP		
0.5 + 0.25	16%	+
1.0 + 0.5	35%	+
1.5 + 0.75	55%	++
2.5 + 0.5	78%	+++
3.0 + 1.0	80%	++++
4.0 + 1.0	60%	++
2,4-D +BAP+ Proline + CEH + Tryptophan		
2.0 +0.5+500 +300 +250	92%	+++
3.0 +0.5+ 500+ 300 + 250	97%	++++
NAA + BAP + Proline + CEH* + Tryptophan		
2.0 +0.5+500 +300 +250	75%	+++
3.0+ 0.5+500+ 300 + 250	88%	++++

\*CEH: casein enzyme hydrozylate

Table 2: Effect of growth regulators on shoot and root formation from embryogenic callus on MS medium (each treatment had 3 replicates)

Phytohormones mgLG <sup>l</sup>	No.of shoots/ plantlets (Mean ± SE)	No.of Roots (Mean ± SE)
<b>KN+NAA</b>		
0.5+0	38.3±4.2	33.6±3.2
1.0+0	59.5±6.4	24.5±2.5
2.0+0	42.1±2.1	38.6±7.2
0.5+0.025	48.3±6.2	18.1±3.1
1.0+ 0.25	67.3±7.2	39.8±5.2
2.0+ 0.55	32.5±5.4	26.5±6.4
<b>BA+NAA</b>		
0.5 +0	44.6±3.8	20.7±7.5
1.0 +0	62.6±7.2	26.4±3.5
2.0+0	56.8±8.1	30.6±2.4
0.5+0.025	56.6±4.2	19.7±6.5
1.0+ 0.25	64.3±8.2	33.9±8.3
2.0+ 0.55	41.6±3.5	23.2±1.6
<b>KN +2, 4-D</b>		
0.5+0.025	34.0±0.5	22.6±2.1
1.0+ 0.25	58±0.5	38.6±2.1
2.0 + 0.75	12.5±2.8	28.6±2.1
<b>BA+2,4-D</b>		
0.5+0.025	26±2.1	20.6±2.1
1.0+ 0.25	55±3.6	32.2±1.5
2.0 + 0.75	41.53±4.2	18.6±2.7




Fig. 1: Callus induction and plant regeneration from shoot tips of *Eleusine coracana*  
 A- Callus initiation on MS + 2.5 mgLG<sup>l</sup> 2,4-D and 0.5 mgLG<sup>l</sup> BAP  
 B- Callus formation and shoot buds induction on MS +2,4-D (3 mgLG<sup>l</sup>) +BAP( 0.5 mgLG<sup>l</sup> )+ Proline (500 mgLG<sup>l</sup>) + CEH (300 mgLG<sup>l</sup> + Tryptophan (250 mgLG<sup>l</sup>).  
 C, D & E- shoot growth and root formation from embryogenic calli on MS supplemented with 1 mgLG<sup>l</sup> Kinetin and 0.25 mgLG<sup>l</sup> NAA  
 F - Regenerated plants transferred to pots

of BAP. However, the highest degree of callus formation and highest per cent (97%) of explants that induced callus was achieved with 2,4-D (3 mgLG<sup>-1</sup>) + BAP (0.5 mgLG<sup>-1</sup>) supplemented with Proline (500 mgLG<sup>-1</sup>), Tryptophan (250 mgLG<sup>-1</sup>) and CEH (300 mgLG<sup>-1</sup>) (Table 1). Auxins play an important role in callus induction. The induction of callus in cereals and millets is commonly achieved by 2,4-D [14, 15]. Our results revealed that a combination of auxins (2,4-D) and low concentrations of cytokinins (BAP) are better inducers of callus than auxins alone. Supplementation of proline, tryptophan and CEH (casein enzyme hydrozylate) have further enhanced the efficiency of callus formation and the percent of explants that have given rise to callus. 2,4-D and KN have also been used for callus induction and somatic embryogenesis in the related species *Eleusine indica* [16]. Similar observations were noted by Kothari and Chandra [17] and Latha *et al.* [12] in finger millet.

**Plantlet Regeneration from Embryogenic Callus:** The well developed embryogenic calli were transferred to regeneration medium for shoots and roots formation. Among different combinations and concentrations of growth regulators employed, the highest mean number of shoots (67%) and roots (39 %) were formed when MS medium was supplemented with KN (1 mgLG<sup>-1</sup>) and NAA (0.25 mgLG<sup>-1</sup>) as shown in Table 2. Cytokinins play an important role in shoot growth. In our study KN with low concentration of NAA resulted in high mean number of shoots and good rooting. Similar reports were noted by others in finger millet [18, 19]. KN and BAP have also been used for efficient shoot induction and plant regeneration in pearl millet [20-23].

## REFERENCES

1. Rangan, T.S. and I.K. Vasil, 1984. In Ammirato PV *et al* (eds) Handbook of plant cell culture, vol, 3, Millets, Macmillan, New York, pp: 126-150.
2. Sastri, B.N., 1952. The Wealth of India. CSIR, New Delhi.
3. Malleshi, N.G. and H.S.R. Desikachar, 1986. Studies on comparative malting characteristics of some tropical cereals and millets. *J. Inst. Brew.*, 92: 174-176.
4. Magonja, M.A., J.M. Lenne, E. Manyasa, S. Sreenivasaprasad, 2007. Finger Millet Blast Management in East Africa. Creating opportunities for improving production and utilization of finger millet. Proceedings of the First International Finger Millet Stakeholder Workshop, Projects R8030 & R8445 UK Department for International Development-Crop Protection Programme. International Crops Research Institute for the Semi-Arid Tropics, pp: 196.
5. Park, C.H. and P.D. Walton, 1989. Embryogenesis and plant regeneration from tissue culture of Canada wild rye *Elymus Canadensis* L. *Plant Cell Reports*, 8: 289-291.
6. Rangan, T.S., 1976. Growth and plant regeneration in tissue culture of some Indian millets; *Paspalum scorbiculatum* (L.); *Eleusine coracina* (L.) Gaertn. and *Pennisetum typhoideum* Pers. *Pflanzphysiol.*, 78: 208-216.
7. Thiru, A.N. and Mohan Ram, 1990. Tissue culture studies in Ragi (*Eleusine coracana* (L.) Gaertn. *Plant Cell Reports*, 9: 93-96.
8. Mohanty, B.D., S.D. Gupta and P.D. Ghosh, 1985. Callus initiation and plant regeneration in ragi (*Eleusine coracana* (L.) Gaertn.). *Plant Cell Tissue Organ Cult.*, 5: 147-15.
9. Wakizuka, T. and T. Yamaguchi, 1987. The induction of enlarged apical domes invitro and multi-shoot formation from finger millet (*Eleusine coracana*), *Annals of Botany*, 60: 331-336.
10. Vasil, I.K., 1987. Developing cell and tissue culture systems for the improvement of cereal and grass crops. *J. Plant Physiol.*, 128: 193-218.
11. Murashige, T. and F. Skoog, 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures, *Physiol. Plant*, 15: 473-497.
12. Sivadas, P., S.L. Kothari and N. Chandra, 1990. High frequency embryoid and plantlet formation from tissue culture of the Finger millet- *Eleusine coracana* (L.) Gaertn. *Plant Cell Rep.*, 9: 93-96.
13. Latha M.A., V. Dasvantha Reddy, A. Madavi Latha and K. Venkateswara Rao, 2005. Production of transgenic plants resistant to leaf blast disease in finger millet (*Eleusine coracana* (L.) Gaertn.). *Plant Sci.*, 169: 657-667.
14. Eapen, S. and L. George, 1990. Influence of phytohormones, carbohydrates, amino acids, growth supplements and antibiotics on somatic embryogenesis and plant differentiation in finger millet. *Plant Cell Tissue Organ Cult.*, 22: 87-93.
15. Jabeen, M. and S. Siriwardana, 1987. Selection for salt tolerance in rice (*Oryza sativa* L.) *Mod. Trends. Pl. Sci. Res. Pak.*, pp: 164-169.

16. George, L. and S. Eapen, 1990. High frequency plant regeneration through direct shoot development and somatic embryogenesis in immature inflorescence cultures of Finger millet (*Eleusine coracana* (L.) Gaertn). *Euphytica*, 48: 269-274.
17. Chowdhry, C.N., A.K. Tyagi, N. Maheshwari and S.C. Maheshwari, 1993. Effect of L-proline and L-tryptophan on somatic embryogenesis and plantlet regeneration of rice (*Oryza sativa* L.). *Plant Cell Tissue Organ Culture*, 32(3): 357-361.
18. Evans, D.A., W.R. Sharp and C.E. Flick, 1981. Plant regeneration from cell cultures. *Hort. Rev.*, 3: 214-314.
19. Kothari, S.L. and N. Chandra, 1995. Advances in tissue culture and genetic transformation of cereals. *J. Indian Bot Soc.*, 74: 323-342.
20. Yemets, A., L. Klimkina, L. Tarassenko and Y. Blume, 2003. Efficient callus formation and plant regeneration of goose grass (*Eleusine indica* (L.) Gaertn.). *Plant Cell Rep.*, 21: 503-510.
21. Mythili, P., A. Madhavi, V.D. Reddy and N. Seetharam, 2001. Efficient regeneration of pearl millet *Pennisetum glaucum* (L.) from shoot tip cultures. *Indian J. Exp. Biol.*, 39: 1274-1279.
22. Srivastav, S. and S.L. Kothari, 2002. Embryonic callus induction and high frequency plant regeneration in pearl millet. *Cer. Res. Commun*, 30: 69-74.
23. Nayak, P. and S.K. Sen, 1989. Plant regeneration through somatic embryogenesis from suspension cultures of a minor millet, *Paspalum scorbiculatum* (L.). *Plant Cell Rep.*, 8: 296-299.