

Feeding Value Potential of Mulberry (*Morus alba*) Leaf Meal to Replace Concentrate Mix

¹Gebrekidan Tesfay, ²Berhan Tamir and ²Gebreyohannes Berhane

¹Department of Animal Production and Technology,
Adigrat University, P.O. Box 50, Tigray, Ethiopia

²Department of Animal Production Studies,
Addis Ababa University, P.O. Box 34, Debre Zeit, Ethiopia

Abstract: This experiment was carried out to estimate the feeding value of mulberry (*Morus alba*) leaf meal by evaluating its chemical composition, *in vitro* organic matter digestibility and *in sacco* dry matter (DM) degradability. The objective of the study was to assess the potential of mulberry foliage in supplementing the feed of ruminant animals during the dry season when other feed resources are scarce and their quality generally fall short of animal requirements. Dry matter (DM), crude protein (CP), ash, ether extract (EE), crude fibre (CF), neutral detergent fibre (NDF), acid detergent fibre (ADF) and acid detergent lignin (ADL) contents of the different diets was determined. *In vitro* organic matter digestibility was determined by the two-stage method. *In sacco* rumen degradability was also measured using three rumen fistulated Boran-Holstein Friesian cross steers at 0, 6, 12, 24, 48, 72 and 96 hours. The composition of mulberry leaf meal compares favorably with that of the concentrate mixture in most of the nutrients. The ash content of mulberry leaf meal was more than double to that of the concentrate mixture. Similarly, the calcium value of mulberry leaf meal was more than threefold to that of calcium content of concentrate mixture. Mulberry leaf meal alone had the greatest values for slowly degradable fraction (*b*) than the diets with less proportion of mulberry. Whereas mulberry leaf meal alone (T5) and 75g concentrate mix + 259.7g mulberry leaf (T4) had significantly ($p < 0.05$) less soluble fraction (*a*) and effective degradability (*ED*) values as compared to the diets with less proportion of mulberry. In a nutshell, all the treatment diets recorded more than 66% DM degradability at 24 h, which implied that they were all greatly degradable in the rumen.

Key words: Chemical composition • Degradability • Digestibility • *Morus alba*

INTRODUCTION

In areas like Ethiopia where livestock are closely integrated with crop production, crop residues are considered as valuable sources of ruminant feeds. However, crop residues are fibrous having low digestible organic matter and low crude protein content and generally are low in nutritive value. This low nutritive values restrict its utilization by rumen microorganisms and consequently by the host animal. An adequate supply of nitrogen (N) to rumen microbes is crucial to obtain maximum rate of plant cell digestion as well as a high microbial protein synthesis [1- 3]. There are several alternatives for improving the performance of ruminants

fed these low nutritional quality fibrous diets. Supplementation with concentrate could improve the nutritional value of low quality roughage such as straws and could be one alternative. However, the concentrate diets are often expensive and not accessible to smallholder farmers. In many tropical countries and regions, there has been a focus on indentifying and using locally available shrubs and tree leaves as cheaper alternative supplements for ruminants because of their high nutritive value and positive effects on rumen function [4].

Additionally, several works in different places have reported the potential of the mulberry (*Morus alba*) forage for sustainable ruminant production [5-10]. Hence,

mulberry might be potentially used as supplement to low quality roughage diets due to its high protein content [11] and degradability [12]. Beside its high protein content, the degradability of its organic matter is also high [12] so it can supply fermentable energy in the rumen. Moreover, it can create favorable condition in the rumen for plant cell wall degrading microorganisms [5].

Therefore, this experiment was carried out to compare the chemical composition, *in vitro* digestibility as well as *in sacco* degradability of sole mulberry leaf meal, concentrate mix and their mixtures.

MATERIALS AND METHODS

Sample Preparation and Experimental Design: Leaves from mulberry that were harvested from nearby farmers and nursery areas were dried under a shade for 4-5 days till the leaves were easily crushed when pressed in a hand and were packed in a sack for later use. The concentrate mixture which was purchased from animal feed shop was comprised of wheat bran and noug seedcake at the ratio of 2:1 on dry matter (DM) bases, in that order. Five treatment diets were designed in such a way that concentrate mix was progressively replaced by mulberry leaf meal from 0% to 100% at iso-nitrogenous level. The diets include, 300g concentrate mix alone (T1), 225g concentrate mix + 86.55g mulberry leaf (T2), 150g concentrate mix + 173.1g mulberry leaf (T3), 75g concentrate mix + 259.7g mulberry leaf (T4) and 346.2g mulberry leaf alone (T5).

Chemical Analysis: Chemical composition of the feed samples was determined at Holeta Agricultural Research Center, animal nutrition laboratory, Ethiopia. Samples of partially dried feeds were dried overnight at 105 °C in a forced draft oven for determination of total dry matter content. Samples were analyzed for DM, ash and nitrogen (N) using the procedure of AOAC [13] and crude protein (CP) was calculated as N x 6.25. Neutral detergent fiber (NDF), acid detergent fiber (ADF) and acid detergent lignin (ADL) were analyzed using the procedures of Van Soest *et al.* [14].

***In vitro* Organic Matter Digestibility:** *In vitro* organic matter digestibility was determined by the two-stage method of Tilley and Terry [15]. Samples were incubated for 48 hours with rumen fluid and buffer followed by another 48 hour digestion with pepsin and HCl and the residue was ashed in a muffle furnace at 550 °C for 5 hour. Rumen fluid was obtained from rumen fistulated Boran x

Holstein Friesian steers kept at maintenance dietary condition with diets containing hay supplemented daily with 4 kg of concentrate mixture comprised of 74, 25 and 1% wheat bran, noug seed cake and salt, respectively.

***In Sacco* Dry Matter Degradability:** Rumen degradability of the samples was determined by incubating about 3 gm of sample in a nylon bag (41µm pore size and 6.5x14 cm dimension) in three rumen fistulated Boran x Holstein Friesian steers kept at Holeta Agricultural Research Station. The steers were offered grass hay supplemented daily with 4 kg of concentrate mixture comprised of 74, 25 and 1% wheat bran, noug seed cake and salt, respectively. The steers were kept in individual pens. The feed samples were incubated for 0, 6, 12, 24, 48, 72 and 96 hours. Duplicate nylon bag containing samples were incubated in three rumen fistulated animals by placing the samples at different hours and taking them out at the same time (sequential addition). At the end of the incubation period, sample containing bags, including zero hour bags were hand washed in a running tap water. The washed bags were then dried in an oven at 105 °C for 24 hours to determine dry matter. The dried bags were then taken out of the oven and allowed to cool down in desiccators and weighed immediately. The ruminal *in sacco* DM degradability at each incubation time was determined as following $DMD (g/kg DM) = (DM \text{ in feed sample} - DM \text{ in residue})/DM \text{ in feed sample}$.

The DMD data was fitted to the exponential equation $p = a + b(1 - e^{-ct})$ as described by Ørskov and McDonald [16], where **p** is the amount of nutrient degraded (%) at time **t**, **a** is the intercept of the degradation curve at time zero and represent as degradability of soluble fraction (%), **b** is the rumen-insoluble, but slowly degradable fraction (%), **c** is base for natural logarithm **c** is the rate constant for degradation of the **b** fraction (%/h), **t** is the incubation time (h) and the lag phase (L) = $(1/c) \log_e [b/(a+b - A)]$ [17]. Potential degradation (PD) was estimated as (A + B), while effective degradability (ED) of DM was calculated using the formula: $ED = A + [B * c / (c + k)]$, where **A**, **B** and **c** are as described above and **k** is rumen outflow rate, which is assumed to be 0.03/h for roughage feeds [16]. The calculation of the equation was carried out using the NEWAY program [18].

Statistical Analysis: Data from the experiments on *in vitro* organic matter digestibility and *in sacco* dry matter degradability were subjected to the analysis of variance (ANOVA) in a randomized complete block design using

the general linear model procedure of SAS [19]. Individual differences between means were tested using Tukey HSD test. In all the comparisons, the level of significance was set at $\alpha = 0.05$.

RESULTS

Chemical Composition of the Feeds Used in the Experiment: The diets were formulated to meet the Crude protein (CP) requirements of yearling Tigray highland lambs. The composition of mulberry leaf meal compares favorably with that of the concentrate mixture in most of the nutrients and even it was better in its ash content than the concentrate mixture (Table 1). The dry matter content seems similar across the different treatments where as the organic matter content declined as the proportion of concentrate mixture decreases. The ash content of mulberry leaf meal in the current study was more than double to that of the concentrate mixture. Similarly, the calcium value of mulberry leaf meal was more than threefold to that of concentrate mixture.

In vitro Organic Matter Digestibility and in Sacco Dry Matter Degradability: There was less difference in *in vitro* organic matter digestibility (76.42-77.78%) across the different treatments in the current study. All the tested diets resulted more than 76% *in vitro* organic matter digestibility. The *in sacco* dry matter degradability characteristics of sole mulberry foliage and concentrate mix or their mixtures are presented in Table 2. The greatest and least soluble fraction (A) was recorded in the diet with higher proportion of concentrate mix and in the sole mulberry leaf meal, respectively. The amount of washable materials (A) significantly increased as the proportion of concentrate mix increased.

The washable materials were significantly higher in T1 and T2, than the remaining treatments. Unlike the washable materials, the reverse is true in the case of the potential degradation for non-water soluble materials (B). The insoluble but fermentable fractions increased as the proportion of mulberry leaf meal advanced. The slowly degradable fraction (B) value obtained in mulberry leaf meal was superior (53.2%) as compared to values recorded for the sole concentrate mix. The potential degradability of T4 and T5 in the current study was significantly less than those of T1, T2 and T3. There was less difference in the rate of degradation across the different treatments in this study. Slight increase in degradation rate constant was observed until T3, which declined steadily at higher mulberry leaf meal level.

Table 1: Chemical composition of the feed ingredients

Chemical composition	Treatments				
	T1	T2	T3	T4	T5
DM (%)	90.2	89.5	89.2	89.5	91.8
OM (%DM)	92.34	89.70	86.06	85.59	84.30
Ash (%DM)	7.66	10.30	13.94	14.41	15.70
CP (%DM)	22.03	22.13	20.01	18.51	18.49
CF (%DM)	13.27	12.84	13.56	13.78	14.17
NDF (%DM)	35.18	34.19	35.23	36.78	37.98
ADF (%DM)	20.17	19.47	21.66	19.91	22.32
ADL (%DM)	3.39	3.42	3.36	3.83	4.19
EE (%DM)	7.16	6.33	5.91	5.34	4.15
Ca (%DM)	0.6	0.76	1.08	1.45	2.11
P (%DM)	1.02	0.81	0.94	0.84	0.77

MLM= Mulberry leaf meal; CM= Concentrate mix (2:1 of wheat bran to noug seed cake); DM=Dry matter; OM=Organic matter; CP=Crude protein; CF= Crude fiber; NDF=Neutral detergent fiber; ADF = Acid detergent fiber; ADL = Acid detergent lignin; EE= ether extract; Ca= calcium; P= phosphorus

Table 2: *In sacco* dry matter degradability characteristics of sole mulberry foliage and concentrate mix or their mixtures (%DM)

Parameters	Treatments					SEM	SL
	T1	T2	T3	T4	T5		
A	26.2 ^a	26.1 ^a	24.1 ^b	22.7 ^c	22.2 ^c	0.25	***
B	51.1 ^b	51.1 ^b	52.5 ^{ab}	52.6 ^{ab}	53.2 ^a	0.52	*
PD	77.0 ^a	77.2 ^a	76.6 ^a	75.3 ^b	75.3 ^b	0.30	***
ED	63.6 ^a	63.5 ^a	62.8 ^a	60.5 ^b	58.9 ^c	0.36	***
C/hr	0.087	0.089	0.090	0.086	0.084	0.00	NS
L (hr)	4.27 ^a	4.43 ^a	5.03 ^b	4.90 ^b	4.87 ^b	0.08	***
<i>In vitro</i> DOM	77.46	77.47	76.98	77.78	76.42	1.34	0.861

^{a-c}Means with different superscripts in a column within a category differ; *** = $\bar{N} < 0.001$; * = $\bar{N} < 0.05$; NS = not significant; SEM = standard error of the mean; SL= significant level A = rapidly soluble fraction; B = insoluble but fermentable fraction; PD = potential degradability; C = the rate of degradation of B; L = lag time; ED = effective degradability; DOM = Digestibility of organic matter; hr = hour

The trend of degradation of different levels of mulberry leaf meal in combinations with concentrate mixture showed a time dependent increase (Figure 1). The trend of degradability of T1 and T2 increased consistently up to 24 hr but it continues increasing steadily up to 48hrs. However, the progress seems constant beyond the 48 hrs. Nevertheless, the trend of degradation for T3, T4 and T5 was slow up to 6 hours; later T3 increased rapidly to 12 hours compared to T4 and T5 and continues to increase exactly the same as T1 and T2. From 6 – 24 hours, T4 and T5 showed similar increment. After 24 hrs T1, T2, T3 and T4 started to decrease their degradation rate, while T5 (sole mulberry leaf meal) keeps increasing up to 48 hrs and continued increasing slowly till 72 hours. As shown in Figure 1, the extent of degradation of all the treatment




Fig. 1: *In sacco* dry matter disappearances of the different treatment feeds.

diets beyond 72 hours seems similar. After 48 hours incubation the total dry matter degradability values showed non-significant effect across the different treatment diets. Additionally, T1, T2 and T3 showed significantly higher degradability values than that of T4 and T5 in 12 to 48 h incubation. In a nutshell, all the treatment diets recorded more than 66% DM degradability at 24 hours, which implied that they were all potentially degradable in the rumen.

DISCUSSION

Chemical Composition: Since the feeds used in the current experiment were dried under sun, their dry matter was expectedly high (Table 1). The organic matter content declined as the proportion of concentrate mixture decreases. This might be due to the higher ash content of mulberry leaf meal. The NDF, ADF and ADL compositions of the treatment diets were not deviated largely, justifying the comparable potential of mulberry foliage to that of concentrate mix. The CP content of mulberry leaf meal obtained in this study was within the range of values (18 to 25%) obtained by Ba *et al.* [20], Kandylis *et al.* [7] and Vu *et al.* [21]. The NDF content of sole mulberry leaf meal was to some extent higher than 31.1% and 26.4% reported by Vu *et al.* [21] and Habib *et al.* [22] respectively. Whereas, its ADF content was within the range of values (17.4 – 24.7 %) detected by Kabi and Bareeba [23] as well as Habib *et al.* [22]. The ADL content of mulberry leaf meal in the present study is in accordance with of the results that Kandylis *et al.* [7] and Atiso *et al.* [24] have reported (4.1% and 4.65%, respectively). In three clones

of mulberry leaves, Schmidek *et al.* [25] noted comparable NDF (30.2–39.3%) and ADF (17.2–21.7%) results with the present finding.

Feeds with high ADF content could lower the availability of nutrients since there is a negative relationship between ADF and digestibility of feeds [26]. Moreover, Kabi and Bareeba [23] also demonstrated that NDF, ADF and ADL contents increased with increasing maturity. However, the values for fibre fractions of mulberry leaf meal obtained in this experiment were low. This idea further corroborated the comparative supplementary feeding value of mulberry for ruminants. According to Lonsdale [27] feeds that have <12%, 12-20% and >20% CP are classified as low, medium and high protein sources, respectively. Hence, mulberry leaf meal in the current study could be categorized among the medium upper limit protein source feed that can serve as a protein supplement for low quality crop residues particularly during the dry season. Similarly, Benavides [28] also noted mulberry leaves have a high potential as a protein-rich forage supplement to be used in feed for ruminants, monogastrics and rabbits.

The ash content (15.4%) of mulberry reported by Habib *et al.* [22] was comparable with the current finding. Similarly, Singh and Makkar [29] stated that since mulberry leaves are rich in mineral matters like calcium, nitrogen, sulphur and other minerals they have the potential to be used as a supplementary feed for improving livestock productivity and this conforms to the current study. According to McDowell [30], the recommended Ca to P ratio for ruminants is 2:1 to 4:1. Hence, the ratio obtained in the current study was within

this range and the good mineral contents of mulberry foliage in this study further provoke its use as good alternative source of feed for ruminants. Ether extract obtained from the current study was higher than the value reported by Hurtado *et al.* [31] (2.1%) whereas, it was comparable with 4.21% and 3.69% reported by Wang *et al.* [32] and Doran *et al.* [33], respectively. This suggests mulberry is not an energy deficient plant. Therefore, the leaf from this plant can be considered as a cheap and affordable supplement for ruminant animals particularly in herds of small scale rural farmers.

In vitro Organic Matter Digestibility: The comparable digestibility of mulberry with the commercial concentrate mixture ensures the potential of the leaf supplementing the less quality crop residues in ruminant production. The results that we have obtained for *in vitro* organic matter digestibility are relatively higher than the results for *in vitro* digestibility of browse species during wet season (68%) and dry season (72%) reported by Gizaw *et al.* [3]. Moreover, Shayo [34] reported that *in vitro* digestibility of mulberry leaves was 82.1% which is to some extent larger than the current study. Similarly, Bakshi and Wadhwa, [35] also noted that *in vitro* digestibility of mulberry leaves was higher than other forage species. The better digestibility of mulberry could be attributed to high CP level and increased concentration of ammonia nitrogen in rumen [36]. Higher ammonia nitrogen in rumen improves microbial activity and growth of fibrolytic bacteria resulting in more DM digestibility [37]. The narrow range of *in vitro* digestibility observed in the current study among the different treatment feeds might be a reflection of similarities in their chemical composition and potential of the mulberry being comparable with concentrate mix.

The threshold level of NDF in tropical plants beyond which feed intake of ruminants is affected is 60% [38]. Tree forages with a low NDF concentration (20–35%) are usually of great digestibility [35]. The NDF result of this study is in accordance with the results reported by Schmidek *et al.* [25] and Cheema *et al.* (2011) for multipurpose trees and shrub species. Therefore, the NDF (37.98%) value obtained in the present experiment could be regarded as a medium one. This attribute can induce even greater fermentation rate, therefore, improving its digestibility [39].

In sacco Degradability: The sole mulberry leaf meal (T5) had the greatest value for slowly degradable fraction (*b*) than the diets with less proportion of mulberry. Additionally, T5 and T4 had significantly ($p < 0.05$) less

soluble fraction (*a*) and effective degradability (ED) values compared to the diets with less proportion of mulberry (Table 1). The higher washable materials in T1 than T5 might indicate relatively higher content of dusty and small particles which could easily pass or wash from the bag, rather than a greater solubility. Similarly, Promkot and Wanapat [40] reported that the result of the rumen bag technique depends on the way the feed is prepared and the pore size of the material from which the bag is made. Unlike the washable materials, the opposite is true in the case of the potential degradation for non-water soluble materials (B).

According to Belachew *et al.* [41], multipurpose trees could be assigned to great (> 450 g/kg DM), medium (400–450 g/kg DM) and low (< 400 g/kg DM) quality groups based on their ED values. All the experimental diets in the current study including the mulberry foliage belonged to great quality group (Table 2). This further clarifies the comparable nutritional potential of mulberry foliage with that of concentrate mix and justifies the nutritive potential of mulberry which could supplement the less quality cereal crop residues in ruminant feed.

Six hours post incubation; the greatest DM disappearance was recorded for T1 and T2, which have higher proportion of concentrate mix in the diet. Even though DM disappearances of mulberry leaf meal were lower compared to concentrate mix, six hours post incubation; these results were comparable with other multipurpose trees. Belachew *et al.* [41] noted that six hours post incubation, the greatest DM disappearances was recorded for the leaves of multipurpose trees like *E. capensis* (32.12%) and *M. lanceolata* (32.60%) and the least DM disappearances was determined for *F. sycomorus* (15.82%) and *R. glutinosa* (16.87%) leaves. In the present study, dry matter disappearance increased with rumen incubation time for all experimental feeds. This is in accordance with the findings of Belachew *et al.* (2013[41], Lebopa *et al.* [42], Vranic *et al.* [43], Lanyasunya *et al.* [44] and Tesema *et al.* (2003) which confirmed that *in sacco* degradation of dry matter increase as incubation time advances.

The obtained results for potentially degradable and rate of degradation are higher than the results reported by Suchitra and Wanapat [45]. However, the washable (39.2), potential degradability (84.7) and effective degradability (64.2) values were higher than the values obtained in the current study. The rate of degradations obtained in the current study (0.084 to 0.090) were higher than most of the multipurpose trees reported by Belachew *et al.* [41] and to

some extent is in accordance with that of Suchitra and Wanapat [45]. They have reported that the rate of degradation for Cassava hay (0.086) was highest compared to the other treatments resulting in low gut fill which could lead to higher intake and animal production. Furthermore, the lag time was at its minimum for T1 (4.27 h) and then increased and showed maximum value for T3 (5.03 h). In short, the amount of dry matter found degraded in all tested feeds at 12 and 24 hours were more than 50 % and 66 %, respectively, which implied that they were all greatly degradable in the rumen.

Generally, the results of the current study indicated that mulberry foliage appears to be rich in protein and minerals and low in fiber contents. Mulberry leaf meal by virtue of having high CP, good Ca to P ratio, promising digestibility and digestion rate as well as good extent of digestion is compatible to concentrate mix. In conclusion, mulberry leaf meal could potentially be used as a supplement to poor quality roughage diets, particularly during the dry season and could substitute the costly and not easily accessible protein concentrates by poor farmers in Ethiopia.

REFERENCES

1. Adugna Tolera, 2008. Feed resources and feeding management: A manual for feedlot operators and development workers. Ethiopian Sanitary and Phytosanitary Standards and Livestock and Meat Marketing Program (SPS-LMM), Addis Ababa, pp: 43.
2. Gül, S., M. Keskin and S. Kaya, 2010. Olive Cake usage as an Alternative to Cotton Seed Meal in Dairy Goat Feeding. African Journal of Agricultural Research, 5: 1643-1646.
3. Gizaw, S, A. Tegegne, B. Gebremedhin and D. Hoekstra, 2010. Sheep and goat production and marketing systems in Ethiopia: Characteristics and strategies for improvement. IPMS (Improving Productivity and Market Success) of Ethiopian Farmers Project Working Paper 23. ILRI (International Livestock Research Institute), Nairobi, Kenya, pp: 58.
4. Yao, J., B. Yan, X.Q. Wang and J.X. Lui, 2000. Nutritional evaluation of mulberry leaves as feeds for ruminants. Livest. Res. Rural Dev., 12(2): <http://www.lrrd.org/lrrd12/2/yao122.htm>
5. Yulistiani, D., Z.A. Jelan and J.B. Liang, 2008. Degradability of mulberry (*Morus alba*) and rice bran in the rumen of sheep fed different diets. JITV, 13(4): 264-272.
6. Martín, G.J., Y. Noda, G. Pentón, D.E. García, F. García, González E, Ojeda F, Milera M, Lopez O, Ly J, Leiva L, Arece J., 2007. Mulberry (*Morus alba*, Linn.): a species of interest for livestock production. Pastosy Forrajes, 30, 3–19.
7. Kandyliis, K., I. Hadjigeorgiou and P. Harizanis, 2009. The nutritive value of mulberry leaves (*Morus alba*) as a feed supplement for sheep. Trop. Anim. Prod., 41: 17-24.
8. Soca, M., F. Ojeda, D.E. García and M. Soca, 2010. Effects of feeding mulberry (*Morus alba*) forage on the productive and health indicators of grazing calves. Pastosy Forrajes, 33: 1-10.
9. Salinas-Chavira, J., O. Castillo-Martínez, Ramirez, J.E. Bribiesca and M. Mellado, 2011. Effect of increasing levels of white mulberry leaves (*Morus alba*) on ruminal dry matter degradability in lambs. Tropical Animal Health and Production, 43(5): 995-999.
10. González-García, E. and G. Martín Martín, 2016. Biomass yield and nutrient content of a tropical mulberry forage bank: effects of season, harvest frequency and fertilization rate. Grass and Forage Science, (Online DOI: 10.1111/gfs.12227).
11. Sanchez, M.D., 2002. World distribution and utilization of mulberry and its potential for animal feeding. FAO Animal Production and Health Paper 147, Rome, Italy; pp: 345.
12. Saddul, D., Z.A. Jelan, J.B. Liang and R.A. Halim, 2005. Evaluation of Mulberry (*Morus alba*) as Potential Feed Supplement for Ruminants: The Effect of Plant Maturity on In situ Disappearance and In vitro Intestinal Digestibility of Plant Fractions. Asian-Aust. J. Anim. Sci., 18(11): 1569-1574.
13. AOAC (Association of Official Analytical Chemists), 1990. Official method of analysis. 15th edition. AOAC, Inc., Arlington, Virginia, USA, pp: 12.
14. Van Soest, P.J., B.J. Robertson and B.A. Lewis, 1991. Methods for dietary fiber, neutral detergent fiber and non-starch polysaccharides in relation to animal nutrition. Journal of Dairy Science, 74: 3583-3597.
15. Tilley, J.M.A. and R.A. Terry, 1963. A two stage technique for the in vitro digestion of forage crops. Journal of British Grassland Society, 18: 108-112.
16. Ørskov, E.R. and I. McDonald, 1979. The estimation of protein degradability in the rumen from incubation in measurements weighted according to rate of passage. Journal of Animal Science, 88: 645-650.
17. Ørskov, E.R. and M. Ryle, 1990. Energy Nutrition in Ruminants. Elsevier, Oxford, pp: 149.

18. Chen, X.B., 1996. An Excel Application Program for Processing Feed Degradability Data. User Manual, Rowett Research Institute, Buchsburn, Aberdeen, UK.
19. SAS, 2008. SAS/STAT Guide to personal computers, version 9.2. Statistical Analysis System Institute Inc., NC., USA.
20. Ba, N.X., V.D. Giang and L.D. Ngoan, 2005. Ensiling of mulberry foliage (*Morus alba*) and the nutritive value of mulberry foliage silage for goats in central Vietnam. Livestock Research For Rural Development, 17(15).
21. Vu, C.C., M.W.A. Verstegen, S.W.H. Hendrik and K.C. Pham, 2011. The nutritive value of mulberry leaves (*Morus alba*) and partial replacement of cotton seed in rations on the performance of growing Vietnamese cattle. Asian-Australasian Journal of Animal Science, 24: 1233-1242.
22. Habib G., N.A. Khan, A. Sultan and M. Ali, 2016. Nutritive value of common tree leaves for livestock in the semi-arid and arid range lands of Northern Pakistan. Livestock Science, 184: 64-70.
23. Kabi, F. and F.B. Bareeba, 2008. Herbage biomass production and nutritive value of mulberry (*Morus alba*) and Calliandra calothyrsus harvested at different cutting frequencies. Animal Feed Science and Technology, 140: 178-190.
24. Atiso, M., Y. Tesfay and T. Alemayehu, 2012. Substituting mulberry leaves (*Morus indica L.*) for concentrate mix in the diet of lactating Holstein Friesian cows. J. Dry lands, 5: 372-376.
25. Schmidek, A., R. Takahashi, A.N. Medeiros and K.T. Resende, 2000. Bromatological Composition and Degradation rate of Mulberry in Goats. Pp. 207-210. In: Sanchez (ed.), Proceedings of electronic conference, August, 2002, Rome, Italy.
26. McDonald, P., R.A. Edwards, J.E.D. Greenhalgh and C.A. Morgan, 2002. Animal Nutrition, 6th ed., (Prentice Hall, USA). pp: 607.
27. Lonsdale, C., 1989. Raw Materials for Animal Feed Compounders and Farmers. Straights. (Book) Chalcombe publications, Great Britain.
28. Benavides, J.E., 2000. Utilisation of Mulberry in Animal Production Systems, In: Sanchez (ed.), Mulberry for Animal Production, Proceeding of Electronic Conference.
29. Singh, B. and H.P.S. Makkar, 2002. The potential of mulberry foliage as feed supplement in India. pp: 139-155. In: Sanchez (ed.), Proceedings of electronic conference, August 2002, Rome, Italy.
30. McDowell, L.R., 1997. Minerals for Grazing Ruminants in Tropical Regions. University of Florida. Gainesville.
31. Hurtado, D.I., S. Nocua, W. Narvaez-Solarte and E. Vargas-Sanchez, 2012. Nutritional value of mulberry (*Morus sp.*), matarratón (*Gliricidia sepium*), Indian grass (*Panicum maximum*) and arboloco (*Montanoa quadrangularis*) in the feeding of guinea pigs (*Cavia porcellus*). Veterinaria y Zootecnia, 6: 56-65.
32. Wang, W.X., H.J. Yang, Y.K. Bo, S. Ding, B.H. Cao, 2012. Nutrient composition, polyphenolic contents and in situ protein degradation kinetics of leaves from three mulberry species. Livestock Science, 146: 203-206.
33. Doran, M.P., E.A. Laca and R.D. Sainz, 2007. Total tract and rumen digestibility of mulberry foliage (*Morus alba*), alfalfa hay and oat hay in sheep. Animal Feed Sciences and Technology, 138: 239-253.
34. Shayo, C.M., 1997. Uses yield and nutritive value of mulberry (*Morus alba*) trees for ruminants in the semi-arid areas of central Tanzania. Trop. Grasslands., 31: 599-604.
35. Bakshi, M.P.S. and W. Wadhwa, 2007. Tree leaves as complete feed for goat bucks. Sma Rum Res., 69: 74-78.
36. Hristov, A.N., K.L. Grandeen, J.K. Ropp, D. Greer, 2004. Effect of Yucca schidigera based surfactant on ammonia utilization *in vitro* and in situ degradability of corn grain. Animal Feed Science and Technology, 115: 341-355.
37. Griswold, K.E., G.A. Apgar, J. Bouton and J.L. Firkins, 2003. Effects of urea infusion and ruminal degradable protein concentration on microbial growth, digestibility and fermentation in continuous culture. J. Anim. Sci., 81: 329-336.
38. Meissner, H.H., M.O. Viljoen and W.A. van Niekerk, 1991. Intake and digestibility by sheep of Anthephora, Panicum, Rhodes and Smooth finger grass. In: Proc. 4th International Rangeland Congress, September 17th-21st, Montpellier, France.
39. Van Soest, P.J., 1994. Nutritional ecology of the ruminant. 2nd ed. Cornell University Press. pp: 476.
40. Promkot, C. and M. Wanapat, 2003. Ruminal degradation and intestinal digestion of crude protein of tropical protein resources using nylon bag technique and three-step in vitro procedure in dairy cattle; Livestock Research for Rural Development (15)11.

41. Belachew, Z., K. Yisehak, T. Taye and G.P.J. Janssens, 2013. Chemical Composition and *in Sacco* Ruminal Degradation of Tropical Trees Rich in Condensed Tannins. *Czech. J. Anim. Sci.*, 58: 176-192.
42. Lebopa, C.K., E.A. Boomker, M. Chimonyo and S.D. Mulugeta, 2011. Factors affecting the feeding behaviour of free ranging Tswana and Boer goats in the False Thornveld of the Eastern Cape, South Africa. *Life Science Journal*, 8: 70-80.
43. Vranic, M.K. Mladen, B. Kresimir, L. Josip, P. Goran, K. Hrvoje and M. Ivana, 2009. Maize silage supplementation to lower quality grass silage improves the intake, apparent digestibility and N retention. *Mljekarstvo*, 59: 302-310.
44. Lanyasunya, T.P., H.R. Wang, E.A. Mukisira, S.A. Abdulrazak and W.O. Ayako, 2006. Effect of seasonality on feed availability, quality and herd performance on smallholder farms in Ol-Joro-Orok location/Nyandarua district. *Tropical and Subtropical Agroecosystems*, 6: 87-93.
45. Suchitra, K. and M. Wanapat, 2008. Study on ruminal degradability of local plants by using nylon bag technique. *Livestock Research for Rural Development*. Volume 20, supplement. <http://www.lrrd.org/lrrd20/supplement/such1.htm>