

## Phosphorus-Zinc Interaction: Effects on Yield Components and Biochemical Composition and Bread Making Qualities of Wheat

L.K. Mishra and A.B. Abidi

Department of Biochemistry,  
ND University of Agriculture and Technology, Kumarganj, Faizabad-224229, India

**Abstract:** Field trials were conducted to assess the response of conjunctive use of phosphorus and zinc fertilizer on the 1000-seed weight and certain biochemical components of three varieties of wheat (*Triticum aestivum*) viz: PBW-343, HW-2425 and NW-1014. Results indicated that the phosphorus and zinc application had an synergistic impact on the 1000-seed weight and protein content of the wheat varieties. The increase in the doses of phosphorus and zinc had a positive impact on all the physical and biochemical characters of the wheat varieties. Application of phosphorus @ 60 kg/ha and zinc @ 10 kg/ha was found suitable to enhance the nutrient composition and certain bread making qualities of the wheat varieties.

**Key words:** Wheat • Bread making qualities • Biochemical composition • Phosphorus and zinc fertilization

### INTRODUCTION

The role of macro and micronutrients is crucial in crop nutrition and thus important for achieving higher yields. Nitrogen (N), phosphorus (P) and potassium (K), being primary essential nutrient, have prime importance in crop nutrition. Phosphorus (P) is involved in almost all biochemical pathways as a component part of energy carrier compounds, ATP and ADP. Phosphorus is one of the 17 essential elements required for plant growth and reproduction [1]. Phosphorus (P) is used in the plant for energy storage and transfer, maintenance and transfer of genetic code and is structural component of cells and many biochemicals. Phosphorus deficiencies results in poor root growth, stunted top growth, reduced yield and crop quality alongwith delayed maturity. Six micronutrients i.e., Mn, Fe, Cu, Zn, B and Mo are known to be required for all higher plants [2]. These have been well documented to be involved in photosynthesis, N-fixation, respiration and other biochemical pathways [1]. Approximately 60% of the world arable land is considered to be difficult for the plant production due to mineral stress caused by the deficiency, unavailability, or toxicity of some essential nutritive elements [3]. Of the microelements, Zn is thought to be the most widespread [4-6]. Zinc is an important essential element present in plant enzymatic systems. Genc *et al.* [7] reported that zinc has vast numbers of functions in plant metabolism and consequently zinc deficiency has a multitude of effects on

plant growth. Zinc deficiency is a worldwide nutritional constraint for crop production in many types of soil in the world [8, 9]. According to Graham and Welch [2] about 50% of the soil used for cereal production in the world contains a low level of plant available zinc which reduces not only grain yield but also nutritional quality (low in micronutrients essential for good human health). In general zinc application appears to improve the overall field performance of wheat plants.

Among the interactions involving macro and micronutrients the interaction between phosphorus and zinc is of greatest potential significance. This interaction is often negative (antagonistic) especially when a soil is deficient in both the nutrients but only one of them is applied through fertilizers. But it is quite possible to increase productivity in wheat through balanced use of phosphorus and zinc in such a way that they act synergistically. The impact of phosphorus and zinc together on the quality aspects and yield of wheat varieties has been studied in the present investigation.

### MATERIAL AND METHODS

**Experimental Design and Sowing Conditions:** The field trial was conducted to study the effect of zinc and phosphorus fertilization on the nutritional qualities and yield of wheat (*Triticum aestivum*). Three varieties commonly used in India i.e. PBW-343, HW-2425 and NW-1014 were used for the experiment purpose. The trial

was laid out in a split plot design with three phosphorus (control, 60 and 90 kg/hectare) and three zinc (Control, 5 and 10 kg/hectare) doses in the form of Single super phosphate and zinc sulphate respectively. Before fertilization pooled soil samples were analyzed for the available phosphorus and zinc (DTPA extractable) and it was 16.8 mg/kg and 0.5 mg/kg. The crop was sown in the month of December at a seed rate of 120 kg/hectare as per the recommended agronomic practices so as to raise a healthy and vibrant crop.

**Biochemical and Statistical Analyses:** The yield of the crop was obtained after harvest. The crop sample was obtained after the harvest and subjected to various biochemical analyses to ascertain the wheat quality. Bold healthy seeds were used to obtain the thousand seed weight. The sedimentation value and water absorption capacity were measured according to the method of The American Association of Cereal Chemistry [10]. The protein content in the samples was determined by the method of Lowry *et al.* [11]. The amino acid composition such as Tryptophan content, lysine and methionine content was determined by the method of Spies and Chamber [12], Felker *et al.* [13] and Horn *et al.* [14] respectively. The data obtained for different parameters was subjected to statistical analyses as per the method suggested by Gomez and Gomez [15] for split plot design.

## RESULTS

1000 kernel weight showed significant variation among different wheat varieties (Table 1). Thousand kernel weights ranged from 34.58 to 38.55 g among different wheat varieties. Significantly the highest 1000-kernel weight was observed in PBW-343 and the lowest in HW-2425. The application of phosphorus and zinc in increasing doses led to significant increase in the 1000-seed weight during both the years of the field experiment.

Maximum 1000-seed weight was recorded with the application of 90 kgP/ha and 10 kgZn/ha and it was statistically significant over all the other treatments. Statistically significant differences were recorded due to interaction between the phosphorus and zinc doses and phosphorus application @ 90 kg/ha in combination with zinc @ 10 kg/ha recorded significantly higher 1000-seed weight in both the years over all the treatments except phosphorus application @ 60 kg/ha along with zinc @ 10 kg/ha (Table 2).

The chemical composition of whole wheat flour such as protein content were significantly affected by the wheat varieties, phosphorus doses, zinc doses and interaction between phosphorus and zinc doses (Table 3). The amino acid composition of the whole wheat flour such as lysine, tryptophan and methionine content were significantly affected by phosphorus and zinc application during both the years of investigation (Table 5). The impact of varieties and interaction between the phosphorus and zinc doses was non significant.

Table 1: 1000-Seed weight of wheat as influenced by various treatments

Treatments	Test weight (In grams)	
	2003-04	2004-05
<b>Variety</b>		
V <sub>1</sub> (NW-1014)	37.38	37.18
V <sub>2</sub> (HW-2425)	34.78	34.58
V <sub>3</sub> (PBW-343)	38.55	38.35
CD (P = 0.05)	(1.53)	(1.52)
<b>Phosphorus levels</b>		
P <sub>0</sub> (0 kg/ha)	33.72	33.52
P <sub>1</sub> (60 kg/ha)	36.52	36.32
P <sub>2</sub> (90 kg/ha)	40.48	40.28
CD (P = 0.05)	(1.54)	(1.54)
<b>Zinc levels</b>		
Z <sub>0</sub> (0 kg/ha)	34.15	33.95
Z <sub>1</sub> (5 kg/ha)	37.67	37.47
Z <sub>2</sub> (10 kg/ha)	38.84	38.69
CD (P = 0.05)	(1.54)	(1.54)

Table 2: 1000-Seed weight of wheat as influenced by interaction between phosphorus and zinc levels

Zinc levels	Phosphorus levels					
	2003-04			2004-05		
	P <sub>0</sub> (control)	P <sub>1</sub> (60 kg/ha)	P <sub>2</sub> (90 kg/ha)	P <sub>0</sub> (control)	P <sub>1</sub> (60 kg/ha)	P <sub>2</sub> (90 kg/ha)
Z <sub>0</sub> (Control)	31.45	33.50	37.49	31.25	33.30	37.29
Z <sub>1</sub> (5 kg/ha)	34.14	37.16	41.72	33.94	36.96	41.52
Z <sub>2</sub> (10 kg/ha)	35.56	38.89	42.23	35.36	38.69	42.03
CD (P = 0.05)	2.677			2.677		

Table 3: Protein content in wheat as influenced by various treatments

Treatments	Protein content (In per cent)	
	2003-04	2004-05
Variety		
V <sub>1</sub> (NW-1014)	11.19	11.30
V <sub>2</sub> (HW-2425)	11.53	11.65
V <sub>3</sub> (PBW-343)	10.08	10.79
CD (P = 0.05)	(0.54)	(0.17)
Phosphorus levels		
P <sub>0</sub> (0 kg/ha)	10.81	10.92
P <sub>1</sub> (60 kg/ha)	11.16	11.27
P <sub>2</sub> (90 kg/ha)	11.44	11.55
CD (P = 0.05)	(0.44)	(0.44)
Zinc levels		
Z <sub>0</sub> (0 kg/ha)	10.39	10.49
Z <sub>1</sub> (5 kg/ha)	11.41	11.52
Z <sub>2</sub> (10 kg/ha)	11.61	11.73
CD (P = 0.05)	(0.44)	(0.44)

The sedimentation value and water absorption capacity were also significantly affected by varieties, doses of phosphorus and zinc (Table 6). The sedimentation value ranged from 31.53 to 36.21 while water absorption capacity was in the range of 65.62 to 72.48.

## DISCUSSION

The significant differences observed in 1000-kernel weight among wheat varieties may be due to the differences in the genetic make up of the varieties. However, these differences may be partly attributed due to different growing and environmental conditions prevailed during growing periods. The results are comparable with early findings of Finney *et al.* (1973) [16], Slaughter *et al.* (1992) [17] and Butt *et al.* (2001) [18]. The increase in 1000-seed weight due to increase in the doses of phosphorus may be due increase in the amount of available phosphorus during initial stage of growth when it is most needed. Thus, increasing doses of phosphorus

Table 4: Protein content in wheat as influenced by interaction between phosphorus and zinc levels

Zinc levels	Phosphorus levels					
	2003-04			2004-05		
	P <sub>0</sub> (control)	P <sub>1</sub> (60 kg/ha)	P <sub>2</sub> (90 kg/ha)	P <sub>0</sub> (control)	P <sub>1</sub> (60 kg/ha)	P <sub>2</sub> (90 kg/ha)
Z <sub>0</sub> (Control)	9.83	10.33	11.00	9.93	10.44	11.11
Z <sub>1</sub> (5 kg/ha)	11.11	11.51	11.60	11.22	11.63	11.72
Z <sub>2</sub> (10 kg/ha)	11.49	11.63	11.72	11.60	11.75	11.84
CD (P = 0.05)	0.775			0.767		

Table 5: Amino acid composition in wheat as influenced by various treatments (in per cent)

Treatments	Tryptophan (in per cent)		Methionine (in per cent)		Lysine (in per cent)	
	2003-04	2004-05	2003-04	2004-05	2003-04	2004-05
Variety						
V <sub>1</sub> (NW-1014)	2.24	2.22	1.44	1.43	2.20	2.22
V <sub>2</sub> (HW-2425)	2.30	2.27	1.99	1.97	2.08	2.02
V <sub>3</sub> (PBW-343)	2.26	2.24	1.82	1.80	2.40	2.42
CD (P = 0.05)	(NS)	(NS)	(NS)	(NS)	(NS)	(NS)
Phosphorus levels						
P <sub>0</sub> (0 kg/ha)	1.79	1.78	1.66	1.64	2.14	2.16
P <sub>1</sub> (60 kg/ha)	2.44	2.42	1.77	1.75	2.23	2.25
P <sub>2</sub> (90 kg/ha)	2.56	2.53	1.82	1.80	2.31	2.33
CD (P = 0.05)	(0.09)	(0.09)	(0.07)	(0.06)	(0.08)	(0.04)
Zinc levels						
Z <sub>0</sub> (0 kg/ha)	1.96	1.94	1.50	1.49	2.16	2.18
Z <sub>1</sub> (5 kg/ha)	2.33	2.31	1.84	1.82	2.24	2.26
Z <sub>2</sub> (10 kg/ha)	2.50	2.48	1.90	1.88	2.27	2.29
CD (P = 0.05)	(NS)	(NS)	(NS)	(NS)	(0.08)	(0.09)

Table 6: Sedimentation value and Water absorption capacity of wheat as influenced by various treatments

Treatments	Sedimentation (in ml)		Water absorption capacity (in ml)	
	2003-04	2004-05	2003-04	2004-05
<b>Variety</b>				
V <sub>1</sub> (NW-1014)	31.53	31.84	65.62	66.27
V <sub>2</sub> (HW-2425)	35.85	36.21	71.76	72.48
V <sub>3</sub> (PBW-343)	34.11	34.45	66.93	67.60
CD (P = 0.05)	(1.67)	(1.06)	(2.67)	(1.21)
<b>Phosphorus levels</b>				
P <sub>0</sub> (0 kg/ha)	32.86	33.19	66.28	66.95
P <sub>1</sub> (60 kg/ha)	33.77	34.10	67.90	68.57
P <sub>2</sub> (90 kg/ha)	34.86	35.21	70.13	70.83
CD (P = 0.05)	(1.33)	(1.34)	(0.92)	(2.66)
<b>Zinc levels</b>				
Z <sub>0</sub> (0 kg/ha)	33.62	33.95	67.72	68.39
Z <sub>1</sub> (5 kg/ha)	33.86	34.20	68.08	68.76
Z <sub>2</sub> (10 kg/ha)	34.01	34.35	68.51	69.20
CD (P = 0.05)	(1.33)	(1.34)	(0.92)	(2.66)

ensure that it is available in adequate amount and plays the significant role in the carbohydrate synthesis more efficiently. The findings are in agreement with the results obtained by Sharma and Bhardwaj [19].

The increasing trend in test weight due to increase in zinc doses during both the years of experiment may be due to the fact that application of zinc led to increase availability of zinc for plant growth in the zinc deficient local soil. Zinc is an important substrate involved in photo system-II of photosynthesis and plays vital role in energy metabolism process in plant. Thus, the increased availability and efficient absorption of zinc resulted in vibrant metabolism in wheat plant, which is an important reason for increase in test weight of the seeds. Similar results were obtained by Verma and Minhas [20].

The significant response observed due to interaction between the phosphorus and zinc doses may be due to the synergistic action of phosphorus and zinc doses. The synergistic action was mainly promoted due to the fact that the local soil was deficient in zinc and thus its application at the rate of 10 kg/ha could not trigger any antagonistic reaction as is normally witnessed when both these nutrients are applied in combination. The results are in conformity with the findings of Gattani *et al.* [21] and Nayak and Gupta [22].

The variation among the varieties with respect to the protein content may be attributed to the different genetic makeup of the varieties. The varieties having vigorous growth rates and efficient root systems could easily

adapt to the prevailing agro-climatic conditions. Thus, they could absorb the micro and macronutrients at much efficient rate and thereby initiate vibrant metabolism, which ultimately led to increase in protein content and other nutritional characters of wheat plant. The results of this investigation are comparable with early findings of Davis *et al.* [23], Tanija *et al.* [24]. The concomitant increase in the protein content of wheat with increase in dosage of phosphorus during both the years of cultivation might be due to the affect of increase in phosphorus concentration in soil and it is well known that phosphorus nutrition directly and indirectly influences wheat grain protein in many ways. The nutrient is required for absorption and assimilation of 'N' by wheat plants [25], for translocation of 'N' from vegetation to grain which might indirectly affect protein concentration. The results are in agreement with the findings of Mosolov and Volleidt [26]. The increase in the availability of zinc in zinc deficient soil led to increase in protein content of wheat. It is well known that zinc is actively involved in protein synthesis of plants. It is an important structural component of the protein synthesis machinery and is involved in the form of zinc mottifs in protein synthesis of plant. Similar, results have been reported by Jhaw [27]. The synergistic action between phosphorus and zinc may be the reason behind the increasing trend of protein content (Table 4). The deficiency of zinc in soil also might have played indirect role in increasing the protein content in wheat. Application of zinc led to synergistic action with phosphorus in the zinc deficient soil. The results are in conformity with the findings of Webb and Loneragan [28].

The increase in the amount of amino acids due the increase in the doses of phosphorus and zinc may be ascribed to the fact that both these minerals play a vital role in protein synthesis during the grain filling stages in the wheat crop and indirectly enhance the availability of nitrogen to the plants. Thus, increased efficiency of the protein synthesis is manifested in the increase in its structural components which include the amino acids such as lysine, tryptophan and methionine. Similar findings were reported by Lindsay [29], Alessendroni *et al.* [30] and Epperdorfer [31].

The varietal differences with respect to sedimentation value and water absorption capacity of wheat may be attributed to the fact that these varieties show significant variation among themselves in the context of gluten content. The sedimentation value and water absorption capacity of wheat is directly proportional to gluten content of wheat varieties. Therefore, it is natural that

they have variation in their sedimentation values and water absorption capacity also. Similar results have also been reported by Islam *et al.* [32].

The positive impact on sedimentation value and water absorption capacity due to increase in the doses of phosphorus and zinc may be due to the vital role played by these nutrients in increasing the gluten as well as the total protein content of the wheat. The findings of the present investigation are in conformity with the findings of Pratt [33] and Jyurg *et al.* [34].

The results of the present investigation clearly indicate that in zinc deficient soil the interaction between phosphorus and zinc is positive which is in contrary to established negative interaction between these two nutrients. It may also be concluded that these two nutrients play a very important role in improving the bread making qualities and the biochemical composition of the wheat varieties.

#### ACKNOWLEDGEMENT

The University scholarship received by the first author during the course of the present investigation is gratefully acknowledged.

#### REFERENCES

- Marschner, H., 1995. Mineral nutrition for Higher plants (2<sup>nd</sup> Edition). Academic Press Ltd, London, UK, pp: 229-312.
- Welch, R.M., W.H. Allaway, W.A. House and J. Kubota, 1991. Geographic distribution of trace element problems. In: Mortvedt J.J., ed. Micronutrients in agriculture, 2nd Ed. Madison, Wisconsin: SSSA Book Ser. 4. SSSA. pp: 31-57.
- Foy, C.D., 1983. Plant adaptation to mineral stress In problem soils. Iowa State J. Res., 57: 339-354.
- Graham, R.D., J.S. Ascher and S.C. Hynes, 1992. Selecting zinc efficient cereal genotypes for soils of low zinc status. Plant Soil, 146: 241-250.
- Yilmaz A., H. Ekiz, B. Torun, A. Aydin and I. Cakmak, 1995. Determination of zinc application methods in zinc-deficient wheat-growing areas of Central Anatolia. In: Soil fertility and fertilizer management. 9th Int. CIEC Symp. Proc. Kusadasi-Soke, Turkey, pp: 91-98.
- Cakmak I., M. Kalayci, H. Ekiz, H.J. Braun, Y. Kilinc and A. Yilmaz, 1999. Zinc deficiency as a practical problem in plant and human nutrition in Turkey: A NATO-science for stability project. Field Crop Res., 60: 175-188.
- Genc, Y., G.K. McDonald and R.D. Graham, 2006. Contribution of different mechanisms to zinc efficiency in bread wheat during early vegetative stage. Plant and Soil., 281: 353-367.
- Sillanpaa, M., 1982. Micronutrients and Nutrients Status of Soil: A Global Study. FAO Soils Bull No, 48. Rome.
- Rengel, Z. and R.D. Graham, 1995. Importance of seed Zn content for wheat growth on Zn-deficient soil. Plant and Soil., 173: 259-266.
- AOAC: Official Methods of Analysis. 1990. 15th Edn. Association of Official Analytical Chemists Washington, DC, USA.
- Lowry, O.H., N.J. Roserbrough, A.L. Farr and R.J. Randall, 1951. Protein Measurement with the Folin phenol reagent. J. Biol. Chem., 193: 265-275.
- Spies, J.T. and D.C. Chamber, 1949. Chemical determination of tryptophan in protein. Anatyl. Chem., 21: 1249.
- Felker, C., C.K. Libanauskas and G. Wainer, 1978. Estimation of lysine in foods. Crops Sci., 18: 480-490.
- Horn, J.M., D.B. Jones and A.E. Blum, 1946. Colorimetric determination of methionine in protein and foods. J. Biol. Chem., 1160-1173.
- Gomez, K.A. and A.A. Gomez, 1984. Statistical procedures for agricultural research Edn. 2<sup>nd</sup> John Willey and Sons, New York, pp: 680.
- Finney, P.L., G.S. Bains, R.C. Hosney and D.R. Lineback, 1973. Quality of Indian wheats. Cereal Sci. Today, 18: 392-7.
- Slaughter, C.D., H.N. Carl and R.H. William, 1992. Quality and classification of hard red wheat. Cereal Chem., 69: 428.
- Butt, M.S., F.M. Anjum, D.J. Van-Zuilichem and M. Shaheen, 2001. Development of predictive models for end use quality through canonical analysis. Int. J. Food Sci. Tech., 36: 433-40.
- Sharma, C.M. and S.K. Bhardwaj, 1998. Effect of phosphorus and zinc fertilization on yield and nutrient uptake in wheat (*Triticum aestivum* L.). Indian J. Agronomy, 43(3): 426-430.
- Verma, T.S. and R.S. Minhas, 1987. Zinc and phosphorus interaction in a wheat-maize cropping system, Fertilizer Res., 13: 77-86.
- Gattani, P.D., R.L. Jain and C.P. Vinayak, 1976. Zinc phosphorus interaction in wheat. J. the Indian Society of Soil Sci., 24(2): 208-210.
- Nayak, A.K. and M.L. Gupta, 1995. Phosphorus, zinc and organic matter interaction in relation to uptake, tissue concentration and absorption rate of phosphorus in wheat J. the Ind. Soc. of Soil Sci., 43(4): 633-636.

23. Davis, K.R., R.F. Cain, L.J. Peters, D. Le. Tourneau and J. McGiunis, 1981. Evaluation of the nutrient composition of wheat. In: Proximate analysis, B1, B2, B3 and pyridoxine. *Cereal Chem.*, 48: 116.
24. Tanija, S., K. Gupta, D.S. Wagle and K.S. Dhinda, 1983. Biological evaluation of wheat varieties. *J. Food Sci. Tech.*, 20: 319.
25. Harper, J.E. and G.M. Paulsen, 1969. Nitrate reductase activity in corn seedlings as affected by light and nitrate content in nutrient media. *Plant Physiol.*, 35: 700-708.
26. Mosolov, I.V. and L.P. Volleidt, 1962. Effect of doses and ratios of nitrogen and phosphorus on metabolism, yield and quality of spring wheat grain. *Plant Physiol.*, (U.S.S.R.) 9: 136-141.
27. Jhaw, G.C., 1991. Effect of different fertilizer strategies on yield of winter wheat. *Beijing Agric. Sci.*, 9(1): 24-29.
28. Webb, M.J. and J.F. Loneragan, 1990. Zinc translocation to wheat roots and its implications for a phosphorus. Zinc interaction in wheat plants. *J. Plant Nutr.*, 13: 1499-1512.
29. Lindsay, W.L., 1972. Zinc in soils and plant nutrition. *Adv. Agron.*, 24: 147-186.
30. Alessendroni, A.A., Baroccio and F. Incorvaia, 1971. Effect of cultivar, environment and nitrogen fertilizer on amino acid composition and grain protein in durum wheat. *Corenlicoltura*, 7: 105-125 (Abst. F.C.A., 32, 645).
31. Epperdorfer, W.H., 1978. Effect of N, P and K on amino acid composition and on relationship between nitrogen and amino acids in wheat and oat grains. *J. Fd. Agric. Sci.*, 29: 995-1001.
32. Islam, Q., F.M. Anjum, M.S. Butt and M. Hinnai, 1998. Suitability of Local Wheat Varieties for the Production of Pizza. *Pakistan J. Food Sci.*, 8: 8-11.
33. Pratt, D.B., Jr. 1971. Criteria of flour quality. *Wheat: Chemistry and Technology* IInd ed. (ed. Pomeranj, Y.). Amer Assoc. of Cereal Chemists, Incorporated. St. Paul, Minnersota. pp: 211-212.
34. Jyurg, W.H., A. Ehmann, K.K. Schlender and J. Scala, 1974. Differential response of two bean varieties to zinc as revealed by electrophoretic protein pattern. *Crop Sci.*, 12: 26-29.