

## The Effect of Menstrual Cycle on Ventilatory Equivalents in Two Mode of Exercise in Active Females

*Mahsa Mohsenzadeh*

Department of Physical Education and Sport Science.  
Islamic Azad University, Karaj Branch, Iran

**Abstract:** The purpose of this study was to determine the ventilatory responses during two incremental exercises in both the early follicular and luteal phases of the menstrual cycle in twenty healthy active females. Subjects were healthy women that were 20-25 years old with no known history of cardiopulmonary, metabolic, or musculoskeletal disease. Early follicular was in the fourth day of menstrual cycle and the luteal phase was determined by the level of Gnadotropin and Progesterone hormones in serum blood samples. The participants were randomly classified in two equal groups (Treadmill; T and Cycle Ergometer; CE). The participants performed an incremental exercise on a cycle ergometer and treadmill during their early follicular and luteal phases until they were exhausted. The pulmonary gas analyzer (K4B2) was applied to measure the minute ventilation ( $V_E$ ), Ventilatory equivalents for oxygen consumption ( $V_E/VO_2$ ) and Ventilatory equivalents for carbon dioxide production ( $V_E/VCO_2$ ) in early follicular and luteal phases of the menstrual cycle. Paired t-tests were performed to detect baseline differences across menstrual phases. The results indicated no significant differences ( $p > 0.05$ ) in  $V_E$ , ventilatory equivalents ( $V_E/VO_2$ ,  $V_E/VCO_2$ ) across the menstrual cycle of in active females. In conclusion, these results suggest that menstrual cycle hormones did not affect ventilatory responses at incremental exercise in active females. However, Ventilatory threshold during treadmill running was higher than cycle ergometer.

**Key words:** Ventilatory equivalents • Menstrual cyle • Mode of exercise • Incremental exercise

### INTRODUCTION

The effects of menstrual cycle hormones on exercise performance have been studied previously. However, the results remain controversial. The primary effects of estradiol and progesterone are related to reproductive behavior, but a number of reviews have addressed the role both progesterone and estradiol play in stimulating minute ventilation ( $V_E$ ) [1,2]. Exercise  $V_E$  plays a critical role in providing  $O_2$  to exercising muscles. However, exercise  $V_E$  is not considered to be a limiting factor during exercise, given that normal healthy individuals rarely approach mechanical or diffusion limitations even at maximal exercise intensities [3]. Particularly, the respiratory responses to progressive intensity exercise have shown inconsistent results. Several studies show higher exercise minute ventilation ( $V_E$ ) during the luteal phase compared to the follicular phase [4-7] while other studies show no difference [8-11]. The effect of cyclic

variations in ovarian hormones on exercise  $V_E$  continues to be unclear. The luteal and the follicular phase have shown differences for maximal  $V_E$  as high as 12 L/min with no effect on  $VO_{2max}$  [6,12]. Numerous studies have also suggested the impact of menstrual cycle hormones on  $VO_{2max}$  or exercise performance appears to be minimal [6,9-15]. There are inconsistent reports in the literature on the effect of sex on the ventilatory equivalents for oxygen ( $V_E/VO_2$ ) and carbon dioxide ( $V_E/VCO_2$ ) during exercise. Some have demonstrated no differences [6,16] and others have shown a significantly higher  $V_E/VCO_2$  in women compared to men [17]. Neither the Blackie *et al.* (1991) nor the Habedank *et al.* (1998) studies accounted for menstrual cycle phase [16,17]. The differences in  $V_E/VCO_2$  can be greater during the luteal phase [11]; however, this is also controversial [12]. De Souza *et al.* (1990) observed no differences in oxygen uptake,  $V_E$ , heart rate, respiratory exchange ratio (RER), rating of perceived exertion (RPE), time to fatigue and plasma lactate

following maximal and sub maximal exercise tests between the follicular and luteal phases in eumenorrheic and amenorrheic runners [10]. Also, the results suggest that neither the follicular or luteal phases alter exercise performance in female athletes [10]. The  $V_E/VCO_2$  relationship can be stated as the absolute change in  $V_E$  per unit change in  $VCO_2$  or as the y-intercept and has been used to evaluate the appropriateness (efficiency) of minute ventilation during exercise. There are inconsistent reports of the effect of gender on the exercise  $V_E/VCO_2$  where some have demonstrated no differences [16] and others have shown a significantly higher  $V_E/VCO_2$  in women compared to men [17]. However, neither of the two studies accounted for the menstrual cycle phase and the differences can possibly be greater throughout the luteal phase [6,11]. Furthermore, Schoene *et al.* (1981) has demonstrated that  $V_E/VO_2$  during progressive exercise on a bicycle ergometer was significantly increased throughout the luteal phase compared to the follicular phase [5]. In particular, the ventilatory responses to progressive intensity exercise have shown mixed results. It is thought that the higher levels of circulating estrogen and progesterone are the stimuli for altered ventilatory responses during the luteal phase of the cycle. The purpose of this investigation was to compare different mode of testing ventilatory threshold in the early follicular and luteal phases of menstrual cycle of inactive women. The results of the study could enhance our understanding of the hormonal influences on breathing during exercises that occur during the menstrual cycle. The results could have an impact on exercise rehabilitation programs for patient populations, exercise prescription for disease prevention in healthy individuals and training strategies for competitive athletes.

**MATERIAL AND METHODS**

Twenty healthy, active Females between the age of 19 and 25 yr with no known history of cardiopulmonary, metabolic or musculoskeletal disease were recruited in this study. All subjects demonstrated a normal menstrual cycle during the previous 4 months as determined by the medical history form and were not currently using birth control medications. The participants included 20 basketball players who were students of Islamic Azad University Central Branch. Subjects were randomly classified in two equal groups (Treadmill; T and Cycle Ergometer; CE). The age, weight and height of

participants were respectively 22±3 yrs, (52.31±2.44) kg and (165 ±4.50) cm in T: active group and 22±3 yrs, (53.43±4.19) kg and (165±3.77) cm in C: active group.

Early follicular was in the fourth day of menstrual cycle and the luteal phase was determined by the level of Progesterone, Prolactin, FSH and LH hormones in serum blood samples(RIA-method). An incremental graded exercise test (GXT) was conducted on a cycle ergo meter (Techno Gym) and Treadmill (Cosmed-Italy). On the cycle ergometer: After a 5-min warm up at 50 W, the subjects rode a progressive exercise test (25 W/min), until they were unable to continue [18]. The pulmonary gas analyzer (K4B2-Casmed) was applied to measure the, minute ventilation ( $V_E$ ), ventilatory equivalents ( $V_E/VO_2$ ,  $V_E/VCO_2$ ) in two phases. All testing was conducted in accordance with the guidelines of the American College of Sports Medicine (2000) in the measurement center of Olympic committee [19]. Paired T-test was used to determine the differences in minute ventilatory, ventilatory equivalents  $V_E/VO_2$  and  $V_E/VCO_2$  during the follicular and luteal phases at incremental exercise. Significance was set at the 0.05 level. All statistical analysis was performed utilizing SPSS, version 13.

**RESULTS**

The results indicated no significant differences in maximal  $V_E$  across the menstrual cycle ( $p > 0.05$ ). The ventilatory equivalents  $V_E/VO_2$  and  $V_E/VCO_2$  showed no significant differences ( $p > 0.05$ ) across the menstrual cycle, but in T: group higher than C: group.  $V_E/VO_2$  at maximal exercise for the follicular and luteal phases in T: group was 44.98 ± 4.67 and 45.09 ± 4.53, C: group was 39.89 ± 4.87 and 40.09 ± 4.17 respectively (Table1).  $V_E/VCO_2$  at maximal exercise for the follicular and luteal phases in T: active group was 34.56 ± 4.50 and 34.75 ± 3.64, C: group was 31.02 ± 4.05 and 30.22 ± 3.56 respectively (Table1).

Table 1: Maximal Exercise Test Variables

Groups	Variable	Follicular	LutealTreadmill
	$VE(L/min)$	91.19±6.87	97.67±14.89
	$VE/VO_2$	44.98 ± 4.67	45.09 ± 4.53
	$VE/VCO_2$	34.56 ± 4.50	34.75 ± 3.64
Cycle Ergometer	$VE(L/min)$	73.65±18.50	71.15±19.27
	$VE/VO_2$	39.89 ± 4.87	40.09 ± 4.17
	$VE/VCO_2$	31.02 ± 4.05	30.22 ± 3.56




Fig. 1: Minute Ventilation (VE) at incremental exercise during the follicular and luteal phases of the menstrual cycle. Values are means + SD.




Fig. 2: Ventilatory equivalents for oxygen consumption (VE/VO<sub>2</sub>) at incremental exercise during the follicular and luteal phases of the menstrual cycle. Values are means + SD.




Fig. 3: Ventilatory equivalents for carbon dioxide production (VE/VCO<sub>2</sub>) at incremental exercise during the follicular and luteal phases of the menstrual cycle. Values are means + SD.

### CONCLUSION

The major finding of this study suggests that menstrual hormones did not affect breathing responses at maximal exercise. Additionally, there were no significant differences in gas exchange variables during a maximal graded exercise test between the follicular and luteal phases of the menstrual cycle. Menstrual cycle hormones did not affect respiratory stimulation across the menstrual cycle at maximal  $V_E$ . Our results are in agreement with Beidleman *et al.* (1999), Casazza *et al.* (2002), Dean *et al.* (2003), Bembem *et al.* (1995), De Souza *et al.* (1990), Hackney *et al.* (1991), Stephenson *et al.* (1982) and Lebrun *et al.* (1995) [8,9,10,14,15,20,21]. The Beidleman *et al.* (1999) study showed that peak  $V_E$  and sub maximal  $V_E$  were not affected by menstrual cycle phase [12].

However, other studies have shown significant differences in exercise  $V_E$  during the luteal phase of the menstrual cycle [4-7]. The varying results between these studies and the current study could be related to individual responses and differences in progesterone receptor sensitivity [9]. In addition, it is widely accepted that ventilatory measures demonstrate large within-subject daily variability. The ventilatory equivalents ( $V_E/VO_2$  and  $V_E/VCO_2$ ) were not elevated during the luteal phase in the incremental exercise compared to the follicular phase. Our study is one of the few studies to account for menstrual cycle phase for these variables. Beidleman *et al.* (1999) and Bembem *et al.* (1995) had similar findings that showed no significant differences ( $p > 0.05$ ) in the ventilatory equivalents in the follicular and luteal phases of the menstrual cycle [8-12]. Nonetheless, Shoene *et al.* (1981) results showed  $V_E/VO_2$  was significantly greater at all levels of exercise in the luteal phase [5]. However, the study by Schoene *et al.* (1981) did not show a significant correlation between respiratory variables and plasma progesterone [5]. It is not surprising that  $V_E/VO_2$  and  $V_E/VCO_2$  were not elevated in the current study as neither  $V_E$ ,  $VO_2$  nor  $VCO_2$  were different between menstrual phases at maximal level of exercise. Controversy regarding the effect of ovarian hormones on ventilatory chemosensitivity is probably due to 1) a wide range of estradiol and progesterone among subjects in the same study, as well as between studies, 2) individual responsiveness to a given ovarian hormone level and 3) the relatively large within-subject, between-day variability inherent in measures of ventilator chemosensitivity [22]. Additionally, the differences between average of ventilator equivalent in two modes of exercise were significant in both groups of participants and, it was more in T than CE ( $P < 0.05$ ). The greater slow component observed in cycling compared to running may be related to differences in the muscle contraction regimen that is required for the two exercise modes. It seems that because of the physical depression during menstrual cycle, which is due to the female untrained, the replacement of running with cycling, with the aim of making physical activity simpler and encouraging female to continue their activity during this time, is not recommendable. In conclusion, these results suggest that the menstrual cycle phase did not affect breathing responses at maximal exercise. Additionally, there are no differences in maximal exercise test variables between the follicular and luteal phases of the menstrual cycle. These results provide additional data suggesting that the timing of the menstrual cycle phase may not be as critical as once thought when designing future exercise ventilation

studies. The results of the study could enhance our understanding of the hormonal influences on breathing during exercise that occur during the menstrual cycle. The results could have an impact on exercise rehabilitation programs for patient populations, exercise prescription for disease prevention in healthy individuals and training strategies for competitive athletes.

### REFERENCES

1. Bayliss, D.A. and D.E. Millhorn, 1992. "Central neural mechanisms of progesterone action: application to the respiratory system". *J. Applied Physiol.*, 73: 393-404.
2. Lyons, H.A., 1976. "Centrally acting hormones and respiration". *Pharmacol. Ther.*, B2: 743-751.
3. Dempsey, J.A., 1986. "Is the lung built for exercise?". *Med. Sci. Sports Exerc.*, 18: 143-155.
4. Jurkowski, J.E., N.L. Jones, C.J. Toews and J.R. Sutton, 1981. "Effects of menstrual cycle on blood lactate, O<sub>2</sub> delivery and performance during exercise". *J. Applied Physiol.*, 51: 1493-99.
5. Schoene, R.B., H.T. Robertson, D.J. Pierson and A.P. Peterson, 1981. "Respiratory drives and exercise in menstrual cycles of athletic and nonathletic women". *J. Applied Physiol.*, 50: 1300-1305.
6. Sheel, A.W., J.C. Richards, G.E. Foster and J.A. Guenette, 2004. "Sex differences in respiratory exercise physiology". *Sports Med.*, 34: 567-579.
7. William, T.J. and G.S. Krahenbuhl, 1997. "Menstrual cycle phase and running economy". *Medicine Science Sports Exercise*. 29: 1609-1618.
8. Bembien, D.A., P.C. Salm and A.J. Salm, 1995. "Ventilatory and blood lactate responses to maximal treadmill exercise during the menstrual cycle". *J. Sports Medicine and Physical Fitness*. 35: 257-262.
9. Casazza, G., S. SangHoon, B. Miller, F. Navazio and Brooks, G. 2002. "Effects of oral contraceptives on peak exercise capacity". *J. Applied Physiol.*, 93: 1698-1702.
10. De Souza, M.J., M.S. Maguire and K.R. Rubin, 1990. "Effects of menstrual phase and amenorrhea on exercise performance in runners". *Medicine Science Sports Exercise*. 22: 575-580.
11. Dombrov, M.L., H.W. Bonekat and T.J. Williams, 1987. "Exercise performance and ventilatory response in the menstrual cycle". *Medicine Science Sports Exercise*. 19: 111-117. 1987.
12. Beidleman, B.A., P.B. Rock, S.R. Muza, C.S. Fulco, V.A. Forte and A. Cymerman, 1999. "Exercise VE and physical performance at altitude are not affected by menstrual cycle phase". *J. Applied Physiol.*, 86: 1519-1526.
13. Bryner, R.W., R.C. Toffle and I.H. Ullrich, 1990. "Effect of low dose oral contraceptives on exercise performance". *British J. Sports Med.*, 30: 36-40.
14. Hackney, A.C., C.S. Curley and B.J. Nicklas, 1991. "Physiological responses to submaximal exercise at the mid-follicular, ovulatory and mid-luteal phases of the menstrual cycle". *Scand J. Medicine Science Sports*. 1: 94-98.
15. Lebrun, C.M., D.C. McKenzie and J.C. Prior, 1995. "Effects of menstrual cycle phase on athletic performance". *Medicine Science Sports Exercise*. 27: 437-444.
16. Blackie, S.P., M.S. Fairbairn and N.G. McElvaney, 1991. "Normal values and ranges for ventilation and breathing pattern at maximal exercise". *Chest*. 100: 136-142.
17. Habedank, D., I. Reindl and G. Vietzke, 1998. "Ventilatory efficiency and exercise tolerance in 101 healthy volunteers". *European Journal of Applied Physiology Occup. Physiol.*, 77: 421-426.
18. Hopkins, S.R., R.C. Barker, T.D. Brutsaert, T.P. Gavin, P. Entin, I.M. Olfert, S. Veisel and P.D. Wagner, 2000. "Pulmonary gas exchange during exercise in women: effects of exercise type and work increment". *J. Appl. Physiol.*, 89: 721-730.
19. American College of Sports Medicine, Guidelines for Exercise Testing and Prescription (2000). (6<sup>th</sup> Eds). Chapter 4, 57-90. Lippincott, Williams and Wilkins: Philadelphia, PA.
20. Dean, T.M., L. Perreault, R.S. Mazzeo and T.J. Horton, 2003. "No effect of menstrual cycle phase on lactate threshold". *J. Applied Physiol.*, 95: 2537-2543.
21. Stephenson, L.A., M.A. Kolka and J.E. Wilkerson, 1982. "Metabolic and thermoregulatory responses to exercise during the human menstrual cycle". *Med. Sci. Sports Exerc.*, 14: 270-275.
22. Sahn, S.A., C.W. Zwillich, N. Dick, R.E. McCullough, S. Lakshminarayan and J.V. Weil, 1977. "Variability of ventilator responses to hypoxia and hypercapnia". *J. Applied Physiol.*, 43: 1019-1025.