

## Trigonometric Curve Fitting Based on Multi-step Differential Transformation Method and the Application of Non-linear Oscillatory Systems

<sup>1</sup>Ahmet Gökdoğan, <sup>1</sup>Mehmet Merdan and <sup>2</sup>Syed Tauseef Mohyud-Din

<sup>1</sup>Department of Mathematics Engineering, Gümüşhane University, 29100-Gümüşhane, Turkey

<sup>2</sup>HITEC University Taxila Cantt, Pakistan

**Abstract:** In this article, trigonometric curve fitting on the multi-step differential transformation method (MsDTM) is implemented to give approximate analytical solutions of nonlinear ordinary differential equation such as non-linear oscillation systems. In addition, the solution obtained from MsDTM applied trigonometric curve fitting. Our proposed approach reveals that results to approximate analytical solutions of non-linear oscillation systems. Some plots are presented to show the reliability and simplicity of the methods.

**Key words:** Curve fitting . the multi-step differential transformation method . non-linear oscillatory systems

### INTRODUCTION

This study will consider the following general non-linear oscillations [1-4]:

$$y'' + w_0^2 y + \lambda g(y) = f(t) \quad (1)$$

with initial conditions

$$y(0) = A, y'(0) = 0 \quad (2)$$

Here  $g$  is a nonlinear function depending on  $y$  and its derivatives.

A review of the recently developed analytical methods is given in review article and the comprehensive book by He [5, 6]. In the last decade, several computational methods have been applied to solve nonlinear oscillation systems. Some of these well-known methods are such as: variational iteration method [7], homotopy perturbation method [8], homotopy analysis method [9], Exp-function method [10, 11], linearized perturbation method [12], modified Lindstedt-Poincare methods [13], iteration perturbation method [14], bookkeeping parameter perturbation method [15], energy balance method [16], amplitude frequency formulation [17], max-min approach [18, 19], rational harmonic balance method [20], Mickens iteration procedure [21].

The differential transform method (DTM) is a numerical as well as analytical method for solving integral equations, ordinary, partial differential

equations and differential equation systems. The concept of the differential transform was first proposed by Zhou [22] and its main application concern with both linear and nonlinear initial value problems in electrical circuit analysis. The DTM gives exact values of the  $n$ th derivative of an analytic function at a point in terms of known and unknown boundary conditions in a fast manner. This method constructs, for differential equations, an analytical solution in the form of a polynomial. It is different from the traditional high order Taylor series method, which requires symbolic computations of the necessary derivatives of the data functions. The Taylor series method is computationally taken long time for large orders. The DTM is an iterative procedure for obtaining analytic Taylor series solutions of differential equations. Different applications of DTM can be found in [23-26].

The aim of this Letter is to apply DTM and multi-stage DTM [27] to solve the non-linear oscillator equation. The numerical solutions are compared with the multi-stage DTM and by classical RK4.

This paper is organized as follows:

In section 2, 3 and 4, we describe DTM and multi-stage DTM briefly. To show in efficiency of this method, we give some examples and numerical results in Section 5. The conclusions are then given in the final section 6.

### SOLUTION APPROACHES

Consider a general equation of  $n$ -th order ODE

$$f(t, y, y', \dots, y^{(n)}) = 0 \quad (3)$$

subject to the initial conditions

$$y^{(k)}(0) = d_k, k = 0 \dots n-1 \quad (4)$$

system in (3), we will next present the solution approaches of (3) base on standard DTM and MsDTM separately.

### SOLUTION OF BY DTM

To illustrate the differential transformation method (DTM) for solving differential equations systems, the basic definitions of differential transformation are introduced as follows. Let  $y(t)$  be analytic in a domain  $D$  and let  $t = t_i$  represent any point in  $D$ . The function  $y(t)$  is then represented by one power series whose center is located at  $t_i$ . The differential transformation of the  $k$ th derivative of a function  $y(t)$  is defined as follows:

$$Y(k) = \frac{1}{k!} \left[ \frac{d^k y(t)}{dt^k} \right]_{t=t_i}, \forall t \in D \quad (5)$$

In (5),  $y(t)$  is the original function and  $Y(K)$  is the transformed function. As in [22-28] the differential inverse transformation of  $Y(K)$  is defined as follows:

$$y(t) = \sum_{k=0}^{\infty} Y(k) (t - t_i)^k, \forall t \in D \quad (6)$$

In fact, from (5) and (6), we obtain

$$y(t) = \sum_{k=0}^{\infty} \frac{(t - t_i)^k}{k!} \left[ \frac{d^k y(t)}{dt^k} \right]_{t=t_i}, \forall t \in D \quad (7)$$

Eq. (7) implies that the concept of differential transformation is derived from the Taylor series expansion. From the definitions of (5) and (6), it is easy to prove that the transformed functions comply with the following basic mathematics operations (Table 1)

In real applications, the function  $y(t)$  is expressed by a finite series and (5) can be written as

$$y(t) = \sum_{k=0}^N Y(k) (t - t_i)^k, \forall t \in D \quad (8)$$

Equation (8) implies that  $\sum_{k=N+1}^{\infty} Y(k) (t - t_i)^k$  is negligibly small.

Table 1: Operations of the one dimensional differential transform

Original function	Transformed function
$y(t) = f(t) \mp g(t)$	$Y(k) = F(k) \mp G(k)$
$y(t) = \xi f(t)$	$Y(k) = \xi F(k)$
$y(t) = \frac{df(t)}{dt}$	$Y(k) = (k+1) F(k+1)$
$y(t) = \frac{d^n f(t)}{dt^n}$	$Y(k) = \frac{(k+n)!}{k!} F(k+n)$
$y(t) = e^{\lambda t}$	$Y(k) = \frac{\lambda^k}{k!}$
$y(t) = \sin(wt + \phi)$	$Y(k) = \frac{w^k}{k!} \sin\left(\frac{k\pi}{2} + \phi\right)$
$y(t) = \cos(wt + \phi)$	$Y(k) = \frac{w^k}{k!} \cos\left(\frac{k\pi}{2} + \phi\right)$
$y(t) = \int_0^t f(t) dt$	$Y(k) = \frac{F(k-1)}{k}$
$y(t) = t^r$	$Y(k) = \delta(k-r) = \begin{cases} 1, & k=r, \\ 0, & \text{otherwise} \end{cases}$
$y(t) = f(t)g(t)$	$Y(k) = \sum_{k_1=0}^k F(k_1)G(k-k_1)$
$y(t) = f(t)g(t)h(t)$	$Y(k) = \sum_{k_2=0}^k \sum_{k_1=0}^{k_2} F(k_1)G(k_2-k_1)H(k-k_2)$

### SOLUTIONS BY MSDTM

Let  $[0, T]$  be the interval over which we want to find the solution of the initial value problem (3). In actual applications of the DTM, the approximate solution of the initial value problem (3)-(4) can be expressed by the finite series,

$$y(t) = \sum_{i=0}^N b_i t^i, t \in [0, T] \quad (9)$$

Assume that the interval  $[0, T]$  is divided into  $N$  subintervals  $[t_{n-1}, t_n]$ ,  $n = 1, 2, \dots, N$  of equal step size  $h = T/N$  by using the nodes  $t_n = nh$ . The main ideas of the multi-step DTM are as follows [43]. First, we apply the DTM to Eq. (3) over the interval  $[0, t_1]$ , we will obtain the following approximate solution,

$$y_1(t) = \sum_{i=0}^N b_i t^i, t \in [0, t_1] \quad (10)$$

using the initial conditions  $y_1^{(k)}(0) = d_k$ . For  $n \geq 2$  and at each subinterval  $[t_{n-1}, t_n]$ , we will use the initial conditions  $y_n^{(k)}(t_{n-1}) = y_{n-1}^{(k)}(t_{n-1})$  and apply the DTM to Eq. (3) over the interval  $[t_{n-1}, t_n]$ , where  $t_i$  in Eq. (3) is

replaced by  $t_{n-1}$ . The process is repeated and generates a sequence of approximate solutions  $y_n(t)$ ,  $n = 1, 2, \dots, N$  for the solution  $y(t)$ ,

$$y_n(t) = \sum_{i=0}^N b_{ni}(t - t_{n-1})^i, \quad t \in [t_{n-1}, t_{n+1}] \quad (11)$$

In fact, the multi-step DTM assumes the following solution,

$$y(t) = \begin{cases} y_1(t), & t \in [0, t_1] \\ y_2(t), & t \in [t_1, t_2] \\ \vdots \\ y_N(t), & t \in [t_{N-1}, t_N] \end{cases} \quad (12)$$

### TRIGONOMETRIC CURVE FITTING

We consider data obtained from the multi-step DTM. The problem of data fitting consist of finding a function  $f$  that “best” represents the data which are subject to errors. A reasonable way to approach the problem is to plot data points in an  $xy$ -plane and try to recognize the shape of a guess function  $f(t)$ . Trigonometric fitting is a method to approximate a function  $f$  by series of trigonometric functions. Trigonometric functions in general form can be written.

$$\begin{aligned} f(t) = & A_1 \cos(w_1 t + \varphi_1) + A_2 \cos(w_2 t + \varphi_2) \\ & + \dots + A_i \cos(w_i t + \varphi_i) + \dots + D \end{aligned} \quad (13)$$

$(i = 1, 2, \dots, n)$

As a function approximately describing the data, in which

$$M = (A_1, w_1, \varphi_1, A_2, w_2, \varphi_2, \dots, A_i, w_i, \varphi_i, \dots, D)^T$$

is the unknown coefficients. Set  $y_i$  is the found value at the point  $x_i$  ( $i = 1, 2, \dots, n$ ),  $y'_i$  is the found value at the point  $x_i$ , is the result of the calculation by the fitting function at point, the mean square error of the  $n$  data points

We will apply classic DTM and the multistage DTM to nonlinear ordinary differential Eq. (15). Applying classic DTM for Eq. (15)

$$\begin{aligned} (k+1)Y(k+1) = & Z(k)(k+1)Z(k+1) + \frac{3}{4} \sum_{k_2=0}^k \sum_{k_1=0}^{k_2} (k-k_2+1)Y(k_1)Y(k_2-k_1)Z(k-k_2+1) \\ & + \frac{3}{4} \sum_{k_2=0}^k \sum_{k_1=0}^{k_2} Y(k_1)Z(k_2-k_1)Z(k-k_2) + \frac{3g}{1} \sum_{k_1=0}^k Y(k_1)G(k-k_1) = 0, Y(0) = A, Z(0) = 0 \end{aligned} \quad (16)$$

Trigonometric nonlinearity:

$$\delta = \sqrt{\sum_{i=1}^n (y_i - y'_i)^2}$$

obviously the smaller the value of this function, the better results are. Under general conditions, according to the properties of boundary of trigonometric functions, the correlation between the maximum pure trigonometric function ( $A$ ), the maximum(max) and minimum(min) of data is  $A \leq (\max + \min)/2$ ; as the periodic characteristics of the trigonometric function (period  $2\pi$ ) the  $w_i t + \varphi_i$  data will repeat every  $2\pi$  every  $\pi$  positive and negative values will be changed. So,  $w_i t$  and  $\varphi_i$  's absolutely value should be less than  $\pi$ , considering the accuracy, the minimum spacing between numerical as  $T$  is determined to be maximum absolute value of  $w_i$  for the  $\pi/T$ . D range of value;  $\min \leq D \leq \max$  [49, 52-56].

### APPLICATIONS

**Example 1:** (The motion of a rigid rod rocking back). In this section, we consider a complicated and practical case of the motion of a rigid rod rocking back and forth on the circular surface without slipping [50] as follow:

$$\begin{aligned} \frac{d^2 y}{dt^2} + \frac{3}{4} y^2 \frac{d^2 y}{dt^2} + \frac{3}{4} y \left( \frac{dy}{dt} \right)^2 + \frac{3g \cos(y)}{1} &= 0 \\ y(0) = A, y'(0) &= 0 \end{aligned} \quad (14)$$

where  $A$  represents the amplitude of the oscillation. Motion is assumed to start from the position of maximum displacement with zero initial velocity. Eq. (14) can be expressed in the form of two simultaneous first-order differential equations in terms of  $y(t)$  and  $z(t)$ , i.e.

$$\begin{aligned} \frac{dy}{dt} &= z \\ \frac{dz}{dt} + \frac{3}{4} y^2 \frac{dz}{dt} + \frac{3}{4} y z^2 + \frac{3g \cos(y)}{1} &= 0 \\ y(0) = A, z(0) &= 0 \end{aligned} \quad (15)$$

$$f(x) = \sin(x), g(x) = \cos(x)$$

Similarly, replacing  $k-1$  by  $k$  gives:

By definition,

$$\begin{aligned} F(0) &= [\sin(x(t))]_{t=0} = \sin(x(0)) = \sin(X(0)) \\ G(0) &= [\cos(x(t))]_{t=0} = \cos(x(0)) = \cos(X(0)) \end{aligned} \quad (17)$$

$$\begin{aligned} F(k) &= \sum_{k_1=0}^{k-1} \left( \frac{k-k_1}{k} \right) G(k_1) X(k-k_1), \quad k \geq 1 \\ G(k) &= - \sum_{k_1=0}^{k-1} \left( \frac{k-k_1}{k} \right) F(k_1) X(k-k_1), \quad k \geq 1 \end{aligned} \quad (20)$$

To find other transformed functions, we differentiate

Combine Eqs.(17) and (20) to give the recursive relation:

$$f(x) = \sin(x), g(x) = \cos(x)$$

obtaining:

$$F(k) = \begin{cases} \sin(X(0)), & k=0 \\ \sum_{k_1=0}^{k-1} \left( \frac{k-k_1}{k} \right) G(k_1) X(k-k_1), & k \geq 1 \end{cases} \quad (21)$$

$$\begin{aligned} \frac{df(x)}{dt} &= \cos(x) \left( \frac{dx(t)}{dt} \right) = g(x) \left( \frac{dx(t)}{dt} \right) \\ \frac{dg(x)}{dt} &= -\sin(x) \left( \frac{dx(t)}{dt} \right) = -f(x) \left( \frac{dx(t)}{dt} \right) \end{aligned} \quad (18) \quad \text{and}$$

$$G(k) = \begin{cases} \cos(X(0)), & k=0 \\ - \sum_{k_1=0}^{k-1} \left( \frac{k-k_1}{k} \right) F(k_1) X(k-k_1), & k \geq 1 \end{cases} \quad (22)$$

Applying the differential transform to Eq. (18) obtain:

$$\begin{aligned} (k+1)F(k+1) &= \sum_{k_1=0}^k (k+1-k_1)G(k_1)X(k+1-k_1) \\ (k+1)G(k+1) &= - \sum_{k_1=0}^k (k+1-k_1)F(k_1)X(k+1-k_1) \end{aligned} \quad (19)$$

are given by [48].

where  $Y_i(n)$ , for  $n = 1 \dots M$ , satisfy the following recurrence relations,

$$\begin{aligned} (k+1)Y_i(k+1) &= Z(k)(k+1)Z_i(k+1) + \frac{3}{4} \sum_{k_2=0}^k \sum_{k_1=0}^{k_2} (k-k_2+1)Y_i(k_1)Y_i(k_2-k_1)Z_i(k-k_2+1) \\ &+ \frac{3}{4} \sum_{k_2=0}^k \sum_{k_1=0}^{k_2} Y_i(k_1)Z_i(k_2-k_1)Z_i(k-k_2) + \frac{3g}{1} \sum_{k_1=0}^k Y_i(k_1)G(k-k_1) = 0 \\ Y_0(0) &= A, Y_0(1) = 0, Y_{i+1}(0) = Y_i(t^*), Y_{i+1}(1) = Y_i(t^*), t \in [t_p, t_{p+1}], t^* = t_i, i = 0, \dots, M-1 \end{aligned} \quad (23)$$

By applying the multistage DTM to Eq.(15) is obtained Eq.(24) as following:

$$y(t) = \begin{cases} \sum_{k=0}^N Y_0(k)t^k, t \in [t_0, t_1] \\ \sum_{k=0}^N Y_1(k)t^k, t \in [t_p, t_2] \\ \vdots \\ \sum_{k=0}^N Y_M(k)t^k, t \in [t_{M-1}, t_M] \end{cases} \quad (24)$$

On the function (13) taken  $n = 1$ ,  $t \in [0, 7]$ ,  $g = 1 = 1$ , solution of curve fitting

$$y_{cf}(t) = 0.15737\cos(1.715664t + 0.0003702) - 0.14184e^{-4}$$

is obtained and presented in the following in Fig. 1. RK4 method is also compared with trigonometric curve fitting solution.

In (13) expression,  $n = 2$ ,  $t \in [0, 8]$  and  $g = 1 = 1$  are taken, solution of

$$y_{cf}(t) = 0.47835\cos(1.5939t + 0.3643e^{-4}) + 0.7651e^{-3}\cos(6.6469t - 28.6595) - 0.20276e^{-3}$$




Fig. 1: Plots of (a) displacement  $y$  versus time  $t$  and (b) phase portrait for  $A = 0.05\pi$




Fig. 2: Plots of (a) displacement  $y$  versus time  $t$  and (b) phase portrait for  $A = 0.05\pi$




Fig. 3: Plots of (a) displacement  $y$  versus time  $t$  and (b) phase portrait for  $A = 0.05\pi$

is obtained and presented in the following in Fig. 2. RK4 method is also compared with trigonometric curve fitting solution.

In (13) equation,  $n = 3$ ,  $t \in [0, 9]$  and  $g = 1 = 1$  are taken, solution of

$$y_{cf}(t) = 0.4781\cos(1.594t - 0.00557) + 79.95\cos(0.000701t + 3.143) + 0.00228\cos(0.973t - 2.961) + 79.95$$

is obtained and presented in the following in Fig. 3. RK4 method is also compared with trigonometric curve fitting solution.

**Example 2:** In this section, we consider physical model of nonlinear equation [1] as following showed in Fig. 4:

$$\frac{d^2y}{dt^2} + \frac{4r}{3m} \sin(y) - \frac{3F_0}{ml} \sin(w_0 t) = 0 \quad (25)$$

$$y(0) = A, y'(0) = 0$$

where  $A$  represents the amplitude of the oscillation. Motion is assumed to start from the position of maximum displacement with zero initial velocity. We will apply classic DTM and the multistage DTM to nonlinear ordinary differential Eq. (25). Applying classic DTM for Eq. (25) and from the trigonometric nonlinear in the example 1 given above,

$$(k+1)(k+2)Y(k+2) = -\frac{4r}{3m}F(k) + \frac{3F_0w_0^k}{mlk!} \sin\left(\frac{k\pi}{2}\right) \quad (26)$$

$$Y(0) = A, Y(1) = 0$$

where  $Y_i(n)$ , for  $n = 1 \dots M$ , satisfy the following recurrence relations,

$$(k+1)(k+2)Y_i(k+2) = -\frac{4r}{3m}F_i(k) + \frac{3F_0w_0^k}{mlk!} \sin\left(\frac{k\pi}{2}\right)$$

$$Y_0(0) = A, Y_0(1) = 0, Y_{i+1}(0) = Y_i(t^*), Y_{i+1}(1) = Y_i(t^*) \quad (27)$$

$$t \in [t_i, t_{i+1}], t^* = t_i, i = 0, \dots, M-1$$

is obtained and presented in the following in Fig. 5. RK4 method is also compared with trigonometric curve fitting solution.

In (13) expression,  $n = 1, t \in [0, 5], r = 1000, m = 10, F_0 = w_0 = 1 = 1$  and  $A = \pi/8$  solution of

$$y_{cf}(t) = 0.3926990818 \cos(3.64014563935407\pi t - 0.612387145462451292e-4)$$

is obtained and presented in the following in Fig. 6. RK4 method is also compared with trigonometric curve fitting solution.




Fig. 5: Plots of (a) displacement  $y$  versus time  $t$  and (b) phase portrait for  $A = \pi/12$




Fig. 4: The physical model of nonlinear equation [51]

By applying the multistage DTM to Eq. (25) is obtained Eq. (28) as following:

$$y(t) = \begin{cases} \sum_{k=0}^N Y_0(k) t^k, t \in [t_0, t_1] \\ \sum_{k=0}^N Y_1(k) t^k, t \in [t_1, t_2] \\ \vdots \\ \sum_{k=0}^N Y_M(k) t^k, t \in [t_{M-1}, t_M] \end{cases} \quad (28)$$

On the function (13) taken  $n = 1, t \in [0, 4], r = 1000, m = 10, F_0 = w_0 = 1 = 1$  and  $A = \pi/12$  solution of

$$y_{cf}(t) = 0.2617993878 \cos(3.65979516413592\pi t + 0.442004866524034105e-4)$$




Fig. 6: Plots of (a) displacement  $y$  versus time  $t$  and (b) phase portrait for  $A = \pi/8$




Fig. 7: Plots of (a) displacement  $y$  versus time  $t$  and (b) phase portrait for  $A = \pi/5$

In (13) equation,  $n = 1$ ,  $t \in [0, 6]$ ,  $r = 1000$ ,  $m = 10$ ,  $F_0 = w_0 = 1 = 1$  and  $A = \pi/5$  solution of

$$y_{cr}(t) = 0.6283185308 \cos(3.58502086997380\pi t - 0.625140793577042644e-4)$$

is obtained and presented in the following in Fig. 7. RK4 method is also compared with trigonometric curve fitting solution.

**Example 3:** In this section, we have Duffing equation with constant coefficient [1] as following showed in Fig. 8:

$$\frac{d^2x}{dt^2} + \frac{K_1}{m}x + \frac{K_2}{2mh^2}x^3 = \frac{F_0}{ml} \sin(w_0 t), x(0) = A, x'(0) = 0 \quad (29)$$

where  $A$  represents the amplitude of the oscillation. Motion is assumed to start from the position of maximum displacement with zero initial velocity. We will apply classic DTM and the multistage DTM to nonlinear ordinary differential Eq. (29). Applying classic DTM for Eq. (29),

$$(k+1)(k+2)X(k+2) = -\frac{K_1}{m}X(k) - \frac{K_2}{2mh^2} \sum_{k_2=0}^k \sum_{k_1=0}^{k_2} X(k_1)X(k_2-k_1)X(k-k_2) + \frac{F_0 w_0^k}{mk!} \sin\left(\frac{k\pi}{2}\right) \quad (30)$$

$$X(0) = A, X(1) = 0$$

where  $Y_i(n)$ , for  $n = 1 \dots M$ , satisfy the following recurrence relations,

$$(k+1)(k+2)X_i(k+2) = -\frac{K_1}{m}X_i(k) - \frac{K_2}{2mh^2} \sum_{k_2=0}^k \sum_{k_1=0}^{k_2} X_i(k_1)X_i(k_2-k_1)X_i(k-k_2) + \frac{F_0 w_0^k}{mk!} \sin\left(\frac{k\pi}{2}\right)$$

$$X_0(0) = A, X_0(1) = 0, X_{i+1}(0) = X_i(t^*), X_{i+1}(1) = X_i(t^*)$$

$$t \in [t_i, t_{i+1}], t^* = t_i, i = 0, \dots, M-1$$
(31)

By applying the multistage DTM to Eq.(29) is obtained Eq.(32) as following:

$$x(t) = \begin{cases} \sum_{k=0}^N X_0(k) t^k, t \in [t_0, t_1] \\ \sum_{k=0}^N X_1(k) t^k, t \in [t_1, t_2] \\ \vdots \\ \sum_{k=0}^N X_M(k) t^k, t \in [t_{M-1}, t_M] \end{cases}$$
(32)

On the function (13) taken  $n = 1, t \in [0, 3]$ ,

$$L = 1 \text{ m}, h = 0.9 \text{ m}, m = 10 \text{ kg}, K_1 = 1000 \text{ N/m}, K_2 = 1100 \text{ N/m}, F_0 = 1 \text{ N}, w_0 = 1 \text{ rad/s}$$




Fig. 8: The physical model of Duffing equation with constant coefficient [51]




Fig. 9: Plots of (a) displacement  $x$  versus time  $t$  and (b) phase portrait for  $A = 0.9 \tan(\pi/36)$






Fig. 10: Plots of (a) displacement  $x$  versus time  $t$  and (b) phase portrait for  $A = 0.9 \tan(\pi/18)$




Fig. 11: Plots of (a) displacement  $x$  versus time  $t$  and (b) phase portrait for  $A = 0.9 \tan(\pi/12)$

and  $A = 0.9 \tan(\pi/36)$  solution of

$$y_{cf}(t) = 0.7873979720e-1 \cos(10.01653269t + 0.1005230648e-3)$$

is obtained and presented in the following in Fig. 9. RK4 method is also compared with trigonometric curve fitting solution.

In (13) expression,  $n = 1, t \in [0, 4]$ ,

$$L = 1 \text{ m}, h = 0.9 \text{ m}, m = 10 \text{ kg}, K_1 = 1000 \text{ N/m}, K_2 = 1100 \text{ N/m}, F_0 = 1 \text{ N}, w_0 = 1 \text{ rad/s}$$

and  $A = 0.9 \tan(\pi/18)$  solution of

$$y_{cf}(t) = 0.1586942826 \cos(10.06478006t + 0.3279565577e-3)$$

is obtained and presented in the following in Fig. 10. RK4 method is also compared with trigonometric curve fitting solution.

In (13) equation,  $n = 1, t \in [0, 5]$ ,

$$L = 1 \text{ m}, h = 0.9 \text{ m}, m = 10 \text{ kg}, K_1 = 1000 \text{ N/m}, K_2 = 1100 \text{ N/m}, F_0 = 1 \text{ N}, w_0 = 1 \text{ rad/s}$$

and  $A = 0.9 \tan(\pi/12)$  solution of

$$y_{cr}(t) = 0.2411542732 \cos(10.14819210t + 0.7173231348e-3)$$

is obtained and presented in the following in Fig. 11. RK4 method is also compared with trigonometric curve fitting solution.

**Example 4:** In this section, we consider the motion equation of the pendulum with harmonic stringer point [1] as following showed in Fig. 12:

$$\begin{aligned} \frac{d^2\theta}{dt^2} + \left( \frac{g}{l} - \frac{w_0^2 Y}{1} \cos(w_0 t) \right) \sin \theta &= 0 \\ \theta(0) &= A, \quad \dot{\theta}(0) = 0 \end{aligned} \quad (33)$$

where  $A$  represents the amplitude of the oscillation. Motion is assumed to start from the position of maximum displacement with zero initial velocity. We will apply classic DTM and the multistage DTM to nonlinear ordinary differential Eq. (33). Applying classic DTM for Eq. (33) and from the trigonometric nonlinear in the example 1 given above,

$$\begin{aligned} (k+1)(k+2)\Theta(k+2) &= F(k) \left( \frac{g}{l} - \frac{Y \cos(0.5\pi k) w_0^{k+2}}{lk!} \right) \\ \Theta(0) &= A, \quad \Theta(1) = 0 \end{aligned} \quad (34)$$

where  $\Theta_i(n)$ , for  $n = 1 \dots M$ , satisfy the following recurrence relations,

On the function (13) taken  $n=1, t \in [0, 5], l=1 \text{ m}, w_0=1 \text{ rad/s}, Y=0.25 \text{ m}, g=9.81 \text{ m/s}^2$  and  $A = \pi/12$  solution of

$$y_{cr}(t) = 0.261895058654144197 \cos(0.979981126902420784\pi t - 0.431628482521539199e-4)$$

is obtained and presented in the following in Fig. 13. RK4 method is also compared with trigonometric curve fitting solution.




Fig. 13: Plots of (a) displacement  $\theta$  versus time  $t$  and (b) phase portrait for  $A = \pi/12$




Fig.12: Pendulum with harmonic stringer point:  $y(t) = Y \cos w_0 t$  [51]

$$\begin{aligned} (k+1)(k+2)\Theta(k+2) &= F(k) \left( \frac{g}{l} - \frac{Y \cos(0.5\pi k) w_0^{k+2}}{lk!} \right) \\ \Theta_0(0) &= A, \quad \Theta_0(1) = 0, \quad \Theta_{i+1}(0) = \Theta_i(t^*), \quad \Theta_{i+1}(1) = \Theta_i(t^*) \\ t &\in [t_i, t_{i+1}], \quad t^* = t_i, \quad i = 0, \dots, M-1 \end{aligned} \quad (35)$$

By applying the multistage DTM to Eq. (33) is obtained Eq. (36) as following:

$$\theta(t) = \begin{cases} \sum_{k=0}^N \Theta_0(k) t^k, & t \in [t_0, t_1] \\ \sum_{k=0}^N \Theta_1(k) t^k, & t \in [t_1, t_2] \\ \vdots \\ \sum_{k=0}^N \Theta_M(k) t^k, & t \in [t_{M-1}, t_M] \end{cases} \quad (36)$$




Fig. 14: Plots of (a) displacement  $\theta$  versus time  $t$  and (b) phase portrait for  $A = \pi/4$




Fig. 15: Plots of (a) displacement  $\theta$  versus time  $t$  and (b) phase portrait for  $A = 5\pi/12$

In (13) expression,  $n = 1, t \in [0, 6], l = 1 \text{ m}, w_0 = 1 \text{ rad/s}, Y = 0.25 \text{ m}, g = 9.81 \text{ m/s}^2$  and  $A = \pi/4$  solution of

$$y_{cr}(t) = 0.788238375005162582 \cos(0.946384386107354980\pi + 6.28291712268189606)$$

is obtained and presented in the following in Fig. 14. RK4 method is also compared with trigonometric curve fitting solution.

In (13) equation,  $n = 1, t \in [0, 7], l = 1 \text{ m}, w_0 = 1 \text{ rad/s}, Y = 0.25 \text{ m}, g = 9.81 \text{ m/s}^2$  and  $A = 5\pi/12$  solution of

$$y_{cr}(t) = -1.32267015669864163 \cos(0.879650367165896506\pi - 28.2751089644735885)$$

is obtained and presented in the following in Fig. 15. RK4 method is also compared with trigonometric curve fitting solution.

We can observe that local changes and the phase plane trajectories trigonometric curve fitting obtained using the multi-stage DTM are in high agreement with local changes and the phase plane trajectories obtained using RK4 method. In calculations of trigonometric curve fitting calculations were used command maple fit.

## CONCLUSIONS

In this paper, we carefully applied the multistage DTM, a reliable modification of the DTM that

improves the convergence of the series solution to the nonlinear vibrating equations. For data obtained from the multi-step DTM, generated the most appropriate trigonometric curve fitting. The method provides immediate and visible symbolic terms of analytic solutions, as well as numerical approximate solutions to both linear and nonlinear differential equations. The validity of the proposed method has been successful by applying it for the nonlinear vibrating equations. As understood from the above examples, the comparison between multistage DTM and RK4, both results obtained by both methods are accurate and close to each other.

## REFERENCES

1. Mehdipour, I., D.D. Ganji and M. Mozaffari, Application of the energy balance method to nonlinear vibrating equations. *Current Applied Physics*, 10: 104-112.
2. Akbarzade, M., D.D. Ganji and H. Pashaei, 2008. *Progress in Electromagnetic Research*, C 3: 57-66.
3. Rafei, M., D.D. Ganji, H. Daniali and H. Pashaei, 2007. The variational iteration method for nonlinear oscillators with discontinuities. *Journal of Sound and Vibration*, 305: 416-620.
4. Hagedorn, P., 1981. *Nonlinear Oscillations translated by Wolfram Stadler*. Clarendon Press, Oxford.
5. He, J.H., 2006. Some asymptotic methods for strongly nonlinear equations. *Int. J. Mod. Phys. B* 10: 1141.
6. He, J.H., 2006. *Non-perturbative Methods for Strongly Nonlinear Problems*. Dissertation, de-Verlag im Internet GmbH.
7. He, J.H., 1999. Variational iteration method: A kind of nonlinear analytical technique: Some examples. *Int. J. Nonlinear Mech.* 34 (4): 699-708.
8. He, J.H., 1999. Homotopy perturbation technique. *Comput. Method Appl. Mech. Eng.*, 178 (3/4): 257-262.
9. Liao, S.J., 1992. *The homotopy analysis method and its applications in mechanics*. Ph.D. Dissertation, Shanghai Jiaotong University.
10. Zhang, S., 2008. Application of Exp-function method to high-dimensional nonlinear evolution equation. *Chaos Solitons Fractals*, 38: 270-276.
11. Bekir, A. and A. Boz, 2008. Exact solutions for nonlinear evolution equations using Exp-function method. *Phys. Lett. A*, 372: 1619-1625.
12. He, J.H., 1999. Modified straightforward expansion. *Meccanica*, 34 (4): 287-289.
13. He, J.H., 2002. Modified Lindstedt-Poincare methods for some strongly nonlinear oscillations. Part I: A new transformation. *Int. J. Nonlinear Mech.*, 37: 315.
14. He, J.H., 2001. Iteration perturbation method for strongly nonlinear oscillations. *J. Vib. Control*, 7 (5): 631-642.
15. He, J.H., 2001. Bookkeeping parameter in perturbation methods. *Int. J. Nonlinear Sci. Numer. Simul.*, 2 (3): 257-264.
16. He, J.H., 2002. Preliminary report on the energy balance for nonlinear oscillations. *Mech. Res. Commun.*, 29: 107-111.
17. He, J.H., 2006. Some asymptotic methods for strongly nonlinear equations. *Int. J. Mod. Phys. B* 20: 1141-1199.
18. He, J.H., 2008. Max-min approach to nonlinear oscillators. *Int. J. Nonlinear Sci. Numer. Simul.*, 9 (2): 207-210.
19. He, J.H., 2004. Chengtian's inequality and its applications. *Appl. Math. Comput.*, 151 (3): 887-891.
20. Mickens, R.E., 1996. *Oscillations in Planar Dynamics Systems*. World Scientific, Singapore.
21. Mickens, R.E., 1987. Iteration procedure for determining approximate solutions to non-linear oscillator equation. *J. Sound Vib.*, 116: 185-188.
22. Zhou, J.K., 1986. *Differential Transformation and Its Applications for Electrical Circuits*. Huazhong University Press, Wuhan, China, (In Chinese).
23. Fatma Ayaz, 2004. Solutions of the system of differential equations by differential transform method. *Appl. Math. Comput.*, 147: 547-567.
24. Fatma Ayaz, 2004. Application of differential transform method to differential-algebraic equations. *Appl. Math. Comput.*, 152: 649-657.
25. Arikoglu, A. and I. Ozkol, 2005. Solution of boundary value problems for integro-differential equations by using differential transform method. *Appl. Math. Comput.*, 168: 1145-1158.
26. Bildik, N., A. Konuralp, F. Bek and S. Kucukarslan, 2006. Solution of different type of the partial differential equation by differential transform method and Adomian's decomposition method. *Appl. Math. Comput.*, 127: 551-567.
27. Arikoglu, A. and I. Ozkol, 2006. Solution of difference equations by using differential transform method. *Appl. Math. Comput.*, 173 (1): 126-136.
28. Arikoglu, A. and I. Ozkol, 2006. Solution of differential difference equations by using differential transform method. *Appl. Math. Comput.*, 181 (1): 153-162.
29. Liu, H. and Y. Song, 2007. Differential transform method applied to high index differential-algebraic equations. *Appl. Math. Comput.*, 184 (2): 748-753.
30. Momani, S. and M. Noor, 2007. Numerical comparison of methods for solving a special fourth-order boundary value problem. *Appl. Math. Lett.*, 191 (1): 218-224.
31. Hassan, I.H., 2008. Comparison differential transformation technique with Adomian decomposition method for linear and nonlinear initial value problems. *Chaos Solitons Fractals*, 36 (1): 53-65.
32. Hassan, I., 2008. Application to differential transformation method for solving systems of differential equations. *Appl. Math. Model.*, 32 (12): 2552-2559.
33. El-Shahed, M., 2008. Application of differential transform method to non-linear oscillatory systems. *Commun. Nonlinear Sci. Numer. Simul.*, 13 (8): 1714-1720.

34. Odibat, Z., 2008. Differential transform method for solving Volterra integral equation with separable kernels. *Math. Comput. Modelling*, 48 (7-8): 144-149.
35. Momani, S., Z. Odibat and V. Erturk, 2007. Generalized differential transform method for solving a space-and time-fractional diffusion-wave equation. *Phys. Lett. A* 370 (5-6): 379-387.
36. Erturk, V., S. Momani and Z. Odibat, 2008. Application of generalized differential transform method to multi-order fractional differential equations. *Commun. Nonlinear Sci. Numer. Simul.*, 13 (8): 1642-1654.
37. Odibat, Z. and S. Momani, 2008. Generalized differential transform method for linear partial differential equations of fractional order. *Appl. Math. Lett.*, 21 (2): 194-199.
38. Momani, S. and Z. Odibat, 2008. A novel method for nonlinear fractional partial differential equations: Combination of DTM and generalized Taylor's formula. *J. Comput. Appl. Math.*, 220 (1-2): 85-95.
39. Odibat, Z., S. Momani and V. Erturk, 2008. Generalized differential transform method: Application to differential equations of fractional order. *Appl. Math. Comput.*, 197 (2): 467-477.
40. Kuo, B. and C. Lo, 2009. Application of the differential transformation method to the solution of a damped system with high nonlinearity. *Nonlinear Anal. TMA* 70 (4): 1732-1737.
41. Al-Sawalha, M. and M. Noorani, 2009. Application of the differential transformation method for the solution of the hyperchaotic Rössler system. *Commun. Nonlinear Sci. Numer. Simul.*, 14 (4): 1509-1514.
42. Chen, S. and C. Chen, 2009. Application of the differential transformation method to the free vibrations of strongly non-linear oscillators. *Nonlinear Anal. RWA*, 10 (2): 881-888.
43. Zaid M. Odibat, Cyrille Bertelle, M.A. Aziz-Alaoui and Gérard H.E. Duchamp, 2010. A multi-step differential transform method and application to non-chaotic or chaotic systems. *Computers and Mathematics with Applications*, 59: 1462-1472.
44. Gökdoğan, A. and M. Merdan, 2010. A numeric-analytic method for approximating the Holling Tanner model. *Studies in Nonlinear Science*, 1 (3): 77-81.
45. Merdan, M. and A. Gökdoğan, 2011. Solution of Nonlinear Oscillators with Fractional Nonlinearities by using The Modified Differential Transformation Method. *Mathematical and Computational Applications*, 16 (3): 761-772.
46. Merdan, M., A. Gökdoğan and A. Yildirim, 2011. On the numerical solution of the model for HIV infection of CD4+T Cells. *Computers and Mathematics with Applications*, 62 (1): 118-123.
47. Merdan, M., A. Gökdoğan and V.S. Ertürk, 2011. A numeric-analytic method for approximating the chaotic three-species food chain models. *International Journal of the Physical Sciences*, 6 (7): 1822-1833.
48. Chang, S.H. and I.L. Chang, 2008. A New Algorithm for Calculating One-Dimensional Differential Transform of Nonlinear Functions. *Appl. Math. Comput.*, 195 (2): 799-808.
49. Baolei Gu, 2009. Trigonometric Curve Fitting Based on Genetic Algorithm and the Application of Data Processing in Geography. *Communications in Computer and Information Science*, 51 (2): 104-109.
50. Ganji, S.S., D.D. Ganji, A.G. Davodi and S. Karimpour, 2010. Analytical solution to nonlinear oscillation system of the motion of a rigid rod rocking back using max-min approach. *Applied Mathematical Modelling*, 34: 2676-2684.
51. Rao, S.S., 2006. *Mechanical Vibrations Book*. Fourth Ed., ISBN: 978-964-9585-5-0.
52. Mohyud-Din, S.T., M.A. Noor and K.I. Noor, 2009. Some relatively new techniques for nonlinear problems. *Mathematical Problems in Engineering*, Hindawi, Article ID 234849, 25 pages, doi:10.1155/2009/234849.
53. Mohyud-Din, S.T., M.A. Noor, K.I. Noor and M.M. Hosseini, 2010. Solution of singular equations by He's variational iteration method. *International Journal of Nonlinear Sciences and Numerical Simulation*, 11 (2): 81-86.
54. Mohyud-Din, S.T. and M.A. Noor, 2009. Homotopy perturbation method for solving partial differential equations. *Zeitschrift für Naturforschung A-A Journal of Physical Sciences*, 64a: 157-170.
55. Mohyud-Din, S.T., A. Yildirim and G. Demiri, 2011. Analytical solution of wave system in  $\mathbb{R}^1$  with coupling controllers. *International Journal of Numerical Methods for Heat and Fluid Flow*, Emerald, IF= 0.674, 21 (2): 198-205.
56. Mohyud-Din, S.T., 2010. Variational iteration techniques for boundary value problems. *VDM Verlag*, ISBN 978-3-639-27664-0.