

**An Assessment of the Accumulation Potential of Lead(pb), Zinc(zn) and Cadmium(cd) by *Avicennia Marina* (Forssk.) Vierh. In Vamleshwar Mangroves near Narmada Estuary, West Coast of Gujarat, India**

<sup>1</sup>J.I. Nirmal Kumar, <sup>1</sup>P.R. Sajish, <sup>2</sup>Rita N. Kumar, <sup>1</sup>Basil George and <sup>1</sup>Shailendra Viyol

<sup>1</sup>P.G. Department of Environmental Science and Technology, Institute of Science and Technology for Advanced Studies and Research (ISTAR), Vallabh Vidya Nagar, Gujarat- 388120, India

<sup>2</sup>Department of Biological Science and Environmental Science, N.V. Patel College of Pure and Applied Science, Vallabh Vidya Nagar, Gujarat- 388 120, India

**Abstract:** The study deals with the accumulation of Pb, Zn and Cd in an important mangrove species, *Avicennia marina* (Forssk.) Vierh. in Vamleshwar mangrove ecosystem, near Narmada estuary, West coast of Gujarat, India at a height differences of 0.5, 1.5, 2.5 mtrs and carried out under field conditions during October 2009. The site was located on 21°30'11.55" N latitude and 72°43'53.68" E longitude. Mangrove receives heavy metal pollution from upstream areas of Narmada estuary and highly populated settlements. However, little is known about the capacity of mangrove plants to take up and store heavy metals. Water, sediment and plant parts such as roots, stems and leaves were analyzed for finding the trace metal accumulation of different height groups by Inductive Coupled Plasma Analyser (ICPA). The amounts of metals found in the water, sediment and plant parts were in the order of Pb>Zn>Cd. The average concentrations of heavy metals in the water were 57.83 mg LG<sup>-1</sup> for Pb, 3.89 mg L<sup>-1</sup> for Zn and 0.42 mg LG<sup>-1</sup> for Cd. It was observed that the average concentrations of Pb (73.6 mg LG<sup>-1</sup>), Zn (8.1 mg LG<sup>-1</sup>) and Cd (0.73 mg LG<sup>-1</sup>) in the sediments were below the critical soil concentrations. The concentrations of heavy metals in different parts of *Avicennia marina* were in the order Roots>stem>leaf except for Cd, but Cd found higher in leaf. The ranges of heavy metals in plants were 18.5-102.2 mg LG<sup>-1</sup> for Pb, 3.5-19.5 mg LG<sup>-1</sup> for Zn and 0.2-4.1 mg LG<sup>-1</sup> for Cd. The concentrations of all heavy metals in *Avicennia marina* except Pb were falling within normal range and were much more in the plants having highest height. This study has therefore shown the potential of *Avicennia marina* as a phytoremediation species for selected heavy metals in many mangrove ecosystems.

**Key words:** Heavy metal accumulation % *Avicennia marina* (Forssk.) Vierh % Mangrove plant parts % Sediments % Vamleshwar mangrove ecosystem

## INTRODUCTION

Mangrove ecosystems are highly productive and play a vital role as a major primary producer within estuarine systems. The uniqueness of *A. marina* root systems serve as habitat and nursery area for many juvenile fish and crustaceans, which have both direct and indirect socio-economic significance and are of great importance to many scientific studies. They also provide erosion mitigation and stabilization for adjacent coastal landforms [1]. In a plant-soil system, strong absorption and fixation of heavy metals by soil can easily cause

residual accumulation in the soil, resulting in over-absorption of heavy metals by growing plants [2, 3]. These plant products are harmful to the health of humans [4].

The uptake Pb, Zn and Cd is passive and its translocation from roots to other plant organs is generally low [5-7]. Many studies have been carried out on various plants to determine its heavy metal accumulation capability in different aquatic and forest environments and vegetable crops [8-12]. There are several studies on heavy metal contaminations in mangrove sediments and their effects on organisms but little is known about heavy

metals uptake by mangrove plants [13, 14]. Therefore, an attempt was made to understand the metal phytoremediation potential of *A. marina* in a mangrove ecosystem at Vamleshwar, Gujarat, India, that would benefit in the wise-management and utilization of this ecosystem. The present study presents data on the concentrations of three metals (Pb, Zn and Cd) in water, sediment and in leaves, stems and roots of *A. marina*.

## MATERIALS AND METHODS

**Sampling Site:** Sampling was conducted at Vamleshwar *A. marina* plantation, which is located 45 km to west of Ankleshwar, Gujarat, India and facing the Arabian Sea at a latitude of 21°30'11.55" N and longitude of 72°46'53.68" E. The study area has various natural resources, vast array of biological diversity and coastal and riverine fishing activities (Figure 1). The plant parts of different height groups viz., 0.5M, 1.5M and 2.5M were collected in three replicates from the study site (n=10) during October 2009. The roots were carefully removed from the sediments while the leaves and stems were collected using a pair of scissors. The samples were immediately wrapped with an aluminium foil and labeled. All the samples were kept in a cooler box with ice at 4°C for storage during transportation and brought to the

laboratory [15]. After reaching the laboratory, root, stem and leaves samples were washed with double distilled water before storage. Water and sediment samples were collected from same sites where plant samples were collected. Soon after collection, the water samples were filtered through 0.45 Fm millimeter filter and preserved in plastic bottles by the addition of a few drops of nitric acid. Sediment samples were preserved in air-dry plastic bags. The samples were labeled carefully and brought to the laboratory for further analysis.

**Analysis of Plant Parts, Water and Sediment for Heavy Metals:** The plant parts of roots, stems and leaves samples were dried at 60°C for two days to achieve constant weight, homogenised and grounded to a fine powder. Samples then stored in plastic vials with labels and kept in desiccators. The samples of water, sediment and plant-parts were chemically analysed for detection for Pb, Zn and Cd. Accurately 1g of dry powder of each sample was weighed and digested with con. HNO<sub>3</sub>, H<sub>2</sub>SO<sub>4</sub> and H<sub>2</sub>O<sub>2</sub> (2:6:6) as prescribed by [16]. Towards the end of the digestion, the flasks were brought to near dryness. The solutions were made to 20ml each in measuring cylinder with double distilled water and analysed for Pb, Zn and Cd by Inductive Coupled Plasma Analyser (ICPA) (Perkin-Elmer ICP Optima 3300 RL, U.S.A) at Sophisticated




Fig. 1: Location of the Study Area.

Instrumentation Centre for Applied Research and Testing (SICART), Vallabh Vidyanagar, Gujarat, India. Mean values of triplicate of each sample of the water, sediment and plant samples were calculated and considered.

**Statistical Analysis:** Pearson correlation coefficient analysis was done between metal-pairs in plants to check if differences exist between different metal combinations in either root, stem or leaf system. The products of the correlation coefficient (r) were evaluated as follows:

0-0.3: No correlation; 0.3-0.5: Low correlation; 0.5-0.7 Medium correlation; 0.7-0.9 High correlation; 0.9-1.0 Very high correlation.

Further the comparison of the concentration of an element in an aquatic organism with that of the same element in the water in which the organism lives was made. This is the ratio between the concentration of the element in the organism and that of in the water, which is known as Concentration Factor [17].

Further the data was analyzed using ANOVA to detect if any significant differences in means of each heavy metal exist between mangrove sediments, roots, stems and leaves and of different heights. Significance value was set at 5%.

## RESULT AND DISCUSSION

### Heavy Metal Accumulation in the Mangrove Sediments, Water and Mangrove Plant Parts:

The average concentration of Pb (73.6 mg LG<sup>-1</sup>) was higher in the sediments followed by Zn (8.1 mg LG<sup>-1</sup>) and Cd (0.73 mg LG<sup>-1</sup>) (Figure 2). Like sediment the concentration of Pb (57.83 mg LG<sup>-1</sup>) in water was greater than the other two elements (Zn-3.89 mg LG<sup>-1</sup> and Cd-0.42 mg LG<sup>-1</sup>) (Figure 2). However, when compared to the critical soil concentrations, the total concentrations for all selected heavy metals present in the sediments were below the critical soil concentration values (18 and Table 1). Meanwhile, the comparison of the total concentrations of the selected heavy metals in the sediments with those from other countries showed that the concentration of Cd in this study was greater than those measured in Buloh River and Khatib Bongsu River at Singapore by [19], Brisbane River, Australia by [20] and Mai Po River, Hong Kong by [21]. The values of the ratio between element concentrations in the sediments and those in the water were low for Pb (1.27) where as that of Zn was observed high (2.08) (Table 2).




Fig. (2A-2C): The mean concentrations of Pb, Zn and Cd in the roots, stems, leaves, water and sediments of *A. marina* (n=3) plantation of different height groups viz., 0.5M, 1.5M and 2.5M.

Table 1: The total concentrations of heavy metals (mg LG<sup>1</sup>) in mangrove sediments of Vamleshwar mangrove ecosystem (present study) relative to other countries. Critical soil concentrations follow [18]

	Pb	Zn	Cd	References
Vamleshwar Mangrove Ecosystem, Gujarat, India	73.6±4.72	8.1±2.94	0.73±0.153	Present Study
Singapore (Buloh River)	12.28±5.18	51.24±39.97	0.181±0.189	[19]
Singapore (Khatib Bongsu River)	30.98±6.16	120.23±13.90	0.266±0.171	[19]
Australia (Brisbane River)	20.1-81.9	40.8-144.0	<0.1-1.9	[20]
Hong Kong (Mai Po)	161.6-219.8	277.2-321.2	0.5-0.6	[21]
Critical Soil Concentrations	100-400	70-400	3-8	

Table 2: Elemental concentration ratios between sediments and water

	Sediment (mg LG <sup>1</sup> )	Water (mg LG <sup>1</sup> )	Sediment/Water
Pb	73.6	57.83	1.27
Zn	8.1	3.89	2.08
Cd	0.73	0.42	1.74

Table 3: Concentration factors calculated for different parts of *A. marina* and elements

	Pb	Zn	Cd	Mean
Leaf	0.84	2.89	5.71	3.15
Stem	0.38	1.49	1.83	1.23
Root	1.57	4.0	4.05	3.21
Mean	0.93	2.79	3.86	
SD	0.60	1.26	1.95	

Table 4: Pearson correlation coefficient between concentrations of heavy metal pairs in root, stem and leaf systems of *A. marina*

Analysis metal-pair	Root system	Stem system	Leaf system
Zn × Pb**	0.655	0.844	0.635
Cd × Pb**	-0.982	0.990	0.980
Zn × Cd**	-0.500	0.911	0.775

\*\* High Correlation (r=0.6-0.9)

Heavy metals concentrations were detected in the mangrove parts prove that Pb, Zn and Cd were distributed in all three parts of *A. marina*. From the observations, *A. marina* accumulated higher concentration of heavy metals in the root system compared to the other parts of plant except Cd which found higher in leaves followed by root and stem. More accumulation was observed in the plant parts of tallest plant group i.e. 2.5M. The average concentration of Pb in leaf, stem and root were 48.64, 21.94 and 91.07 mg LG<sup>1</sup> while for Zn 11.24, 5.78 and 15.53 mg LG<sup>1</sup> whereas for Cd 2.40, 0.77 and 1.70 mg LG<sup>1</sup> respectively (Figure 2). The average concentration values of the elements in the plants declined according to this sequence: Pb>Zn>Cd. The critical ranges of heavy metals in plants were 30-300 mg LG<sup>1</sup> for Pb, 100-400 mg LG<sup>1</sup> for Zn and 10-30 mg LG<sup>1</sup> for Cd. The concentrations of all heavy metals in *A. marina* except Pb were within the normal range in the present study.

**Statistical Analysis:** The concentration factor (C.F) for all three heavy metals and that of different components of *A. marina* also were evaluated (Table 3). The mean C.F. value of elements in the different parts declined according to this sequence: Cd> Zn > Pb. The mean C.F. for the various elements calculated for root is rather high followed by leaf and stem (Table 3). The concentration distribution in plant parts like leaves, stems and roots may vary depending on the concentration of heavy metals in the sediment, height of the plants, the types of heavy metals and also the tolerance of the species and its parts towards the heavy metals. The present study is well corroborated with other works [6, 7].

The output of Pearson correlation coefficient (r) analysis on combinations of different metal-pairs which are present together in either roots, stems or leaves of *Avicennia marina* showed that a high + correlation (r=0.6-0.9) between all the metal-pairs analyzed (Table. 4) except in root system.

The ANOVA also showed significant difference of the heavy metals (Pb, Zn and Cd) concentrations between mangrove sediments and the roots of *A. marina* (Pb,  $p=0.01$ ; Zn,  $p=0.005$ ; Cd,  $p=0.01$ ).

The study showed the transport mechanism of metals from sediment to plant and their accumulation in various parts of the plant. The transport mechanism and accumulation pattern of heavy metals can be elaborated as follows: sediment>root system>stem system>leaf system.

## CONCLUSIONS

This study has showed that *A. marina* possess the capacity to take up selected heavy metals via its roots and storing in its leaves without any sign of injury. As it grows more aged, its capability of accumulating heavy metals is also increasing much fold. This suggests the potential of *A. marina* as a phytoremediation species for many mangrove ecosystems. The luxuriant growth of *A. marina* in comparison to other mangrove species is evident of its adaptability even under polluted conditions.

## ACKNOWLEDGEMENTS

The authors are highly thankful to Ministry of Environment and Forest (MoEF), New Delhi for financial support for the project.

## REFERENCES

1. Harty, C., 1997. Mangrove in New South Wales and Victoria. Vista Publication, Melbourne. pp: 47.
2. Ravikumar, S., G. Prakash Williams, S. Shanthi, N. Anitha Anantha Gracelin, S. Babu and P.S. Parimala, 2007. Effect of heavy metals (Hg and Zn) on the growth and phosphate solubilising activity in halophilic phosphobacteria isolated from Manakudi mangrove. *J. Environ. Biol.*, 28: 109-114.
3. Lian, Y., J. Xu, P. Lin, S. Meguro and S. Kawachi, 1999. Five heavy metals in propagules of ten mangrove species of China. *J. Wood Sci.*, 45: 18-24.
4. Defew, L.H., J.M. Mair and H.M. Guzman, 2005. An assessment of metal contamination in mangrove sediments and leaves from Punta Mala Bay, Pacific Panama. *Marine Pollution Bulletin*, 50: 547-552.
5. Nirmal Kumar, J.I., H. Soni, R.N. Kumar and I. Bhatt, 2008. Biomonitoring of trace elements in some selected aquatic macrophytes with reference to lake contamination: A case study of Pariyej community reserve, Gujarat, India. *Asian J. Microbiol. Biotechnology and Environ. Sci.*, 10(4): 803-810.
6. Baker, A.J. and P.I. Walker, 1990. Ecophysiology of metal uptake by tolerant plants, In: *Heavy metal tolerance in plants evolutionary aspects* (Ed.: A.J. Shaw). CRC, Press, Boca Raton, FL. pp: 155-178.
7. De Lacerda, L.D. and J.J. Abrao, 1986. Heavy metal accumulation by mangrove and saltmarsh intertidal sediments. *Marine Pollution Bulletin*, 17: 246-250.
8. Nirmal Kumar, J.I., H. Soni, R.N. Kumar and I. Bhatt, 2009. Hyper accumulation and mobility of heavy metals in vegetable crops in India. *The J. Agriculture and Environ.*, 10: 29-38.
9. Nirmal Kumar, J.I., H. Soni and R.N. Kumar, 2007. Characterization of Heavy Metals in Vegetables Using Inductive Coupled Plasma Analyzer (ICPA). *J. Appl. Sci. and Environ. Manag.*, 11(3): 75-79.
10. Nirmal Kumar, J.I., H. Soni and R.N. Kumar, 2006. Biomonitoring of selected freshwater macrophytes to assess the lake trace element contamination: A case study of Nal Sarovar Bird Sanctuary, Gujarat, India. *J. Limnol.*, 65(1): 9-16.
11. Silva, C.A.R., A.P. de Silva and S.R. Oliveira, 2006. Concentration, stock and transport rate of heavy metals in a tropical red mangrove, Natal, Brazil. *Marine Chemistry*, 99: 2-11.
12. Yu, K.F., B.S. Kamber, M.G. Lawrence, A. Greig and J.X. Zhao, 2007. High precision analysis on annual variation of heavy metals, lead isotopes and rare earth elements in mangrove tree rings by inductively coupled plasma mass spectrometry. *Nuclear Instruments and Methods in Physics Research, Section B-Beam interactions with materials and atoms*, 255: 399-408.
13. Seng, C.E., P.P.E. Lim and T.T. Ang, 1987. Heavy metal concentrations in coastal sea water and sediment of Prai Industrial Estate, Penang, Malaysia. *Marine Pollution Bulletin*, 18: 611-12.
14. Ismail, A. and M.I.N. Asmah, 1992. Copper, zinc, lead and cadmium in intertidal molluscs and sediment off Seberang Perai coastline, Malaysia. 4th Princess Chulabhorn International Science Congress, Bangkok, Thailand.
15. Prica, M., B. Dalmacija, S. Roncevic, D. Krcmar and M. Becelic, 2007. A comparison of sediment quality results with acid volatile sulfide (AVS) and simultaneously extracted metals (SEM) ratio in Vojvodina (Serbia) sediments. *Science of the Total Environ.*, 389: 235-44.
16. Saison, C., C. Schwartz and J.L. Morel, 2004. Hyperaccumulation of metals by *Thlaspi caerulescens* as affected by root development and Cd-Zn/Ca-Mg interactions. *International J. Phytoremediation*, 6(1): 49-61.
17. De Bortoli, M., P. Gaglione, A. Malvicini and A. Polvani, 1968. Concentration factors for strontium and caesium in fish of the lakes in the region of Varese. *Giorn. Fisica Sanit. Radioprot.*, 12: 324-331.
18. Alloway, B.J., 1990. *Heavy metals in soils*. John Wiley and Sons, Inc, New York.
19. Cuong, D.T., S. Bayen, O. Wurl, K. Subramanian, K.K.S. Wong, N. Sivasothi and J.P. Obbard, 2005. Heavy metal contamination in mangrove habitats of Singapore. *Marine Pollution Bulletin*, 50: 1713-44.
20. Mackey, A.P., M. Hodgkinson and R. Nardella, 1992. Nutrient levels and heavy metals in mangrove sediments from the Brisbane River, Australia. *Marine Pollution Bulletin*, 24: 418-20.
21. Ong Che, R.G., 1999. Concentration of seven heavy metals in sediment and mangrove root samples from Mai Po, Hong Kong. *Marine Pollution Bulletin*, 39: 269-79.