

Antifungal Activity of Essential Oils from Indian Medicinal Plants Against Human Pathogenic *Aspergillus fumigatus* and *A. niger*

Sunita Bansod and Mahendra Rai

Department of Biotechnology, Amravati University, Amravati-444107 India

Abstract: Oils extracted from fifteen medicinal plants were screened for their activity against *A. fumigatus* and *A. niger* by disc diffusion method. Minimum inhibitory concentrations (MICs) of oils (%v/v) against *Aspergillus fumigatus* and *Aspergillus niger* done by agar dilution method and minimum inhibitory concentration (MIC) and minimum cidal concentration (MCCs) data (%v/v) obtained by the broth micro dilution method. The results showed that the maximum antimycotic activity was demonstrated by oils of *Cymbopogon martini*, *Eucalyptus globulus* and *Cinnamomum zylanicum* as compared to control, followed by *Cymbopogon citratus* which showed activity similar to control (miconazole nitrate). The oils of *Mentha spicata*, *Azadirachta indica*, *Eugenia caryophyllata*, *Withania somnifera* and *Zingiber officinale* exhibited moderate activity. The oils of *Cuminum cyminum*, *Allium sativum*, *Ocimum sanctum*, *Trachyspermum copticum*, *Foeniculum vulgare* and *Elettaria cardamomum* demonstrated comparatively low activity against *A. niger* and *A. fumigatus* as compared to control. Mixed oils showed maximum activity as compared to standard. These results support the plant oils can be used to cure mycotic infections and plant oils may have role as pharmaceutical and preservatives.

Key words: Essential oil • Antifungal activity • Minimum inhibitory concentration • *Aspergillus*

INTRODUCTION

Infectious diseases accounts for high proportion of health problems in the developing countries including India. Microorganisms have developed resistance to many antibiotics and as a result, immense clinical problem in the treatment of infectious diseases has been created [1]. The resistance of the organisms increased due to indiscriminate use of commercial antimicrobial drugs commonly used for the treatment of infectious disease. This situation forced the researchers to search for new antimicrobial substance from various sources including medicinal plants [2]. There are alarming reports of opportunistic fungal infections [3]. The infections caused by opportunistic fungi are included under new spectrum of fungal pathogens. Such fungi were earlier reported from various plants as pathogens. But now they are known to cause disease in human beings. There is an increasing awareness amongst clinicians and microbiologists pertaining to importance of infection caused by opportunistic fungi [4]. Aspergillosis is caused due to inhalation of *Aspergillus fumigatus* spores. *Aspergillus fumigatus* is an opportunistic pathogen which

usually affects cavities that have formed in the lungs from preexisting lung diseases. In the lungs, *Aspergillus fumigatus* forms tangled mass of fungus fibers, blood clots. The fungus mass gradually enlarges, destroying lung tissue in the process, but usually does not spread to other areas.

There are 2600 plant species of which more than 700 are noted for their uses as medicinal herbs [5]. In folk medicine, medicinal herbs and plant products were used in treating a wide spectrum of infections and other diseases. Today, a great number of different medicinal tea and other plant products are available in market (including cosmetics and pharmaceuticals), which contains biologically active substances. In recent years, there has been a gradual revival of interest in the use of medicinal and aromatic plants in developed as well as in developing countries, because plant-derived drugs have been reported to be safe and without side-effects.

A survey of literature reveals that there are many essential oils which possesses antifungal activity [6-13]. In tuberculosis, treatment contains high doses of antibiotic due to resistance and side effects of this antibiotic, patients take more time for cure Therefore, we

Table 1: Oils of Indian medicinal plants including the botanical name, Local name, family and traditional use

Botanical name	Local name	Family	Plant part used	Medicinal use
<i>Allium sativum</i>	Garlic oil	Liliaceae	Bulb	expectorant, antibacterial, antifungal
<i>Azadirachta indica</i>	Neem oil	Meliaceae	leaf and seeds	Antiviral activity, antifungal activity
<i>Cuminum cyminum</i>	Jira oil	Umbelliferae	Dried ripe fruit	Stimulant, carminative, antifungal
<i>Cymbopogon martini</i>	Tikhadi oil	Poaceae	Leaves	Rheumatism and skin diseases
<i>Cymbopogon citratus</i>	Lemon grass oil	Poaceae	Leaves	Antimicrobial, Vitamin A
<i>Cinnamomum zylanicum</i>	Dalchini oil	Lauraceae	Bark and leaves	Carminative and antimicrobial
<i>Eucalyptus globulus</i>	Nilgri oil	Myrtaceae	Leaves	Antiseptic and antimicrobial
<i>Eugenia caryophyllata</i>	Clove oil	Myrtaceae	Dry flowers	Antimicrobial
<i>Elettaria cardamomum</i>	Cardamom	Zingiberaceae	Dried fruit	Carminative, antimicrobial
<i>Foeniculum vulgare</i>	Fennel	Umbelliferae	Dried fruit	Carminative, antimicrobial, expectorant
<i>Mentha spicata</i>	Mint oil	Lamiaceae	Leaves	Antimicrobial
<i>Ocimum sanctum</i>	Tulsi oil	Lamiaceae	leaves	Antibacterial, antifungal
<i>Trachyspermum captivum</i>	Ajwain	Umbelliferae	Dried fruits	Antispasmodic throat infection, bronchitis, antibacterial, antifungal
<i>Withania sominifera</i>	Ashwagandha	Solanaceae	Roots and stem	Tonic, antimicrobial
<i>Zingiber officinale</i>	Zinger oil	Zingiberaceae	Rhizome	Antimicrobial, carminative

need to search plant derived antifungal drugs which are safe and without side-effects. Hence, it is of interest to determine the scientific basis for the traditional use of medicinal plants.

The aim of this study was to assess the antifungal activity of 15 essential oils against *Aspergillus niger* and *Aspergillus fumigatus*. In the present study, 15 different medicinal plants have been chosen for the investigation of *in vitro* antifungal activity which is acts as expectorant and not having toxic properties (Table 1). The purpose of this was to create directly comparable, quantitative, antimicrobial data and to generate data for oils for which little data exist and which are used in to developed antituberculosis and antifungal drugs.

MATERIALS AND METHODS

Medicinal Plants Oils: Leaves of *Cymbopogon citratus* Stapf., *C. martinii* Roxb., *Eucalyptus globulus* Labill., *Azadirachta indica* Linn. and *Ocimum sanctum* Linn., were collected from wild fields of Amravati university, Amravati district of Maharashtra India. *Allium sativum* Linn., *Mentha spicata* Linn., *Withania sominifera* Linn., *Zingiber officinale* Roscoe., *Cinnamomum zeylanicum* Nees. collected from PKV Akola, Maharashtra India, Seeds of *Cuminum cyminum* Linn., *Trachyspermum copticum* Linn., *Elettaria cardamomum* Maton., *Foeniculum vulgare* Miller. and *Eugenia caryophyllata* Thunb., were purchased from the plant medicinal shop of Amravati maharashtra india. The seeds and roots of above plants were dried at room temperature and powdered and leaves of above plants were cut into pieces for extraction.

Extraction Procedure: The seeds, roots and leaves for oil were dried and ground to semi-powdered state. The air-dried aerial parts (50 g) were hydro distilled in a clevenger apparatus (sigma chemical company) for 5 h. in accordance with the British pharmacopoeia. The yield was 0.62% dry weight. The aqueous phase was extracted with dichloromethane (Qualigens) (3 x 50 mL). The organic phase was dried with sodium sulphate (Bio-RAD), filtered and the solvent evaporated until dryness by air-dry. The fractions obtained were combined into calibrated flasks, evaporated to dryness and weighted in order to determine the extraction's efficiency. The oils were solublized in DMSO (Bio-RAD) to a final concentration 5 mg/mL [14]. The oils were stored in a sealed glass vial (bijoux bottle) in a refrigerator at 4 °C until required. These all oils of above plants were screened for their antimycotic activity.

Fungus Selection and Growth Condition: In our survey in different hospitals and primary health center of Washim District (Central India) under the Revised National Tuberculosis Control Programmed (RNTCP) of WHO as a senior tuberculosis laboratory supervisor (STLS), we found 1357 patients of tuberculosis were recorded during June 2002 to September 2003 by sputum examination in ten PHC (primary health center) laboratories of tuberculosis in Washim District. These patients were recorded in (tuberculosis unit register) main hospital of washim. Out of 1357 patients, highly chronic pulmonary tuberculosis patients were found to be 500, which on CAT-I and CAT-II category of tuberculosis (DOTS) treatment. We found, out of 500 (CAT-I and CAT-II) patients 200 patients were suffering from mycotic infection and out of 50 patients of symptomatic TB patients' only 23 patients suffering from mycotic infection

rest of negative. The percentage of mycotic infection in pulmonary tuberculosis patients was 46.0%. Mainly for types of fungi, viz. *Aspergillus niger*, *A. fumigatus*, *Histoplasma capsulatum* and *Cryptococcus neoformans* were recovered. Culture characteristic of different fungi analysis were done using sputum sample of chronic tuberculosis patients and symptomatic patients (Non tuberculosis). Selection of *Aspergillus fumigatus* and *Aspergillus niger* on the bases of higher percentage in pulmonary tuberculosis patients.

A.fumigatus and *A. niger* were isolated from the sputum samples of pulmonary tuberculosis patients under the programme DOTS from the civil hospital of Washim, Washim District, maharashtra state, India. Deep sputum samples (2-4ml) tuberculosis patients were obtained under sterile conditions. Organisms were maintained on Czapek's dox agar (diffco) (Sodium nitrate: 02 g; Sucrose: 30 g; Dipotassium phosphate: 1g; Magnesium Sulphate: 0.5 g; Potassium Chloride: 0.5g; Ferrous Sulphate: 0.01g; Agar-Agar:15.0g; Distilled Water:1000 ml; pH:7.3 ± 0.2) in the dark at 28°C for 5-10 days. Culture confirmed by microscopy and PCR. Inocula were prepared by diluting overnight cultures in saline to approximately 10⁵ CFU ml⁻¹. These suspensions were further diluted with saline as required.

Antimycotic Assay by Disc Diffusion Technique:

Oils were screened for their antifungal activity against *A. fumigatus* and *A. niger* by disc diffusion method. Different concentration of each oil and mixed oils (equal ratio of each oil), i.e., 100 µg, 50 µg, 25 µg and 12.5 µg/disc were used for assay. The fungal cultures were grown on czapek dox broth (diffco). The mycelial mat of *A. fumigatus* and *A. niger* of 7-day old culture was washed, suspended in normal saline solution and then filtered through glass wool aseptically. The colony forming units (CFU/ml) of suspension of the test fungus was determined and test inoculum was adjusted 1-5 X 10⁵ ml. These conidia were used for antifungal assay tests. Inocula (0.1ml) were applied on the surface of the Czapek's dox agar (Diffco) plate and spread by using sterile glass spreader. The sterile discs (5mm diameter, Whatmans filter paper No.42) were soaked in added concentrations (100µg / disc, 50µg /disc, 25µg /disc and 12.5µg / disc) of essential oils. The test was performed in triplicate. These dishes were incubated for 48 h at 28°C. Zone of inhibition in mm were determined after 48 h. Standard antibiotics miconazole (Sigma) and itraconazole (Sigma) were used in order to control the sensitivity of the tested fungus. Miconazole was used as positive control because miconazole and itraconazole those

that followed such as itraconazole, fluconazole and itraconazole proved to be important drugs for combating human fungal infections. The clinical efficiency and safety of itraconazole in particular has resulted in wide spread use. The resultant emergence of resistant to azoles has intensified the search for new compounds that are active against resistant organisms [15-18]. DMSO (Sigma) as negative control and filter paper disc which is soaked with distilled water no oil is used as negative control. The zones of different oil were measured. the data of all the parameters were statistically analyzed.

Agar Dilution Method: The agar dilution method followed that approved by the NCCLS with the following modification: a final concentration of 0.5% (v/v) Tween-20 (Sigma) was incorporated into the agar after autoclaving to enhance oil solubility. Briefly, a series of twofold dilutions of each oil, ranging from 2% (v/v) to 0.03% (v/v), was prepared in sabouraud dextrose agar (SDA) (Himedia) with 0.5% (v/v) Tween-20. Plates were dried at 35 °C for 30 min prior to inoculation with 1–2 ml spots containing approximately 10⁵ CFU of each organism. SDA, with 0.5% (v/v) Tween-20 but no oil, was used as a positive growth control. Inoculated plates were incubated at 35°C for 48 h. Minimum inhibitory concentrations (MICs) were determined after 48 h for *Aspergillus fumigatus* and *Aspergillus niger*. The MICs were determined as the lowest concentration of oil inhibiting the visible growth of each organism on the agar plate. The presence of one or two colonies was disregarded.

Broth Micro Dilution Method: The broth micro dilution assay was performed as described previously [19] with the following modifications: Sabouraud dextrose agar (SDA) (Himedia) was used, in tests with *Aspergillus fumigatus* and *A. niger*, sub-cultures were performed after 48 h incubation. For most oils, the highest concentration tested was 4.0% (v/v), although for some this was 8.0% (v/v). The lowest concentration tested was 0.008% (v/v).

Statistical Analysis: Minimum inhibitory concentrations of the essential oils were compared using the Wilcoxon signed rank test (Graph Prism software for Win-dows, GraphPad Software, San Diego, USA).

RESULTS

The antifungal activities of fifteen plant oils obtained by the disc diffusion method are shown in Table 2. All the oils tested exhibited different degrees of antifungal activity against *Aspergillus fumigatus* and *A. niger*

Table 2: Antifungal activity of the fifteen oils of medicinal plants against *Aspergillus fumigatus* and *Aspergillus niger* growth *in vitro*

S. No.	Plant oils	Aspergillus fumigatus 100 50 25 12.5µg				Aspergillus niger 100 50 25 12.5µg			
		/disc /Zone of inhibition (mm)	/disc /Zone of inhibition (mm)	/disc /Zone of inhibition (mm)	/disc /Zone of inhibition (mm)	/disc /Zone of inhibition (mm)	/disc /Zone of inhibition (mm)	/disc /Zone of inhibition (mm)	/disc /Zone of inhibition (mm)
1	Allium sativum (Garlic oil)	15±0.5	13±0.5	12±0.1	10±0.1	14±0.5	12±0.5	10±1.1	8±0.5
2	Azadirachta indica (Neem oil)	16±0.5	13±0.5	12±1.1	10±1.1	14±0.5	13±0.5	11±1.1	8±0.5
3	Cuminum cyminum (Jira Oil)	13±0.5	12±0.5	10.5±0.5	10±0.5	12±0.5	11±0.1	10±0.5	9±0.5
4	Cymbopogon martinii (Tikhadi oil)	24±0.5	22±0.5	21±0.5	19±0.5	22±0.5	20±0.5	19±0.5	15±0.5
5	Cymbopogon citratus (Lemon- Grass Oil)	22±0.5	20±0.5	19±0.5	18±0.5	21±0.5	20±0.5	18±0.5	18±0.5
6	Cinnamomum zylanicum (Kalmi-Dalchini oil)	19±0.5	17±0.5	16±0.5	12±0.5	19±0.5	17±0.5	16±0.5	12±0.5
7	Eucalyptus globulus (Nilgri oil)	22±0.0	20±0.5	19±0.5	18±0.5	21±1.0	20±1.0	19±0.5	18±0.5
8	Eugenia caryophyllata (Clove oil)	15±0.5	13±0.5	12±1.1	10±1.1	14±0.5	12±0.5	10±0.5	8±0.1
9	Elettaria cardamomum (Cardamom seeds)	13±0.5	12±0.5	10.5±0.5	10±0.5	12±0.5	11±0.5	10±0.5	9±0.5
10	Foeniculum vulgare (Fennel fruits oil)	13±0.5	12±0.5	10.5±0.5	10±0.5	12±0.5	11±0.5	10±0.5	9±0.5
11	Mentha spicata (Mint oil)	16±0.5	13±0.5	12±0.5	10±0.5	14±0.5	13±0.5	11±0.5	8±0.5
12	Ocimum sanctum (Tulsi oil)	13±0.5	12.5±0.5	11±0.5	9±0.5	12±0.5	10±0.1	9±0.5	8±0.5
13	Trachyspermum copticum (Ajowan)	13±0.5	12±0.5	10.5±0.5	10±0.5	12±0.5	11±0.5	10±0.5	9±0.5
14	Withania somnifera (Ashwagandha oil)	15±0.5	13±0.5	11±1.1	10±1.1	14±0.5	12±0.5	10±1.1	8±0.0
15	Zingiber officinale (Zinger oil)	15±0.5	13±0.5	12±1.1	10±0.0	14±0.5	12±0.5	10±1.1	8±0.0
16	Mixed oils C. martinii, C. citratus, E. globulus and C. zylanicum)	22±0.5	19±0.5	18±0.5	16±0.5	24±0.5	21±0.5	19±0.5	16±0.5
17	DMSO negative control	No Zone				No Zone			
18	Miconazole (positive control)	24±0.5	20±0.5	19±0.5	18±0.5	25±0.5	22±0.5	20±0.5	18±0.5
19	itraconazole	23±0.5	18±0.5	16±0.5	12±0.5	21±0.5	15±0.5	12±0.5	10±0.5

Table 3: Minimum inhibitory concentrations (MICs) of selected essential oils (%v/v) against *Aspergillus fumigatus* and *Aspergillus Niger* using agar diffusion method

Plant oils	<i>Aspergillus</i>	
	<i>fumigatus</i>	<i>niger</i>
Allium sativum (Garlic oil)	>2	>2
Azadirachta indica (Neem oil)	0.25	0.5
Cuminum cyminum (Jira Oil)	>2	>2
Cymbopogon martinii (Palmarosa)	0.06	0.06
Cymbopogon citratus (Lemon- Grass Oil)	0.06	0.06
Cinnamomum zylanicum (Kalmi-Dalchini oil)	0.12	0.12
Eucalyptus globulus (Nilgri oil)	0.12	0.12
Eugenia caryophyllata (Clove oil)	0.12	0.12
Elettaria cardamomum (Cardamom seeds)	>2	>2
Foeniculum vulgare (Fennel fruits oil)	0.5	0.5
Mentha spicata (Mint oil)	0.5	0.25
Ocimum sanctum (Tulsi oil)	0.5	0.5
Trachyspermum copticum (Ajowan)	0.5	0.5
Withania somnifera (Ashwagandha oil)	0.5	0.5
Zingiber officinale (Zinger oil)	>2	>2
Mixed oils (C. martinii, C. citratus, E. globules and C. zylanicum)	0.03	0.03

isolated from pulmonary tuberculosis patients. The maximum antimycotic activity was shown by *C. martinii* followed by *C. citratus*, *Eucalyptus globulus* and *Cinnamomum zylanicum*. The oils of *Mentha spicata*, *Azadirachta indica*, *Eugenia caryophyllata*, *Withania somnifera* and *Zingiber officinale* exhibited moderate

activity and the oils of *Cuminum cyminum*, *Allium sativum*, *Ocimum sanctum*, *Trachyspermum copticum*, *Foeniculum vulgare* and *Elettaria cardamomum* showed comparatively low activity against *A. niger* and *A. fumigatus*.

The MICs of 15 plant oils obtained by the agar dilution method are shown in Table 3. *Cymbopogon martinii* (Palmarosa), *Cymbopogon citratus* (Lemon-Grass Oil), *Eucalyptus globulus* (Nilgri oil) *Eugenia caryophyllata* (Clove oil) and *Cinnamomum zylanicum* (Kalmi-Dalchini oil) inhibited *Aspergillus fumigatus* and *Aspergillus niger* at = 0.25% (v/v). *Azadirachta indica* (Neem oil), *Foeniculum vulgare* (fennel oil), *Mentha spicata* (Mint oil), *Ocimum sanctum* (Tulsi oil) and *Withania somnifera* (Ashwagandha oil) inhibited *Aspergillus* and *Aspergillus niger* at = 2.0% (v/v). Five oils (*Allium sativum* (Garlic oil), *Cuminum cyminum* (Jira Oil), *Elettaria cardamomum* (Cardamom seed), *Trachyspermum copticum* (Ajowan) and *Zingiber officinale* (Zinger oil) failed to inhibit any organisms at the highest concentration, which was 2.0% (v/v).

Table 4 shows MICs and minimum inhibitory concentrations (MICs) of 15 plant oils and mixed oils (equal ratio of *C. martinii*, *C. citratus* *Eucalyptus globulus* and *Cinnamomum zylanicum*) obtained by the broth micro dilution method. *Cymbopogon martinii* and *Cymbopogon citratus* had lowest MICs of (0.03%v/v) against *Aspergillus fumigatus* and *Aspergillus niger*.

Table 4: Minimum inhibitory concentration and minimum cidal concentration data (% v/v) of *A. fumigatus* and *A. niger* obtained by the broth micro dilution method

Plant oils	Aspergillus fumigatus		Aspergillus niger	
	MIC	MCC	MIC	MCC
Allium sativum (Garlic oil)	>4.0	>4.0	>4.0	>4.0
Azadirachta indica (Neem oil)	0.12	0.12	0.25	0.25
Cuminum cyminum (Jira Oil)	>4	4	>4	>4
Cymbopogon martini (Palmarosa)	0.03	0.03	0.03	0.03
Cymbopogon citratus (Lemon- Grass Oil)	0.03	0.03	0.03	0.03
Cinnamomum zylanicum (Kalmi-Dalchini oil)	0.12	0.12	0.12	0.12
Eucalyptus globulus (Nilgri oil)	0.06	0.06	0.06	0.06
Eugenia caryophyllata (Clove oil)	0.06	0.12	0.12	0.06
Elettaria cardamomum (Cardamom seeds)	>8.0	>8.0	>8.0	>8.0
Foeniculum vulgare (Fennel fruits oil)	0.5	1.0	0.5	0.5
Mentha spicata (Mint oil)	0.12	0.12	0.12	0.12
Ocimum sanctum (Tulsi oil)	0.5	1.0	0.5	1.0
Trachyspermum copticum (Ajowan)	>8	>8	1.0	1.0
Withania somnifera (Ashwagandha oil)	1.0	0.5	1.0	0.5
Zingiber officinale (Zinger oil)	>4	>4	>4	>4
Mixed oils (<i>C. martinii</i> , <i>C. citratus</i> , <i>E. globulus</i> and <i>C. zylanicum</i>)	0.008	0.008	0.06	0.06

Mixed oils had lowest MICs of 0.008% (v/v) against *Aspergillus fumigatus* 0.06% (v/v) against *Aspergillus niger*. Comparison of MICs obtained by agar and broth methods showed that differences exceeding two serial dilutions were seen with *Mentha spicata*, *Allium sativum*, *Azadirachta indica* (Neem oil), *Cuminum cyminum*, *Eucalyptus globulus* and *Eugenia caryophyllata*. The greatest difference was for *Aspergillus fumigatus* against with *Trachyspermum copticum*, where the MIC obtained by agar dilution was 0.5 % (v/v) compared with the MIC by micro dilution >8.0 % (v/v). The lowest minimum inhibitory concentrations were 0.008 % (v/v) of mixed oils against *Aspergillus fumigatus*.

DISCUSSION

The traditional use of plants as medicines provide the basis for indicating which essential oils and plant oils may be useful for specific medical conditions. Historically, many plant oils and extracts, such as tea tree, myrrh and clove, have been used as topical antiseptics, or have been reported to have antimicrobial properties [20,21]. It is important to investigate scientifically those plants which have been used in traditional medicines as potential sources of novel antimicrobial compounds [22]. Also, the resurgence of interest in natural therapies and increasing consumer demand for effective, safe, natural products means that quantitative data on plant oils and extracts are

required. Various publications have documented the antimicrobial activity of essential oils and plant extracts including rosemary, peppermint, bay, basil, tea tree, celery seed and fennel [23-27]. All the oils tested exhibited different degrees of antifungal activity against *Aspergillus fumigatus* and *A. niger*. The maximum antimycotic activity was shown by *C. martinii* followed by *C. citratus*, *Eucalyptus globulus* and *Cinnamomum zylanicum*. Aggarwal, *et al.* [28] reported antimycotic activity of *C. martinii* against *A. niger*. The oil of *C. citratus* was effective against fungal pathogens causing diseases in plants and human beings [29]. Quale, *et al.* [30] treated infections caused by *Candida* in AIDS patients with a drug based on Cinnamon. In our study we also found that essential oil extracted from *Cinnamomum zylanicum* demonstrated strong antifungal activity on both the species of *Aspergillus*. The antimycotic activity of cinnamon bark due to presence of cinnamaldehyde is well known [31]. Similarly, *in vitro* antimicrobial activity of *Cinnamomum zylanicum* (bark) against human pathogenic fungi and commensally bacteria was studied by Chaumont *et al.* [32] and Matan *et al.* [33]. The oils of *Mentha spicata*, *Azadirachta indica*, *Eugenia caryophyllata*, *Withania somnifera* and *Zingiber officinale* exhibited moderate activity. The essential oil of mint was found to have strong antimycotic activity against *Candida albicans* Kishore *et al.* [34] also reported that essential oil of mint showed high antimycotic activity against dermatophytes. The oils of *Cuminum cyminum*, *Allium sativum*, *Ocimum sanctum*, *Trachyspermum copticum*, *Foeniculum vulgare* and *Elettaria cardamomum* showed comparatively low activity against *A. niger* and *A. fumigatus* as compared to control. Nigam and Rao [35] reported that oil of *C. cyminum* was toxic to *Aspergillus*. Similarly, Singh N, [3] reported high antifungal activity of *C. cyminum*. Oil of *Allium sativum* showed significant growth inhibition of fungi including dermatophytes [36]. Singh, [37] found antimycotic activity in oil of *Ocimum sanctum* against pathogenic fungi. Plant oils are important source of fungitoxic compounds and they may provide a renewable source of useful fungicides that can be utilized in antimycotics drugs against *A. fumigatus* and *A. niger* infection in patients suffering from pulmonary tuberculosis. Among the plant oils tested, *C. martinii*, *C. citratus*, *E. globulus* and *C. zylanicum* showed high antimycotic activity. Mixed oils were inhibited the *Aspergillus fumigatus* and *Aspergillus niger* at lowest MICs. These results support the notion that plant essential oils have a role as pharmaceuticals and preservatives.

When comparing data obtained in different studies, most publications provide generalizations about whether or not plant oil possesses activity against fungi. However, not all provide details about the extent or spectrum of this activity. Some publications also show the relative activity of plant oils and extracts by comparing results from different oils tested against the same organism(s). Comparison of the data obtained in this study with previously published results is problematic. First, the composition of plant oils and extracts is known to vary according to local climatic and environmental conditions [38,39]. Furthermore, some oils with the same common name may be derived from different plant species [40]. Secondly, the method used to assess antimicrobial activity and the choice of test organism(s), varies between publications [38]. A method frequently used to screen plant extracts for antimicrobial activity is the agar disc diffusion technique [23,41]. The usefulness of this method is limited to the generation of preliminary, qualitative data only, as the hydrophobic nature of most essential oils and plant extracts prevents the uniform diffusion of these substances through the agar medium [38,24]. Agar and broth dilution methods are also commonly used. The results obtained by each of these methods may differ as many factors vary between assays [28,26]. These include differences in microbial growth, exposure of microorganisms to plant oil, the solubility of oil or oil components and the use and quantity of an emulsifier. These and other elements may account for the large differences in MICs obtained by the agar and broth dilution methods in this study. *In vivo* studies may be required to confirm the validity of some of the results obtained. The need for a standard, reproducible method for assessing oils has been stressed by several authors [42,43]. In view of this, many methods have been developed specifically for determining the antimicrobial activity of essential oils [42,45]. The benefits of basing new methods on preexisting, conventional assays such as the NCCLS methods are that these assays tend to be more readily accepted by regulatory bodies [42,45]. Also, these methods have been designed specifically for assessing the activity of antimicrobial compounds and factors affecting reproducibility have been sufficiently investigated. Although NCCLS methods have been developed for assessing conventional antimicrobial agents such as antibiotics, with minor modifications these methods can be made suitable for the testing of essential oils and plant extracts [42]. For some plant oils, such as wintergreen, eucalyptus, clove and sage, there has been much research and reporting of toxic and irritant properties [21,40]. In spite of this, most of these oils are

available for purchase as whole oils or as part of pharmaceutical or cosmetic products, indicating that toxic properties do not prohibit their use. However, the on-going investigation of toxic or irritant properties is imperative, especially when considering any new products for human use, be they medicinal or otherwise. In summary, this study confirms that many essential oils possess *in vitro* antifungal activity. However, if plant oils are to be used for preservation or medicinal purposes, issues of safety and toxicity will need to be addressed. This obviously justifies the use of above mixed oils in traditional medicine to cure mycotic infections in tuberculosis patients.

ACKNOWLEDGEMENTS

The authors thank to Dr. Mane, District tuberculosis Officer (DTO), Dr. Jadhav, Tuberculosis unit officer, Supervisors, Technicians of civil hospital and ten PHC of washim district, Maharashtra, India for providing tuberculosis patients records, help in sputum examination and isolation of fungi and maintenance of records during survey. This work was supported by Dr. Varma (MD, BAMS), Murtizapur, Akola district Maharashtra INDIA.

REFERENCES

1. Davies, J., 1994. Inactivation of antibiotic and the dissemination of resistance genes. *Sci.*, 264: 375-382.
2. Bauer, A.W., W.M. Kirby, J.C. Sherris and M. Turck, 1996. Antibiotic susceptibility testing by standardized single disc method. *Am. J. Clin Pathol.*, 44: 493-496.
3. Singh, N., 2001. Trends in the epidemiology of opportunistic fungal infections: predisposing factors and the impact of antimicrobial use practices. *Clinical Infect Diseases*, 33(10): 1692-1696.
4. Alex, L.A., C. Korch, C.P. Selitrennikoff and M.I. Simon, 1998. COS1, a two-component histidine kinase that is involved in hyphal development in the opportunistic pathogen *Candida albicans*. *Proc. Natl. Acad. Sci. USA*, 95: 7069-7073.
5. Ali-Shtayeh, M.S. and S.I. Abu Ghdeib, 1999. Antimycotic activity of twenty-two plants used in folkloric medicine in the Palestinian area for the treatment of skin diseases suggestive of dermatophyte infection. *Mycoses*, 42: 665-672.
6. Soliman, K.M. and R.I. Badaea, 2002. Effect of oil extracted from some medicinal plants on different mycotoxigenic fungi. *Food Chem Toxicol.*, 40(11): 1669-75.

7. Thoppil, J.E., J. Minija, A. Tajo and M.J. Deena, 2003. Antimicrobial activities of *Eusteralis deccanensis* and *E. quadrifolia* essential oils. J. Environ. Biol., 24(2): 211-2.
8. Govinden-Soulange, J., N. Magan, A. Gurib-Fakim, A. Gauvin, J. Smadja and H. Kodja, 2004. Chemical composition and *in vitro* antimicrobial activities of the essential oils from endemic *Psiadia* species growing in mauritius. Biol Pharm Bull., 27(11): 1814-8.
9. Romagnoli, C., R. Bruni, E. Andreotti, M.K. Rai, C.B. Vicentini and D. Mares, 2005. Chemical characterization and antifungal activity of essential oil of capitula from wild Indian *Tagetes patula* L. Protoplasma, pp: 57-65.
10. Pinto, E., C. Pina-Vaz, L. Salgueiro, M.J. Goncalves, S. Costa-de-Oliveira, C. Cavaleiro, A. Palmeira, A. Rodrigues and J. Martinez-de-Oliveira, 2006. Antifungal activity of the essential oil of *Thymus pulegioides* on *Candida*, *Aspergillus* and dermatophyte species. J. Med. Microbiol., 55(Pt 10): 1367-73.
11. Tabanca, N., B. Demirci, S.L. Crockett, K.H. Baser and D.E. Wedge, 2007. Chemical composition and antifungal activity of arnica longifolia, Aster hesperius and Chrysothamnus nauseosus Essential Oils. J Agric. Food Chem., 55(21): 8430-8435.
12. Tullio, V., A. Nostro, N. Mandras, P. Dugo, G. Banche, M.A. Cannatelli, A.M. Cuffini, V. Alonzo and N.A. Carlone, 2007. Antifungal activity of essential oils against filamentous fungi determined by broth microdilution and vapour contact methods. J. Appl. Microbiol., 102(6): 1544-50.
13. Dutta, B.K., S. Karmakar, A. Naglot, J.C. Aich and M. Begam, 2007. Anticandidial activity of some essential oils of a mega biodiversity hotspot in India., Mycoses, 50(2): 121- 4.
14. Glowniak, K., T. Wolski and T. Dragan, 1991. Sposób wydobrania Kumaryn Z Surowców rocinnych zwlaszcza Z owoców arcydziegla olesnika. Pat pol., 15: 1-5.
15. Hata, K., J. Ueno and H. Miki, 1995. ER-30346, a novel antifungal triazole. III. *In vitro* activity and the mode of action, abstr. In Program and Abstracts of the 35th Interscience Conference on Antimicrobial Agents and Chemotherapy. American Society for Microbiology, Washington, D.C. Abstract F92:129.
16. Perfect, J.R. and W.A. Schell, 1995. *In vitro* efficacy of the azole, SCH 56592, compared to amphotericin B, fluconazole and itraconazole versus *Cryptococcus neoformans*., In Program and Abstracts of the 35th Interscience Conference on Antimicrobial Agents and Chemotherapy. American Society for Microbiology, Washington, D.C. Abstr. F64:124.
17. Law, D. and D.W. Denning, 1996. *In vitro* activity of Schering 56592, compared with fluconazole and itraconazole against *Candida* spp. In Program and Abstracts of the 36th Interscience Conference on Antimicrobial Agents and Chemotherapy. American Society for Microbiology, Washington, D.C. Abstract F88:115.
18. Munayyer, H., K.J. Shaw, R.S. Hare, B. Salisbury, L. Heimark, B. Pramanik and J.R. Greene, 1996. SCH 56592 is a potent inhibitor of sterol C-14 demethylation in fungi. In Program and Abstracts of the 36th Interscience Conference on Antimicrobial Agents and Chemotherapy. American Society for Microbiology, Washington, D.C. Abstract.F92: 115.
19. Hammer, K.A., C.F. Carson and T.V. Riley, 1996. Susceptibility of transient and commensal skin flora to the essential oil of Melaleuca alternifolia (tea tree oil). Am. J. Infection Control, 24: 186-189.
20. Hoffman, D.L., 1987. The Herb User's Guide. Wellingborough, UK: Thorsons Publishing Group.
21. Lawless, J., 1995. The Illustrated Encyclopedia of Essential Oils. Shaftesbury, UK: Element Books Ltd.
22. Mitscher, L.A., S. Drake, S.R. Gollapudi and S.K. Okwute, 1987. A modern look at folkloric use of anti-infective agents. J. Natural Products, 50: 1025-1040.
23. Morris, J.A., A. Khettry and E.W. Seitz, 1979. Antimicrobial activity of aroma chemicals and essential oils. J. Am. Oil Chemists' Soc., 56: 595-603.
24. Ross, S.A., N.E. El-Keltawi and S.E. Megalla, 1980. Antimicrobial activity of some Egyptian aromatic plants. Fitoterapia, 51: 201-205.
25. Yousef, R.T. and G.G. Tawil, 1980. Antimicrobial activity of volatile oils. Die Pharmazie. 35: 698-701.
26. Hili, P., C.S. Evans and R.G. Veness, 1997. Antimicrobial action of essential oils: the effect of dimethylsulfoxide on the activity of cinnamon oil. Letters in Appl. Microbiol., 24: 269-275.
27. Lis-Balchin, M. and S.G. Deans, 1997. Bioactivity of selected plant essential oils against *Listeria monocytogenes*. J. Appl. Bacteriol., 82: 759-762.

28. Aggarwal, K.K., A. Ahmad, T.R. Santha, N. Jain, V.K. Gupta, K. Sushil and S.P. Khanuja, 2000. Antimicrobial activity spectra of *Pelargonium graveolens* L. and *Cymbopogon winterianus* Jowitt. oil constituents and acyl derivatives. J. Med. Aromatic Plant Sci., 22: 544-548.
29. Singh, H.B., 2000. Antifungal efficiency of some essential oils against *Sclerotium rolfsii*. Ind. Perf., 44: 71-73.
30. Quale, J.M., D. Landman, M.Z. Zaman, S. Burney and S. Sathe, 1996. *In vitro* activity of *Cinnamomum zeylanicum* against azole resistant and sensitive *Candida* Species and a pilot study of Cinnamon for oral Candidiasis. Am. J. Chinese Med., 24: 103-109.
31. Viollon, C. and J.P. Chaumont, 1994. Antifungal properties of essential oils and their main components upon *Cryptococcus neoformans*. Mycopathologia. 128: 151-153.
32. Chaumont, J.P., 2003. Antimycotic essential oils: Impact on skin micro flora, in Plant-Derived Antimycotics: Current Trends and Future Prospects (Eds M.K. Rai and D. Mares). Haworth press, USA, 357-364.
33. Matan, N., H. Rimkeeree, A.J. Mawson, P. Chompreeda, V. Haruthaithanasan and M. Parker, 2006. Antimicrobial activity of cinnamon and clove oils under modified atmosphere conditions. Intl. J. Food Microbiol., 107(2): 180-5.
34. Kishore, N., A.K. Mishra and J.P. Chansouria, 1993. Fungal toxicity of essential oils against dermatophytes. Mycoses. 36: 211-225.
35. Nigam, S.S. and T.S.T. Rao, 1977. Antimicrobial efficacy of some Indian essential oils. In L.D. Kapoor and Ramakrishnan, (Eds), Advance in essential oil Industry, New Delhi: Today's and Tomorrow's printer and publishers, pp: 177-180.
36. Singh, R. and B. Rai, 2000. Antifungal potential of some higher plants against *Fusarium* causing with diseases of *Cajanus Cajan*. Microbios., 102: 165-173.
37. Singh, R., 1998. Antifungal efficacy of volatile constituents of higher plants against sugarcane fungal pathogens. Ind. Perf., 42(2): 82-85.
38. Janssen, A.M., J.J.C. Scheffer and A. Baerheim Svendsen, 1987. Antimicrobial activity of essential oils: a 1976-86 literature review. Aspects of the test methods. Planta Medica., 53: 395-398.
39. Sivropoulou, A., S. Kokkini, T. Lanaras and M. Arsenakis, 1995. Antimicrobial activity of mint essential oils. J. Agril. Food Chem., 43: 2384-2388.
40. Reynolds, J.E.F., 1996. Martindale the Extra Pharmacopoeia 31st edn. London: Royal Pharmaceutical Society of Great Britain.
41. Smith-Palmer, A., J. Stewart and L. Fyfe, 1998. Antimicrobial properties of plant essential oils and essences against five important food-borne pathogens. Letters in Appl. Microbiol., 26: 118-122.
42. Carson, C.F., K.A. Hammer and T.V. Riley, 1995. Broth micro dilution method for determining the susceptibility of *Escherichia coli* and *Staphylococcus aureus* to the essential oil of *Melaleuca alternifolia* (tea tree oil). Microbios, 82: 181-185.
43. Mann, C.M. and J.L. Markham, 1998. A new method for determining the minimum inhibitory concentration of essential oils. J. Appl. Microbiol., 84: 538-544.
44. Remmal, A., T. Bouchikhi, K. Rhayour, M. Ettayebi, and A. Tantaoui-Elaraki, 1993. Improved method for the determination of antimicrobial activity of essential oils in the agar medium. J. Essential Oil Res., 5: 179-184.
45. Smith, M.D. and P.L. Navilliat, 1997. A new protocol for antimicrobial testing of oils. J. Microbiol. Methods, 28: 21-24.