

## Analysis of Chemical Composition of Leaves and Roots of *Ageratum conyzoides*

<sup>1</sup>K.N. Agbafor, <sup>2</sup>A.G. Engwa and <sup>2</sup>I.K. Obiudu

<sup>1</sup>Biochemistry Department, Ebonyi State University, Abakaliki, Nigeria

<sup>2</sup>Chemical Sciences Department, Godfrey Okoye University, Enugu, Nigeria

**Abstract:** *Ageratum conyzoides* has been utilized by herbal medicine practitioners in Eastern Nigeria in treatment and management of several disorders, such as; diabetes, inflammations, spasm, headaches, cancers, among others. The basis of these applications has not been well investigated. This present research studied the phytochemical, proximate, mineral and vitamin composition of leaves and roots of the plant. The chemical analyses were done according to standard procedures. Alkaloids, flavonoids, saponins, tannins, cardiac glycosides and anthraquinones were detected in both samples, while terpenoids were present in the leaves only. The values of these phytochemicals in the leaves were significantly higher ( $P < 0.05$ ) than those in the roots. Proximate analysis indicated that the leaves contained ash (12.64%), crude fat (2.27%), fiber (23.50%), moisture (10.02%), crude protein (14.73%) and carbohydrate (36.84%), while ash (16.95%), crude fat (2.01%), fiber (26.74%), moisture (13.35%), crude protein (9.89%) and carbohydrate (31.06%) were found in the roots. Na, K, Ca, Mg, P, Zn, Mn and Fe were present in the leaves and roots and their concentrations in roots were significantly higher ( $P < 0.05$ ) than in leaves. Both samples were found to contain vitamins B<sub>2</sub>, B<sub>6</sub>, C, E, thiamin and niacin. Their values in the leaves did not differ significantly ( $P > 0.05$ ) from those in roots. These chemical constituents of leaves and roots of *Ageratum conyzoides* may be responsible for their applications in treatment and management of various disorders.

**Key words:** Phytochemicals • Proximate composition • Minerals • Vitamins • *Ageratum conyzoides*

### INTRODUCTION

The use of plants and their products for different purposes has been with man from the beginning. Aside food, plants are often used as medicine [1-3]. All plants produce chemical compounds as part of their normal metabolic activities. These can be divided into primary metabolites such as sugars and fats found in all plants and secondary metabolites, e.g. alkaloids, tannins, saponins, phenols, etc. These secondary metabolites and other chemical constituents of medicinal plants account for their medicinal value [4]. Medicinal plants often exhibit a wide range of biological and pharmacological activities such as; anti-inflammatory, anti-bacterial and anti-fungal properties [5,6]. Extracts, syrups, infusions and concoctions prepared from different parts of these plants are used to remedy different ailments. Such ailments include; typhoid, anaemia, malaria, headache, etc [7]. In Nigeria many diseases were treated and are still being treated with medicinal plants with success. These diseases include malaria, epilepsy, infantile convulsion

diarrhoea, dysentery, bacterial and fungal infections, mental illness, asthma, diabetes, worm infestation, pains and ulcers [8].

*Ageratum conyzoides* belongs to the family of Asteraceae. It is an erect, annual, branched, slender, hairy and aromatic plant which grows to approximately one meter in height. It is a native of central America, Southeast Asia, South China, India, West Africa, etc [1,9]. It has been known since ancient times for its curative properties and has been utilized for the treatment of various ailments, such as burns and wounds, diabetes, headaches, pneumonia, inflammation, asthma, spasmodic and haemostatic diseases, stomach ailments, gynecological diseases, leprosy and other skin diseases [10].

The medicinal applications of various parts of *Ageratum conyzoides* are yet to be fully investigated. In this line of communication, the present research studied and investigated the chemical constituents of leaves and roots of *Ageratum conyzoides*. The pharmacological significance of the chemical composition has been discussed.

## MATERIALS AND METHODS

**Collection and Preparation of Plant Materials:** Fresh leaves and roots of *Ageratum conyzoides* were collected from Okposi in Ohaozara L.G.A of Ebonyi state. The samples were identified and authenticated by Prof. S.C Onyekwelu, a botanist in Applied Biology Department of Ebonyi State University, Abakaliki. They were air-dried and ground into a powder which was stored in an air-tight contain.

**Measurement of Phytochemicals:** Qualitative determination of saponins, alkaloids, tannins, cardiac glycosides, terpenoids, flavonoids and anthraquinones was carried out on the extracts using the method described by Sofowora [11]. The quantitative measurement of the phytochemicals was done according to standard procedures of Trease and Evans [12] and Harbone [13].

**Proximate Analysis:** Proximate analysis was carried out according to the procedure of Association of Official Analytical Chemists [14]. This constitutes the class of food present in samples such as carbohydrate, protein, fat, fibre, ash content and moisture content.

**Determination of Mineral Composition:** The extracts were incinerated into ash, dissolved in 1ml of 2M HCl and diluted to 100ml with deionized water. The resulting solution was used for the determination of Na, Ca, Mg, Mn, Fe, Zn, K, Cr and Cu using atomic absorption spectrophotometer (Buck scientific AAS 200 A) [15]. Phosphorous was measured by the vanadiumolybdate colorimetric method of Pearson [16].

**Measurement of Vitamin Content:** The levels of vitamins A, K, thiamin, C, B<sub>1</sub>, B<sub>2</sub>, B<sub>6</sub>, E and niacin in the extracts were determined according to the methods described by Okwu and Ndu [17].

**Statistical Analysis:** Data generated were expressed as mean  $\pm$  SD. Statistical significance of difference was determined using the program SPSS 12 (SPSS, USA) by performing one-way ANOVA with post-hoc comparisons between the control group and each of the treated groups by Duncan's multiple comparison test. A *p*-value less than 0.05 was considered statistically significant.

## RESULT AND DISCUSSION

The results of the phytochemical screening are presented in table 1. Alkaloids, tannins, saponins, cardiac glycosides, anthraquinones and flavonoids were found in

Table 1: Phytochemical composition of leaves and roots of *Ageratum conyzoides*

| Pytochemicals      | Leaf (mg/100g)                | Root (mg/100g)                |
|--------------------|-------------------------------|-------------------------------|
| Alkaloids          | 26.80 $\pm$ 1.08 <sup>a</sup> | 13.04 $\pm$ 1.14 <sup>d</sup> |
| Flavonoids         | 21.24 $\pm$ 1.85 <sup>a</sup> | 9.50 $\pm$ 2.09 <sup>d</sup>  |
| Tannins            | 4.78 $\pm$ 0.31 <sup>b</sup>  | 2.05 $\pm$ 0.65 <sup>e</sup>  |
| saponins           | 3.16 $\pm$ 0.55 <sup>b</sup>  | 2.02 $\pm$ 0.12 <sup>e</sup>  |
| Cardiac glycosides | 3.05 $\pm$ 0.28 <sup>b</sup>  | 1.60 $\pm$ 0.45 <sup>e</sup>  |
| Terpenoids         | 0.84 $\pm$ 0.06 <sup>c</sup>  | nil                           |
| Anthraquinones     | 0.79 $\pm$ 0.11 <sup>c</sup>  | 0.34 $\pm$ 0.09 <sup>f</sup>  |

Values are mean  $\pm$  SD; n = 3. Values with different superscripts are significantly different (*P*<0.05).

Table 2: Proximate composition of dried leaves and roots of *Ageratum conyzoides*

| Component          | Leaf (%)                      | Root (%)                      |
|--------------------|-------------------------------|-------------------------------|
| Crude carbohydrate | 36.84 $\pm$ 2.05 <sup>a</sup> | 31.06 $\pm$ 1.25 <sup>a</sup> |
| Protein            | 14.73 $\pm$ 1.55 <sup>b</sup> | 9.89 $\pm$ 0.70 <sup>b</sup>  |
| Fibre              | 23.50 $\pm$ 1.02 <sup>c</sup> | 26.74 $\pm$ 0.46 <sup>c</sup> |
| Fat                | 2.27 $\pm$ 0.12 <sup>d</sup>  | 2.01 $\pm$ 0.16 <sup>d</sup>  |
| Ash                | 12.64 $\pm$ 0.64 <sup>b</sup> | 16.95 $\pm$ 1.56 <sup>b</sup> |
| Moisture           | 10.02 $\pm$ 1.15 <sup>b</sup> | 13.35 $\pm$ 2.02 <sup>b</sup> |

Values are mean  $\pm$  SD; n = 3. Values with different superscripts are significantly different (*P*<0.05).

the leaf and root samples, while the presence of terpenoids was detected in the leaves only. This is consistent with the findings of Duru *et al.* [18]. The concentrations of the phytochemicals in the leaf sample were significantly higher (*P*<0.05) than those in the roots. The medicinal uses of *Ageratum conyzoides* may be partly attributed to the identified constituent phytochemicals. Alkaloids have analgesic effects [17]. Morphine alkaloids are powerful pain relievers and narcotics (induces sleep or drowsiness). Atropine, cocaine and other alkaloids are known stimulants of the central nervous system [19]. Tannins prevent urinary tract infection by preventing bacteria from adhering to the walls. Tannins have been shown to be useful in the management of HIV infection and herpes. Combination of tannin and anthocyanins can breakdown cholesterol in the bloodstream and in atherosclerotic plaques. Tannins, along with vitamin C help build and strengthen collagen [20]. Saponins serve as natural antibiotics, which help the body to fight infections and microbial invasions. They also enhance the effectiveness of certain vaccines, lower cholesterol and knock out some tumor cells, particularly lung and blood cancers [17]. Flavonoids act as antioxidants in biological systems. Other properties of flavonoids include protection against allergies, inflammation, free radicals, platelet aggregation microbes, ulcers, hepatoxins, viruses and tumors [17]. The other phytochemicals have various health implications.

Table 3: Mineral constituents of dried leaves and roots of *Ageratum conyzoides*

| Minerals | Leaf (mg/100g)           | Root (mg/100g)            |
|----------|--------------------------|---------------------------|
| Ca       | 48.35± 1.22 <sup>a</sup> | 65.10± 1.02 <sup>m</sup>  |
| Na       | 118.54±1.52 <sup>b</sup> | 152.13±1.30 <sup>n</sup>  |
| Mg       | 10.75±0.93 <sup>c</sup>  | 21.67±1.15 <sup>f</sup>   |
| K        | 0.68± 0.02 <sup>d</sup>  | 1.64 ± 0.11 <sup>i</sup>  |
| Zn       | 0.08± 0.002 <sup>e</sup> | 0.15 ± 0.001 <sup>k</sup> |
| P        | 25.25±1.21 <sup>f</sup>  | 48.52± 1.33 <sup>a</sup>  |
| Fe       | 0.04± 0.001 <sup>g</sup> | 0.09± 0.001 <sup>c</sup>  |
| Mn       | 1.02 ± 0.03 <sup>h</sup> | 2.11 ± 0.02 <sup>l</sup>  |
| Cu       | ND                       | ND                        |
| Cr       | ND                       | ND                        |

Values are mean ± SD; n = 3. Values with different superscripts are significantly different (P<0.05). ND = not detected.

Table 4: Vitamin content of dried leaves and roots of *Ageratum conyzoides*

| Vitamin                     | Leaf (mg/100g)             | Root (mg/100g)            |
|-----------------------------|----------------------------|---------------------------|
| Thiamin                     | 0.73 ± 0.10 <sup>a</sup>   | 0.64 ± 0.06 <sup>a</sup>  |
| B <sub>2</sub> (riboflavin) | 28.47 ± 1.21 <sup>b</sup>  | 20.45 ± 0.11 <sup>b</sup> |
| B <sub>6</sub> (Pyridoxine) | 95.56 ± 1.13 <sup>c</sup>  | 81.78 ± 1.10 <sup>c</sup> |
| C (Ascorbate)               | 108.11 ± 0.55 <sup>c</sup> | 94.65 ± 1.13 <sup>c</sup> |
| E (Tocopherol)              | 36.67 ± 0.35 <sup>b</sup>  | 25.64 ± 1.02 <sup>b</sup> |
| Niacin                      | 64.42 ± 1.35 <sup>d</sup>  | 57.55 ± 0.86 <sup>d</sup> |
| A                           | ND                         | ND                        |

Values are mean ± SD; n = 3. Values with different superscripts are significantly different (P<0.05). ND = not detected.

The proximate composition of the leaves and roots, shown in table 2, revealed that the leaf sample contained ash (12.64%), fat (2.27%), fiber (23.50%), moisture (10.02%), protein (14.73%) and carbohydrate (36.84%), while ash (16.95%), fat (2.01%), fiber (26.74%), moisture (13.35%), protein (9.89%) and carbohydrate (31.06%) were found in the roots. The relatively high fiber and low fat contents may be an important contributing factor to some the medicinal applications of the plant. According to Okaka *et al.* [21], clinical studies suggest fiber can exert a wide range of benefits in areas such as bowel function, gut health, immunity, blood glucose control and serum lipid levels.

Na, K, Ca, Mg, P, Zn, Mn and Fe were present in the leaves and roots in proportions comparable to those of some investigated medicinal plants (Table 3) and their concentrations in roots were significantly higher (P<0.05) than in leaves. These minerals play significant roles in several biological processes. Bone growth and turnover are influenced and regulated by the metabolism of Ca, phosphate and Mg; Fe is important in the formation of haemoglobin. Mg and K are also involved in inducement of calmness [22]. The presence of Zn in the extracts may further explain the utilization of the leaves in wound healing. Zn has inhibitory effect on bacterial growth and

is involved in immune response. There is increased demand for Zn during cell proliferation and protein synthesis [23].

Both samples were found to contain vitamins B<sub>2</sub>, B<sub>6</sub>, C, E, thiamin and niacin (table 4). Their values in the leaves did not differ significantly (P> 0.05) from those in roots. The presence of the observed vitamins in the samples is a boost to their therapeutic potentials. Vitamin B<sub>6</sub> (pyridoxine) helps in the breakdown of blood sugar and dilation of blood vessels [24]; vitamin B<sub>2</sub> (riboflavin) is essential for energy production and in its coenzyme forms (FMN and FAD), serves as hydrogen transport systems; vitamin C, an antioxidant, facilitates wound healing, production of collagen, formation of red blood cells and boosts immune system; vitamin E is an antioxidant and plays a role in cellular respiration; niacin (nicotinic acid) is converted to NAD and NADP, which are coenzymes for various oxidoreductases [22].

## CONCLUSION

Alkaloids, flavonoids, tannins, saponins, cardiac glycosides and anthraquinones are contained in the leaves and roots of the plant, while terpenoids occur in the leaves only. The leaves and roots also contain minerals, vitamins and other compounds of known pharmacological activity.

The various medicinal uses of leaves and roots of *Ageratum conyzoides* may be due to these chemical constituents present in them. Studies on the efficacy of extracts of the leaves and roots against some of disease conditions, as claimed by traditional medicine practitioners, are in progress in our laboratory.

## REFERENCES

1. Iwu, M.M., 2000. Food for medicine: Dietary plants and Masticatories as Sources of Biologically active Substances. University of Ife, Nigeria. Ife Press, pp: 303-310.
2. Ogunkunle, A.T.J. and A.L. Tonia, 2006. Plant and their Natural Pharmacological Effects. African Journal of Biotechnology, 5(21): 2020-2023.
3. Devendran, G. and U. Balasubramanian, 2011. Uses of Medicinal Plants. Asian Journal of Plant Science and Research, 2(4): 44-48.
4. Varadarajan, P., G. Rathinaswamy and D. Asirvatham, 2008. Antimicrobial Properties and phytochemical Constituents of Rheo discolor, Ethnobotanical leaflet, 12: 841-845.

5. Okwu, D.E. and O. Ekeke, 2003. Chemical Profile Analysis of Medicinal Plants. International Journal of Pure and Applied Science, 23(9): 235-238.
6. Oladejo, O.W., I.O. Imosemi, F.C. Osuagwu, O.O. Oluwadara, A. Aiku, O. Adewoyin, O.E. Ekpo, O.O. Oyedele and E.E.U. Akang, 2003. Chemical Profile Analysis of Medicinal Plants. African Journal of Biomedical and Pharmaceutical Science, 3(8): 27-31.
7. Okwu, D.E., 2005. Chemical Profile Analysis. International Journal of Molecular Medicine and Advance Science, 25(4): 375-381.
8. Sofowora, A.E., 1986. The state of medicinal plants in Nigeria, University of Ibadan, Nigeria, Press, pp: 66-80.
9. Prince, L. and P. Prbakaram, 2011. Chemical Profile Analysis of Medicinal Plants. Asian Journal of Plant Science, 20(1): 1-8.
10. Kamboj, A. and A.K. Saluja, 2008. *Ageratum conyzoides* L.: A review on its phytochemical and pharmacological profile. Int J. Green Pharm., 2: 59-68.
11. Sofowora, A., 1993. The state of Medicinal plants in Nigeria. 3<sup>rd</sup> edition, University of Ibadan, Nigeria.
12. Trease, G.E. and W.C. Evans, 1983. Textbook of Pharmacognosy. 12<sup>th</sup> ed. Tindall and Co. London, pp: 343-383.
13. Harborne, I.B., 1973. Phytochemical Methods: A guide to modern Techniques of plant analysis 2<sup>nd</sup> ed. Chapman and Hall, New York, pp: 88-185.
14. AOAC, 1990. Official methods of analysis. Association of Official Analytical Chemists, Washington D.C. 15<sup>th</sup>, Ed.
15. Igwe, C., L.A. Nwaogu and C.O. Ujuwundu, 2007. Assessment of hepatic effects, phytochemical and proximate composition of *phyllanthusamarus*. Afr. J. Biotech., 6(6): 728-731.
16. Pearson, D., 1976. The chemical analysis of foods, 17th ed. Churchill Livingstone, London, pp: 3-4.
17. Okwu, D.E. and C.U. Ndu, 2006. Evaluation of the phytonutrients, mineral and vitamin contents of some varieties of yam. Int. J. Mol. Med. And Advance Sci., 12(2): 199-203.
18. Amadi, B.A., M.K.C. Duru and E.N. Agomuo, 2012. Chemical profiles of leaf, stem, root and flower of *Ageratum conyzoides*. Asian Journal of Plant Science and Research, 2(4): 428-432.
19. Teguja, M.N. and K.U. Omak, 1993. Encyclopedia of medicinal plants. J. Biochemical Sciences, 20: 81-82.
20. Okuda, T., Y. Matsuda, A. Yamanaka and S. Sagisaka, 1991. Abrupt increase in the level of hydrogen peroxide in leaves of winter wheat is caused by cold treatment. Plant Physiology, 97: 1265-1267.
21. Okaka, J.C., E.N.T. Akobundu and A.N.C. Okaka, 2006. Food and Human Nutrition OCJ Academic Publishers, Enugu, Nigeria, 135-150, 104-129.
22. Burtis, C.A. and E.R. Ashwood, 2003. Tietz Fundamentals of Clinical Chemistry, 5<sup>th</sup> ed. Elsevier Pub. India.
23. Williams, L. and D. Leaper, 2005. Nutrition and Wound Healing. Clinical Update, 5(1): 3-5.
24. Trumbo, P., 2014. Pantothenic Acid. In: Modern Nutrition in Health and Disease. 11<sup>th</sup> Edition. Eds A.C. Ross, B. Caballero, R.J. Cousins, K.L. Tucker and T.R. Ziegler, Lippincott Williams and Wilkins, Philadelphia, USA, pp: 351-357.