

Phenotypic Characterization of Indigenous Chicken Ecotypes in North Gondar Zone, Ethiopia

¹Addis Getu, ²Kefyalew Alemayehu and ³Zewdu Wuletaw

¹Department of Animal Production, Faculty of Veterinary Medicine, University of Gondar, Ethiopia

²Department of Animal Production and Technology, Bahir Dar University
College of Agriculture and Environmental Sciences, Ethiopia

³Sustainable Land Management Organization (SLM) Bihar Dar, Ethiopia

Abstract: An exploratory field survey was conducted in north Gondar zone, Ethiopia to identify and characterize the local chicken ecotypes. Seven qualitative and twelve quantitative traits from 450 chickens were considered. Chicken ecotypes such as necked neck, Gargie and Gugut from Quara, Alefa and Tache Armacheho districts were identified, respectively. Morphometric measurements indicated that the body weight and body length of necked neck and Gargie ecotypes were significantly ($p < 0.01$) higher than Gugut ecotypes except in shank circumstances. Sex and ecotype were the significant ($p < 0.01$) sources of variation for both body weights and linear body measurements. The relationship of body weight with other body measurements for all ecotypes in both sexes were highly significant ($r = 0.67$, $p < 0.01$). Some traits like wingspan, body length and super length ($r = 0.64$, $P < 0.01$) for males and for females ($r = 0.59$, $P < 0.01$) of necked neck chickens are significantly correlated with body weight. Therefore, highly correlated traits are the basic indicators for estimation of the continuous prediction of body weight of chicken. Identification and characterization of new genetic resources should be employed routinely to validate and investigate the resources in the country.

Key words: Chicken • Ecotypes • Necked neck • Gargie • Gugut • Ethiopia

INTRODUCTION

Ethiopia takes the lead in livestock population and gateways of domestic animals migration from Asia to Africa, thus rolled for widespread distribution and huge population size in the country [1]. Poultry contribute socio-economic roles for food securities, generation of additional and cash incomes [2] and [3]. Therefore, almost all rural and many peri-urban families keep small flock scavenging chickens [4]. In Ethiopia, the population of chickens estimated about 49.3 million of which 97.3, 0.38 and 2.32 % indigenous, hybrid and exotic breeds respectively [5]. Indigenous chickens have good potential to adapt in different agro ecology through habitual management system [6]. Local chickens are non descriptive type and show a large variation which might be attributed to their widespread distribution; [7, 1, 8].

Indigenous chickens are underestimated because of their poor performance. To this effect, they have

been neglected and little attention has been given from researchers, development workers and policy makers to put them in the research and development agendas [9]. Some researchers have done on phenotypic and genetic characterization of indigenous chicken in some parts of Ethiopia. Poultry production and market system was studied in southern Ethiopia by Mekonnen; [10] characterization of poultry productivity and market system by Bogale, [11] and genetic parameters on horror chickens for weights and egg production trait was conducted by Dana *et al.* [12]. However, comprehensive genetic resources identification in the remote districts northern Gondar zone in general, identification and characterization of new local chicken ecotypes in particular were not studied in the area.

Therefore, the objective of this study was to identify and characterize the new local chicken ecotypes in North Gondar zone, Ethiopia.

MATERIALS AND METHODS

Description of the Study Area: The study was conducted in three districts of north Gondar zone (Quara, Alefa and Tache Armacheho) of Ethiopia. The altitude of the zone is ranged from 528-4620 meter above sea level (m.a.s.l) and rainfall of 880 - 1772 mm with the temperature of 44.5°C to -10°C. Quara district is located western part of north Gondar Zone between 11°47' and 12°21' and latitude and 35°16' and 35°47'E longitude. It is 1123 km far from Addis Ababa and 324 km from Gondar town and elevation ranging 528 - 654 meter above sea level. The annual temperature ranges 25-44°C with mean annual rainfall of 600 - 1000 mm [5]. The same source indicated that Alefa district is located at 162km in southwest of Gondar town and 909 km from Addis Ababa with the temperature of 25 - 30°C and annual rainfall of 900-1400mm. Armacheho district is also found 814 km northwest of Addis Ababa and 65km North West of Gondar town with the altitude of 600-2000 m.a.s.l with the temperature of 25 - 42°C and with annual rainfall of 800-1800 mm [5].

Data Collection Methods: In addition to exploratory field survey, semi-structured questionnaires and participatory rural appraisal (PRA), focus group discussion, field observation, trait characterization and body measurements were employed to dig up the required information. For the morphological and biometrical measurements, all matured chicken ecotypes n = 450, 150 males and 300 females were measured. Qualitative traits such as plumage size, body shape, comb type, shank colour, skin color, head shape and eye colour was documented through direct visualization. Whereas measurable trait like (body weight (kg), Body length, wing span, shank length and circumference, wattle length and width, keel length, super length, beak length, comb length and width) were measured using spring balance and centimeter (cm) in the nearest two digits [13].

Data Management and Statistical Technique: Information from the focused group discussion and personal observation were briefly summarized and synthesized. More over quantitative data was used by General Linear Model (GLM) and imported to [14] version 9. Tukey comparison test was used to compare sub factor brought significant difference. The model was used for body weight and linear body measurement of chickens' ecotypes by considering the fixed effects of sex and ecotype.

$$\text{Model 1. } Y_{ijk} = \mu + A_i + D_j + AD_{ij} + e_{ijk}$$

where:

Y_{ijk} = The observed body weight and linear body measurement of chickens

μ = overall mean

A_i = Fixed effect of i^{th} eco type (I=1, 2 and 3)

D_j = the effect of k^{th} sex (j= male and, I= female)

AD_{ij} = The fixed effect interaction i^{th} of eco type with j^{th} of sex

E_{ijk} = Random residual error

The second used model was multiple linear regression coefficients for continuous predicting body weight of matured cocks and hens using eleven linear body measurements (independent variables) in each ecotype.

$$\text{Model: 2. } Y_j = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \beta_4 X_4 + \beta_5 X_5 + \beta_6 X_6 + \beta_7 X_7 + \beta_8 X_8 + \beta_9 X_9 + \beta_{10} X_{10} + \beta_{11} X_{11} + e_{ijk}$$

where:

Y_j = dependant variable or predicted mean body weight of chickens; β_0 = the intercept; $X_1, X_2, X_3, X_4, X_5, X_6, X_7, X_8, X_9, X_{10}$ and X_{11} are the independent variables for wing span, body length, shank length, shank circumference, keel length, super length, beak length, wattle length, wattle width, comb length and comb width, respectively.

$\beta_1, \beta_2, \beta_3, \beta_4, \beta_5, \beta_6, \beta_7, \beta_8, \beta_9, \beta_{10}$ and β_{11} are the regression coefficient of the variable $X_1,$

$X_2, X_3, X_4, X_5, X_6, X_7, X_8, X_9, X_{10}$ and X_{11} and

e_{ijk} = the residual error

RESULTS

Necked Neck Chickens: Necked neck chickens are found in a very hot ecological zone of Quara district which is located at altitudes ranging from 528-654 meter above sea level with the maximum temperatures of 44°C. They are maintained under scavenging system with small supplementation and sheltered outside the family house (perch). The chickens have predominantly white 28 % and red 20 % body plumage colors. However, they had heterogeneity and diverse additional plumage color like; red-braunish (0.7%), white with red tips (5.3%), black with white tips (10.7%), black (7.3%), multicolor (5.3%) and white black red trips (13.3%). About 53 % of the birds have white skin colour, 66 % single combed followed by

Table 1: Morphological Characteristics of Newly Founded Indigenous Chicken Ecotypes

Character	Attributes	Ecotypes by proportions and their associations					Cramer's V	P-value
		Necked neck=150	Gasgie N=150	Gugut N=150	Overall N=450			
Pc	White black and red ^{ns} trips(<i>Kiy Tikur Teterma</i>)	(0.67)	(1.34)	(2.67)	1.56	0.13	0.323	
	Black with white tips (<i>Tikur Teterma</i>) ^{ns}	(10.70)	(8.67)	(6.00)	8.44	0.13	0.286	
	White with red tips (<i>Kiy Teterma</i>) ^{ns}	(5.33)	(8.67)	(8.00)	7.33	0.11	0.465	
	Grayish-mixture (<i>Gebsema</i>) ^{***}	(10.70) ^b	(22.00) ^a	(10.00) ^b	14.20	0.33	0.005	
	Red-braunish(<i>Kokima</i>) ^{**}	(13.30) ^a	(8.67) ^a	(2.00) ^b	8.00	0.34	0.003	
	Multicolor (<i>Ambesa</i>) [*]	(5.33) ^{ab}	(4.00) ^b	(10.00) ^a	6.44	0.22	0.037	
	Black (<i>Tikur</i>) ^{***}	(7.33) ^b	(5.33) ^b	(22.00) ^a	11.50	0.52	0.001	
	White (<i>Nech</i>) ^{***}	(28.00) ^a	(9.33) ^b	(9.33) ^b	15.60	0.66	0.001	
	Red (<i>Kiy</i>) ^{***}	(18.70) ^b	(32.00) ^a	(30.00) ^a	26.90	0.60	0.001	
	Plain (<i>Ebaberis</i>) ^{ns}	(70.00) ^a	(65.33) ^a	(76.00) ^a	70.40	0.10	0.19	
Hs	Crest (<i>Gutya</i>) ^{ns}	(30.00)	(34.67)	(24.00)	29.60	0.08	0.23	
	Ct	Doublex (V-shape) [*]	NA	(9.33) ^b	(24.00) ^a	32.70	0.21	0.024
Single ^{***}		(66.00) ^a	(27.33) ^b	(30.00) ^b	38.90	0.42	0.001	
Rose ^{***}		(34.00) ^b	(60.70) ^a	(38.70) ^b	44.40	0.33	0.001	
Pea ^{ns}		NA	(2.67) ^a	(7.33) ^a	3.33	0.03	0.045	
Skc	Yellow ^{***}	(66.00) ^a	(24.70) ^b	(67.30) ^a	53.10	0.55	0.001	
	White ^{***}	(23.30) ^b	(66.00) ^a	(20.70) ^b	42.90	0.75	0.001	
	Black [*]	(4.00) ^{ab}	(0.67) ^a	(8.00) ^b	4.22	0.20	0.001	
	Green [*]	(6.00) ^a	NA	(1.33) ^b	1.56	0.18	0.004	
	Red [*]	(0.67) ^a	(8.67) ^b	(3.33) ^a	4.22	0.09	0.05	
Ec	Orange ^{ns}	(0.67) ^a	(4.67) ^a	(4.67) ^a	3.33	0.10	0.088	
	Black ^{ns}	(6.00) ^a	NA	(3.33) ^a	3.11	0.16	0.058	
	Pur ^{***}	(1.33) ^b	NA	(3.33) ^a	1.56	0.13	0.021	
	Red ^{***}	(92) ^b	(95.30) ^a	(88.70) ^b	92	0.42	0.001	
BS	Triangular ^{***}	(40.7) ^a	(15.30) ^b	(30.70) ^a	29.00	0.32	0.001	
	Blocky ^{***}	(58.00) ^a	(31.30) ^b	(49.30) ^a	46.20	0.36	0.001	
	wedge ^{***}	(1.30) ^b	(5.33) ^b	(20.00) ^a	24.90	0.32	0.001	
Snc	Yellow ^{***}	(68.00) ^b	(66.00) ^{ab}	(51.30) ^a	62.00	0.19	0.002	
	White ^{***}	(16.70) ^{ab}	(14.00) ^b	(26.00) ^a	19.10	0.20	0.001	
	Black ^{ns}	(7.33) ^a	(9.30) ^a	(8.00) ^a	8.00	0.07	0.285	
	Green ^{ns}	(7.33)	(10.00)	(8.00)	8.22	0.03	0.814	
	Red [*]	(0.67) ^b	(0.67) ^b	(5.33) ^a	2.67	0.17	0.012	

NA = not available, a, b, c, with different superscript within a row are significantly different (P < 0.05)



Fig. 1: Typical necked neck male (right) and female (left) chicken ecotypes



Fig. 2: Typical Gasgie male (left) and female (right) chicken type



Fig. 3: Typical Gugute male and female chicken type

34 % rose combs and 70 % plain head are facial appearance (Table 1 and Figure 1) of the ecotypes. The other peculiar features were aggressive, high feed requirements, good productive and reproductive performance and tolerance to common diseases and had large carcass (cleaned meat) body weight.

Gasgie Chicken Ecotypes: Their dominant geographical distribution and origin is limited to the mid and plain topography of Alefa district at average altitude of 1400 mere above sea level (Fig. 1). Most of the households keeping these chickens sheltered in the family house during the night, while they spend the day scavenging in the backyards supplemented with grains and food leftovers. The chickens have predominantly red 32 % body plumage colour with other diverse plumage colour such as white (9.3%) red-braunish (9%), white with red tips (9%), black with white tips (9%), black (5.3%), multicolor (4%) and white black red trips (1.3%). About 60.7 % of the ecotypes are rose combed. Long necked (especially males), short weaning time, docile and good productive and reproductive performances are the main unique feature of the ecotype.

Gugut Chickens Ecotypes: Gugut chickens are well dominated in Debresina area of Tache Armacheho district and inherited from early lived parents and transmitted from generation to generation. Most of the households keeping these chickens provided separate shelters for housing during the night in case of suffocation and predators (snack) like that of Quara district. Dense feathers from the neck, absent of wattle in female, low performance, short beak length, ability to resist endemic disease, small body size, passive and easily exposed to predators are the unique behaviors of the ecotype. The chickens have predominantly white (22%) body plumage colour with other diverse plumage colour such as multicolor (10%), red (9.3%), black (9.3%), white with red tips (8%), red-braunish (2%), black with white tips (2%) and white black red trips (2%). In later case they also showed different other qualitative and quantitative variations described from.

The results indicated that the predominant average plumage color of newly identified local chicken ecotypes were 26.90% red followed by 15.60% white and 14.20% Grayish mixture. About 44.4 % are rose comb type, 42.90% white sicken color and 46.20% blocky body shape and

Table 2: The Interaction Effect of Sex and Ecotypes on Linear Body Measurements (Cm) and Body Weight (Kg)

Parameters	Sex	Necked	Gasgie	Gugute			Overall mean	Grand Mean
Sample size	M	50	50	50			150	450
	F	100	100	100			300	
Effects & levels		LSM±SE	LSM±SE	LSM±SE	CV %	P-value	LSM±SE	LSM±SE
WS	M	38.70±2.6 ^a	39.61±0.42 ^a	35.97±0.23 ^b	6.51	.0001**	38.09±0.24 ^a	37.04±.13
	F	37.17±2.36 ^a	37.36±0.26 ^a	35.03±0.18 ^b	6.19	.0001**	36.52±.14 ^b	
SL	M	9.61±1.03 ^a	7.25±0.10 ^b	7.37±0.73 ^b	10.05	.0001**	8.08± 0.11 ^a	7.79±0.15
	F	9.043±1.10 ^a	6.80±0.06 ^c	7.08±0.05 ^b	9.10	.0001**	7.64 ±0.07 ^b	
BL	M	38.12±2.14 ^a	36.10±0.34 ^a	35.2±0.09 ^b	9.49	.0002**	36.77 ± 03 ^a	35.79±0.09
	F	36.90±2.61 ^a	34.60±0.26 ^b	34.37±0.21 ^b	6.93	.0001**	35.29±0.16 ^b	
CL	M	3.25±0.87 ^a	3.16±0.12 ^a	3.08±0.09 ^a	26.24	0.594 ^{ns}	3.16 ± 0.07 ^a	2.76±0.09
	F	2.99±3.68 ^a	2.28±0.07 ^b	2.40±0.06 ^{ab}	35.61	0.0482 [*]	2.55 ±0.13 ^b	
CW	M	2.11±0.82 ^a	1.93±0.13 ^a	2.19±0.05 ^a	38.36	0.255 ^{ns}	2.08± 0.07 ^a	1.68±0.04
	F	1.78±0.85 ^a	1.07±0.06 ^b	1.59±0.06 ^a	45.55	.0001**	1.48 ±0.04 ^b	
WL	M	2.76±0.69 ^a	2.70±0.14 ^a	1.83±0.23 ^b	32.19	.0001**	2.43± 0.07 ^a	1.76 ±0.06
	F	2.44±0.80 ^a	1.84±0.04 ^b	NA	37.23	.0001**	1.42 ±0.07 ^b	
WW	M	2.76±1.01 ^a	2.32±0.16 ^b	1.45±0.09 ^c	44.53	.0001**	2.17± 0.09 ^a	1.51±0.06
	F	2.34±1.03 ^a	1.19±0.05 ^b	NA	56.29	.0001**	1.18 ±0.07 ^b	
bl	M	2.42±0.45 ^a	2.00±0.02 ^b	1.85±0.10 ^c	14.12	.0001**	2.09 ± 0.03 ^a	2.03 ±0.02
	F	2.28±0.60 ^a	1.93±0.0 ^b	1.78±0.02 ^c	18.67	.0001**	1.99 ±0.02 ^b	
sl	M	0.66±0.8 ^a	0.49±0.10 ^a	0.17±0.09 ^b	46.35	.0009**	0.44 ± 0.05 ^a	0.18 ±0.02
	F	0.09±0.32 ^a	0.08±0.02 ^a	NA	48.83	0.0172 [*]	. 011± 0.18 ^a	
SC	M	3.58±0.50 ^b	3.25±0.07 ^b	3.85±0.03 ^a	20.78	.0001**	4.81± 0.18 ^a	3.78±.07
	F	3.31±0.59 ^a	3.11±0.03 ^b	3.38± 0.07 ^a	17.23	.0027**	3.27±.03 ^b	
KL	M	9.11±1.02 ^a	9.55±0.15 ^a	7.62±0.23 ^b	16.81	.0001**	7.51±0.24 ^b	8.24 ±0.09
	F	8.56±0.87 ^b	9.27±0.08 ^a	7.98±0.07 ^c	9.08	.0001**	8.60 ±0.05 ^a	
Wt	M	1.78±0.31 ^a	1.71±0.05 ^a	1.40±0.04 ^b	18.15	.0001**	1.63 ±0.03 ^a	1.46 ±0.01
	F	1.52±0.26 ^a	1.36±0.03 ^b	1.23±0.02 ^c	17.50	.0001**	1.37 ±0.02 ^b	

Ws= wingspan, SL= shank length, BL= body length CL= comb length, CW=comb, width, WW=wattle, width, WL= wattle, length, KL= keel length, sl= sure length, bl= beak length, SC= shank circumference, in the measurement of cm Wt = weight (kg) and NA = not available LSM = least square mean and SE= standard error, a, b, c means different superscripts are significantly different (P<0.05)

70.40% plain headed are the most dominant observable traits of Gugut chickens ecotypes (Table 2). About 34, 60.7 and 38.7 % of chicken from necked neck, Gasga and Gugut ecotype were characterized by rose comp type respectively. The proportion of plain head shape in chicken populations of necked neck, Gasgie and Gugut were comparable from 70%, 65.3% and 76 %, respectively. This variation could be feed availability reflecting, adaptation fitness to their environment [15].

Quantitative Traits of Chickens: A total of 450 adult hens and cocks with twelve measurable parameters such as Wingspan (WS), shank length (SL), shank circumference (SC), Body length(BL), comp length (CL), comp width (CW), wattle length (WL), wattle width (WW), beak length (bl), super length (sl), keel length (KL) (cm) and body weight (Wt) (kg) for different sexes were considered. The least squares mean of body weight and body measurements of necked neck, Gasgie and Gugut chickens with HSD (Honestead significant different) comparison tests are presented in (Table 1). The overall least squares mean of Wingspan, shank length, shank

circumference, Body length, comp length, comp width, wattle length, wattle width, beak length, super length, keel length and body weight were (37.04±0.13), (7.79±0.15), (3.78±0.07), (35.79±0.09), (2.76±0.09), (1.68±0.04), (1.76 ±0.06), (1.51±0.06), (2.03 ±0.02), (0.18 ±0.02), (8.24 ±0.09) (cm) and (1.46 ±0.01) (kg), respectively.

Whereas, overall mean square for body weight obtained for male and female was (1.63±0.03) and 1.37±0.02 (kg), respectively. Despite body weight of male in necked neck 1.78±0.31 (kg) was significantly ($P < 0.01$) heavier than Gugut 1.40±0.04 (kg) but from Gasgie 1.71±0.05 (kg) cocks. Furtherly, the necked neck cocks and hens were found to significantly taller shank length of 9.61±1.03 and 9.043±1.10 (cm) respectively. However, shank circumference of Gugut cocks and hens inversely had superior shank circumferences of 3.85±0.03 and 3.38±0.07 (cm) than 3.58±0.50 and 3.1±0.59 (cm) from necked neck and 3.25±0.07 and 3.11±0.03 (cm) from Gasgie for male and female chickens, respectively. Comp lengths were not significantly different from necked neck chicken ecotypes for males 3.25±0.87 and for female 2.99±3.68 (cm). As to beak length, the necked neck and Gasgie cocks had

Table 3: Correlation Coefficients on Body Weight and Linear body Measurements.

Traits	Variables	Sex and ecotype					
		Necked neck		Gasgie		Gugute type	
		M	F	M	F	M	F
WS	N	50	100	50	100	50	100
	r	0.64**	0.56**	0.35*	0.41**	0.50**	0.39**
SL	N	50	100	50	100	50	100
	r	0.18 ^{ns}	0.20*	0.26*	0.29**	0.54**	0.08 ^{ns}
BL	N	50	100	50	100	50	100
	r	0.59**	0.54**	0.49**	0.59**	0.33*	0.50**
CL	N	50	100	50	100	50	100
	r	0.31*	0.05 ^{ns}	0.54**	0.41**	0.40**	0.35**
CW	N	50	100	50	100	50	100
	r	0.15 ^{ns}	-0.07 ^{ns}	0.64**	0.39**	0.37**	0.21 ^{ns}
WL	N	50	100	50	100	50	100
	r	0.05 ^{ns}	-0.01 ^{ns}	0.67**	0.39**	0.39**	NA
WW	N	50	100	50	100	50	100
	r	0.05 ^{ns}	-0.13 ^{ns}	0.52**	0.47**	0.49**	NA
bl	N	50	100	50	100	50	100
	r	-0.22 ^{ns}	-0.01 ^{ns}	0.20 ^{ns}	0.30**	0.24*	0.22*
sl	N	50	100	50	100	50	100
	r	0.48**	0.27**	0.52**	0.28**	0.21 ^{ns}	0
SC	N	50	100	50	100	50	100
	r	0.31*	0.13 ^{ns}	0.35**	0.18 ^{ns}	0.04 ^{ns}	-0.02 ^{ns}
KL	N	50	100	50	100	50	100
	r	0.37**	0.28**	0.62**	0.33**	0.23*	0.21*

N= number of sample size, r = correlation coefficients

Table 4: Prediction Equations in Multiple Regression Analysis of Body Weight

Ecotype	Male	R ²	Female	R ²
Necked neck	Y= -1.34+0.08WS	0.40	Y= -0.78 +0.06WS	0.31
	Y= -1.12+0.07WS+0.17sl	0.60	Y= -1.48 +0.04WS+ 0.04BL	0.41
	Y= -2.12 + 0.06 WS+0.04 BL+0.14 sl	0.65	Y= -1.60+0.05WS+0.04BL+0.10bl	0.47
Gasgie	Y= Y=1.11+0.22WS	0.45	Y= -0.89+0.06BL	0.35
	Y=0.49+0.21 WS +0.2SC	0.53	Y= -0.77+0.05BL+0.19WW	0.46
Gugut	Y=1.1+0.21WW	0.30	Y= -0.04+0.04BL	0.25
	Y=-0.97+0.16WW+0.06WS	0.44	Y= -0.47+0.04BL+0.22bl	0.31
	Y=-1.05+0.05WS+0.09CL+0.13WW	0.50	Y= -0.47+0.03BL+0.07CL+0.24bl	0.37
	Y=-0.22+0.05WS+0.12CL+0.14WW+0.03KL	0.55	Y= -0.8+0.03BL+0.06CL+0.23bl+0.04KL	0.40
			Y=-1.15+0.03BL+0.06CL+0.2bl+0.05SC+0.07KL	0.42

the longest beak length of, 2.42±0.45 and 2.00±0.02 (cm), respectively; while the smallest beak length was recorded for Gugut cocks 1.85±0.10 (cm). Both the male and female chickens had a non significant ($p > 0.05$) variation beak length with in ecotypes (Table 2).

Correlations of Body Weight and Other Linear Body Measurements: Live weight was positively correlated ($r = 55.5$, $P < 0.01$) with wingspan. Body length and super length in necked neck were positively correlated, males ($r = 0.62$, $P < 0.01$) and females ($r = 0.55$, $P < 0.01$). Whereas

WL is highest correlated trait ($r = 0.67$, $P < 0.01$) with body weight of Gasgie male chickens. The high correlation coefficients between body weight and other body measurements ($P < 0.01$) helped to predicting body weight of chickens.

Prediction Equation Models: A Stepwise multiple regression analysis was first carried out linear regression and conducted to multiple regressions by adding WS, BL and WW at a time in three chicken ecotypes. The regression result of necked neck cocks and females and,

Gasgie cocks were found to the value of 0.40, 0.31 and 0.45 with body weight, respectively. The essence was to determine effects of other body measurements on body weight prediction and applied additional variables to improve the regression coefficient as discussed in (Table 4). To increase meat and egg production requires genetic improvement of body weight of chickens. But, proper measurement of this variable is often hard in villages due to lack of weighing scales. Hence, easily measurable linear body measurements are more relevant for chickens' body weight prediction at farmers' level rather subjectively judging by hand. In addition the present farmers are active at early morning by providing supplementary feed to their chicken before brought to the market.

DISCUSSIONS

More than 70% of the population of chicken ecotypes in the study area carrying the naked neck gene that we studied was new and significantly ($p < 0.001$) higher than those reported in other parts of Ethiopia (7.9%) [16] <2%; [17], Nigeria (6%) [18] and Botswana (3.6%) [19]. The chicken ecotypes in this study have plumage color differed from report of northwest Ethiopian local chicken ecotypes [1]. Furthermore, the overall variations of dominant rose comb types, white skin colored, blocky body shaped and plain head types were the most dominant observable traits of chickens different from the studies done at Bure and Fogera districts in the Amhara region and Dale district in Southern Ethiopia [20]. This variation could be a breed-specific trait, nutritional status of the breeds, genotype and reflecting adaptation fitness to their environment [16, 15]. In this study, complete absence of wattle gene from Gugut female and long necked morphology and early weaning of Gasgie chicken ecotypes makes unique from reports of previous studies in Ethiopia and elsewhere in the tropics reported by [1, 15, 16].

The overall list mean square for body weight obtained for male and for female which were varied from Ethiopian chickens reported by (Danna) [15] which were 1.6 (kg) for males and 1.27(kg) for females [21, 22]. Despite the list mean square for body weight obtained in necked neck and Gasgie chickens were heavier than chicken in central [15] and Northwest Ethiopia [1] of 1.26 and 0.87 kg for adult male and female, respectively [23]. The frequency of chicken ecotypes carrying the naked neck gene that we studied was significantly ($p < 0.001$) higher than those reported in other parts of Ethiopia (<2%) [15] Nigeria (6%) [18] and Botswana (3.6%) [19].

CONCLUSIONS

A key informant was found to be a useful individual to identify distinct animal genetic resource. Necked neck, *Gasgie* and *Gugut* chicken ecotypes were newly identified and dominantly found in *Quara*, *Alefa* and *Tache Armacheho* district northern Ethiopia, respectively. Furthermore phenomic characterization was helped to describing the general uniqueness of the populations. The highest adult body weight was obtained from Naked-neck, followed by Gasgie chicken ecotypes. Necked neck, *Gasgie* and *Gugut* chicken ecotypes were dominantly found in *Quara*, *Alefa* and *Tache Armacheho* district, respectively. In addition qualitatively as Necked neck chickens ecotype was easily identified by their complete absence of feather at neck and chest. Whereas, *Gasgie* chicken ecotype was also characterized by their normal feather (not bold or muffed) and long necked. Generally, complete absence of wattle in hens, smallest of all and dunce feather at neck (muffed) in both sexes were the most famous characteristics of *Gugut* chicken ecotype. Similarly, Naked-neck chickens had the longest shank than those of other current findings. The population of the newly found chicken ecotypes showed heterogeneity in most morphological traits considered. Thus, In-depth molecular characterization using genetic markers should be undertaken to confirm the level of genetic variations and relationships among newly identified and other indigenous chicken ecotypes.

ACKNOWLEDGEMENTS

This piece of work is funded by the German International (GIZ) project to which we are grateful. I thank also all interviewed village chicken owners, district agricultural office experts, key informants and other participants who helped us to carry out every activity on the successful achievement of this research.

REFERENCES

1. Halima, H., 2007. Phenotypic and genetic characterization of indigenous chicken populations in Northwest Ethiopia. PhD. Thesis submitted to the faculty of National and agricultural sciences department of animal Wild life and Grass land Sciences University of the Free State, Bloemfontein and South Africa.

2. Kondombo, S.R., 2005. Improvement of village chicken production in a mixed farming system in Burkina Faso. Ph.D Thesis. Wageningen Institute of Animal Sciences, Animal Nutrition Group, Wageningen University, the Netherlands.
3. Salam, K., 2005. Improvement of village chicken production in a mixed (chicken ram) farming system in Burkina Faso. Ph.D.Thesis. Wageningen Institute of Animal Sciences, Animal Nutrition Group, Wageningen University, the Netherlands.
4. Jens, C.R., P. Anders, V. Charlotte, M.C. Ainsh and F. Lone, 2004. Keeping of village poultry. A technical manual for small-scale poultry production. Copenhagen, Denmark, pp: 34.
5. CSA, 2011. Agricultural sample survey 2010/11. volume. 2: statistical bulletin 05.Report on livestock and livestock characteristics (prevent peasant moldings), Addis Ababa, February 2011, 21.
6. Taddele, D. and Y. Alemu, 1997. Studies on village poultry production systems in the central highlands of Ethiopia. MSc, thesis submitted to Swedish University of Agricultural Sciences.
7. Taddele, D., Y. Alemu and K. Peters, 2003. Village chicken production systems in Ethiopia: Use patterns and performance evaluation and chicken products and socio- economic functions of chicken. *Livestock Research for Rural, Development*, (15)1.
8. Taddele Desse, 2003. Phenotypic and genetic characterization of local chicken ecotypes in Ethiopia. Ph.D. Thesis submitted to Humboldt University of Berlin, Germany.
9. Mekonnen, G., 2007. Characterization of smallholder poultry production and marketing system of Dale, Wonsho and Loka Abaya Weredas of southern Ethiopia. MSc Thesis, Awassa College of Agriculture, Hawassa University.
10. Bogale, K., 2008. In situ characterization of local chicken eco-type for functional traits and production system in Fogera district, Amhara regional state. M.Sc. Thesis Submitted to the department of Animal Science School of graduate Studies, Haramaya University.
11. Dana, N., W.E. vander and A.M. Johan, 2010. Genetic and phenotypic parameter estimates for body weights and egg production in Horro chicken of Ethiopia. Submitted to *Trop. Anim. Health Prod, Animal breeding and genomics Centre*, 14 July 2010, Wageningen University Netherlands.
12. Statistical Analysis System (SAS), 2002. SAS Users? Guide, Version 9.1, (SAS Institute Inc, North Carolina).
13. FAO., 2011. Draft guidelines on phenotypic characterization of Animal genetic Resource. on Genetic Resources for Food and Agriculture Rome, 18-22 July, 2011, pp: 6.
14. Dana, N., 2011. Breeding programs for indigenous chicken in Ethiopia analysis of diversity in production systems and chicken populations; Ph.D. Thesis submitted in fulfillment of the requirements for the degree of doctor at Wageningen University Netherlands.
15. Aberra, M. and N. Tegene, 2011. Phenotypic and morphological characterization of indigenous chicken populations in southern region of Ethiopia. *Animal Genetic Resources*, 49: 19-31.
16. Badubi, S.S., M. Rakereng and M. Marumo, 2006. Morphological characteristics and feed resources available for indigenous chickens in Botswana. CIPAV, Columbia, Livestock Research for Rural Development.
17. Gueye, E.F., 1998. Village egg and fowl meat production in Africa. *World's Poultry Science Journal*, 54: 73-86.
18. Msoffe, P.M., U.M. Minga, J.E. Olsen, M.G.S. Yongolo, H.R. Juul-Madsen, P.S. Gwaksia and M.M.A. Mtambo, 2001. Phenotypes including, immune competence in scavenging local chicken ecotypes in Tanzania. *Trop. Anim. Hlth Prod.*, 33(4): 341-354.
19. Fisseha, M., M. Abera and D. Tadelle, 2010. Assessment of village chicken production system and evaluation of the productive and reproductive performance local chicken ecotype in Bure district North West Ethiopia. *African Journal of Agricultural Research*, 5(13): 739-1748.
20. Dana, N., D. Tadelle, H. V. Elisabeth and A. M. Johan, 2009. Morphological features of indigenous chicken populations of Ethiopia. Animal Breeding and Genomics Center, Wageningen University. *Animal genetic Resources*, 46: 11-23.
21. Panda, B.B., Kalpesh Gaur, M.L. Kori, L.K. Tyagi, R.K. Nema, C.S. Sharma and A.K. Jain, 2009. Anti-Inflammatory and Analgesic Activity of *Jatropha gossypifolia* in Experimental Animal Models, *Global Journal of Pharmacology*, 3(1): 01-05.
22. Parmar Namita, Rawat Mukesh and J. Kumar, 2012. Vijay, *Camellia Sinensis* (Green Tea): A Review *Global Journal of Pharmacology*, 6(2): 52-59.
23. Jagadeeswaran, M., N. Gopal, B. Jayakar and T. Sivakumar, 2012. Simultaneous Determination of Lafutadine and Domperidone in Capsule by High Performance Liquid Chromatography, *Global Journal of Pharmacology*, 6(2): 60-64.