

Effect of Pranidipine on Primphos-Induced Seizure in Mice

Hatefi Farzam

Young Researchers Club, Tabriz Branch, Islamic Azad University, Tabriz, Iran

Abstract: Primphos a synthetic Organophosphorus compound used mainly as insecticides are the property of insecticide. These toxins can be used as insecticides in agriculture and medicine for animals and the destruction of ectoparasites. Studies have shown that Primphos creation seizure effects in different animals. Pranidipine, dihydropyridine calcium channel blockers, widely used for treatment of cardiovascular diseases. Studies have shown that the calcium channel blockers having anticonvulsant effects in different animal models. The aim of this study was to determine the effect of pranidipine on Primphos-induced seizures in mice. In this experiment, the animals were received different doses of pranidipine (2.5, 5, 10, 20 and 40 mg kg b. wt.) intraperitoneally, 30 min before intraperitoneal injection of Primphos (500 mg/kg b. wt.). After Primphos injection, clonic and tonic seizures and death was investigated. Results showed that pranidipine dose dependently reduced the severity of Primphos-induced seizures, so that pranidipine at a dose level of 5 mg/kg and 40 mg/kg, had the lowest ($p < 0.05$) and highest ($p < 0.001$) anticonvulsant effects, respectively. The anticonvulsant activity of pranidipine suggests that possibly due to antagonistic effect on voltage-dependent calcium channel.

Key words: Pranidipine • Primphos • Seizures • Mice

INTRODUCTION

Epilepsy is one of the major neurological diseases in humans and about one percent of the population is involved. It has been shown that epileptic seizure occurs due to occasional discharges in nerve tissue. It is recognized that occasional changes in reversible neuronal function, causing brain electrical activity. In some cases, the seizure occurs due to the entry of calcium ions into nerve cells and reducing intracellular calcium concentration in some epileptic animal models which has inhibitory effect on seizures. During seizures increased intracellular calcium ion concentration while extra cellular calcium concentration decreases [1-6]. Calcium channel antagonists for the treatment of hypertension were produced in the year 1980. Use of these drugs over time to treat other diseases was developed, such as treatment of angina, supraventricular tachycardia attack, hypertrophic cardiomyopathy, pulmonary hypertension and migraine. Recently have shown that calcium channel blockers may have anticonvulsant effects in some animal models. Calcium channel blockers inhibit calcium ion flow through L-type calcium channels sensitive to voltage [1, 3, 7]. It has been shown that calcium channel inhibitors in models of nerve tissue in a large protective effect [3, 8].

Other investigators also reported that calcium channel in some models have anticonvulsant effects [1, 3, 7, 9, 10] but in all animal models of seizures has not demonstrated these effects [10, 11]. Also in rats anticonvulsant effects of calcium channel inhibitors shown but seizure agent has not Primphos. Some medications such as anticonvulsants phenytoin and carbamazepine can induce their effects by inhibiting sodium channels directly and indirectly by preventing the flow of calcium from the membranes of neurons and reduction of excessive concentration of intracellular calcium. Specific drugs used to treat epilepsy are absence seizure kind of like channels as T-type calcium in thalamic neurons are blocked. This reduced calcium ion concentration of the important goals in development of neuroprotective and antiepileptic drugs [5, 6]. Calcium entry into neurons play an important role in creating the seizures and calcium channel inhibitors have different effects on health, including cardiovascular diseases, migraine and headaches caused by vascular changes, nerve regeneration and neuronal regenerative processes [4]. So it seems calcium channel inhibitors used to treat seizures can be useful. Results of these studies for the anticonvulsant effects of calcium channel inhibitors suggests, therefore likely to pranidipine reduce Primphos-induced seizures. Since no research based on the

combined effect of these seizures from Primphos there, in this case study seems necessary. Insecticide use in agriculture and veterinary medicine as strange since World War II and grew during the past 20 years has reached its highest rate constant. While the main consumers of agricultural insecticide industry is also large quantities of other industries use them and their applications in and around homes is considerable. Most of insecticide residues on the remaining products and people exposed to low doses of chemicals through the foods. Numerous incidents of acute insecticide poisoning caused by eating food that mainly followed during storage or transportation had been infected was created [1, 2, 13]. Including the insecticide, which are potential toxicity, are organophosphate. One of the organophosphate is Primphos. The aim of this study was to determine the effect of pranidipine (calcium channel antagonist) on Primphos-induced seizures.


MATERIALS AND METHODS

Male mice NMRI, weighing between 25-30 grams, were purchased and maintained in laboratory of animal breeding center of Tabriz Islamic Azad University and were kept at room temperature, light and humidity constant. Animals' access to food and water were freely. All tests were performed between 10-16 hours. Primphos and pranidipine, both were solved in Twin 80 (5%). Animals were divided randomly and placed in treatment groups (each group n=10). Pranidipine and Twin 80 were


administered intraperitoneally with constant volume and by weight per animal. To remove the effect of injection volume on seizures, all drugs and Twin 80, at 10 ml/kg was set. First, seizures was assessed in animals receiving Primphos and then evaluated the effect of Twin 80 on Primphos-induced seizures with the above injection, 30 minutes before the seizure was determined. More experimental animals, different doses of pranidipine (2.5, 5, 10, 20 and 40 mg/kg) received 30 min before the intraperitoneal injection of Primphos. To create seizures, mice received Primphos (500 mg/kg B.W.) intraperitoneally and then the animals treated for 120 minutes were recorded by video camera. Films from the following four behaviors were recorded: starting time of clonic seizures after injection of Primphos (second), Generation time of death after Primphos injection (second), mortality after injection of Primphos (percentage) and type of seizures induced by injection of Primphos (percentage). After testing data as the mean±SEM expression and to analyze data, ANOVA followed by Tukey multiple comparison tests were used. Value of $p < 0.05$ to determine significance between groups was considered.

RESULTS


Effect of Twin 80 as solvent, on Primphos-induced seizures showed that this substance has no significant effect on seizures. Therefore the results had not presented in graphs and tables have been avoided. Effect of different doses of pranidipine (2.5, 5, 10, 20 and 40 mg/kg



Graph 1: Effect of different doses of pranidipine on the starting time of clonic seizures after injection of Primphos (second). mean±SEM any form of graphs are presented. * $P < 0.05$, ** $P < 0.01$ and *** $P < 0.001$ compared with Primphos group



Graph. 2: Effect of different doses of pranidipine on Generation time of death after Primphos injection (second). mean±SEM any form of graphs are presented. * P<0.05 and *** P<0.001 compared with Primphos group



Graph. 3: Effect of different doses of pranidipine on the mortality after injection of Primphos (percentage)

b. wt.) on Primphos-induced seizures showed that this drug dose-dependently reduced the Primphos-induced seizures (Graphs 1 and 2) Most anticonvulsant effect of pranidipine on the mortality and severity of seizures with a dose of 40 mg/kg was observed (Table 1 and Graph 3).

DISCUSSION

In this study, Primphos cause clonic and tonic seizures and ultimately death. After the mice received intraperitoneal Primphos, some degree of tremor and excessive activity showed that over time the symptoms

became more severe and cause death. In this study, pranidipine dose dependently reduced clonic and tonic seizures and deaths from Primphos. The results showed that the results of different researchers before treatment with calcium channel antagonists, seizure activity creation of different materials down to is in agreement. If pranidipine in preventing tonic convulsions caused by PTZ [4], Aminophyline [9] and pilocarpine [10] have a protective effect, but unlike the above models in all tests anticonvulsant effect has been observed. For example Kainic acid induced seizures, administration of nimodipine before this material could not reduce seizures [7, 8].

Table 1: Effect of different doses of pranidipine on the type of seizures induced by Primphos injection (percentage)

Group	Type of seizures (percentage)		
	Low	Moderate	Severe
Primphos	0	0	100
Pranidipine 2.5+primphos	0	0	100
Pranidipine 5+primphos	0	20	80
Pranidipine 10+primphos	20	40	40
Pranidipine 20+primphos	50	30	20
Pranidipine 40+primphos	60	30	10

In another study showed that calcium channel blockers with values above 80 mg/kg could inhibit tonic seizures from chemicals including PTZ in mice and rats [7, 11, 14], but later showed that high doses of calcium channels blockers cause systemic and cardiac disorders, such as a sharp reduction in coronary blood pressure, decreased movements, imbalance and headache relief [5, 6]. Epilepsy in patients who were resistant to treatment, have reported that nimodipine in an uncontrolled study, seizure frequency is reduced [7, 15], but in another study that two strains were unaware controls, no anticonvulsant effects was observed by nimodipine [16]. Other problems prescription drug, long term administration of drugs with low prescribed intervals (three to four times a day to several weeks) and side effects include headache and hypotension, pronounced the man was from animal models. However, after 24 and 72 hours of administration of nimodipine, percent of alpha and theta waves was increased and vice versa percent in delta waves electroencephalogram was reduced [6, 16]. Other studies have shown that the anticonvulsant effects of calcium channel blockers, with other antiepileptic drugs, increases, For example, in mice and rats with concurrent administration of nimodipine with other drugs can be decreased PTZ-induced tonic seizures, seizures resulting from sound and relieve the electroshock [8, 11-12]. Dihydropyridine calcium channels blockers in experimental seizures by ischemia, bicuculine, electrical cortical shocks, nitrous oxide and alcohol withdrawal syndrome is caused due, have anticonvulsant effects [17]. In another study, calcium channel blockers such as verapamil, nifedipine and Flunarizine to prevent of penicillin-induced seizures and electroencephalogram range have changed [17]. Calcium channel inhibitors on seizures induced by N-methyl-D, L-aspartate (NMDLA) and dihydropyridine calcium channel agonist BAY K 8644 have been effective [5, 18]. In another study on rats have shown that nimodipine in animal models of seizures, nerve

discharge from BAY K 8644 and reduced the decrease in spike-wave EEG [7, 5]. Also have shown that this drug is ischemic brain damage has protective effects [17]. These studies suggest that protective effects of calcium channel antagonists probably due to blocking L-type calcium channels during seizures. These drugs inhibit voltage-dependent calcium channels in seizures, the increase in intracellular calcium to prevent. Well marked that increased Ca²⁺ into the cell in the incidence of certain types of seizures plays a role [4], also marked the loss of calcium outside the cell with reduced flow of calcium from the membranes of neurons for several seconds the discharge of neurons that causes seizures be prevented and the threshold increases [19]. Some of the other antiepileptic drugs such as phenytoin and carbamazepine with a direct effect on neuronal sodium channels act directly or indirectly the flow of calcium ions from the membranes of neurons are inhibited [5, 6].

So it is likely that dihydropyridine calcium channel antagonists to act with similar mechanisms. Also have shown that pranidipine may inhibit calcium, sodium, chloride and potassium and calcium dependent glutamate channels [5].

CONCLUSION

In summary, this study showed that pranidipine (voltage-dependent calcium channel antagonist type L) decreased clonic and tonic seizures from Primphos in mice are probably the main mechanism anticonvulsant related to block calcium channels and reduce calcium flow within neurons. Of course, that this could be generalized to humans rather than question and anticonvulsant effects of dihydropyridine calcium channel antagonists in humans, further investigation is needed.

REFERENCES

1. Khayatnouri, M., F. Hatefi and P. Iagmaie, 2011. Effect of Nifedipine on Primphos-Induced Seizure in Mice. *Global Veterinaria*, 6(6): 551-555.
2. Mohebbi, G.H., A. Jahangiri and P. Hajeb, 2011. Inhibition of Acetyl Cholinesterase Activity Farmers Exposed to Organophosphat Pesticides in Bushehr, Iran. *American-Eurasian J. Toxicological Sci.*, 3(3): 127-129.
3. Khayatnouri, M., P. Iagmaie and F. Hatefi, 2011. Effect of Diltiazem on Primphos-Induced Seizure in Mice. *American-Eurasian J. Toxicological Sci.*, 3(3): 115-119.

4. Khanna, N., S. Bhalla, V. Verma and K.K. Sharma, 2000. Modulatory effects of nifedipine and pranidipine in experimental convulsions. *Indian J. Pharmacol.*, 32: 347-352.
5. Van Luijtelaar, E.L.J.M., N. Ales and A.M.L. Coenen, 1995. Role of L-type calcium channel modulation in nonconvulsive epilepsy in rats. *Epilepsia*, 36(1): 86-92.
6. Kulak, W., W. Sobaniec, K. Wojtal and S.J. Czuczwar, 2004. Calcium modulation in epilepsy. *Pol. J. Pharmacol.*, 56: 29-41.
7. Khayatnouri, M. and S. Bahavarnia, 2011. Effect of Nimodipine on Dichlorvos-Induced Seizure in Mice. *American-Eurasian J. Toxicological Sci.*, 3(3): 204-207.
8. Mikati, M.A., G.L. Holmes, S. Werner, N. Bakker, L. Carmant, Z. Liu and C.E. Stafstrom, 2004. Effects of nimodipine on the behavioral sequelae of experimental status epilepticus in prepubescent rats. *Epilepsy and Behavior*, 5: 168-174.
9. Chakrabarti, A., H. Kaur Saini and S.K. Garg, 1998. Dose-finding study with pranidipine: a selective central nervous system calcium channel blockers on aminophylline induced seizure model in rats. *Brian Res. Bull*, 45: 495-499.
10. Marinho, M.M., V.M. De Bruin, F.C. De Sousa, L.M. Aguiar, R.S. De Pinho and G.S. Viana, 1997. Inhibitory action of a calcium channel blocker (nimodipine) on seizure and brain damage induced by pilocarpine and lithium-pilocarpine in rats. *Neurosci Lett.*, 235: 13-16.
11. Gasior, M., R. Kaminski, T. Burdniak, Z. Kleinrok and S.J. Czuczwar, 1996. Influence of nicardipine, pranidipine and flunarizine on the anticonvulsant efficacy of antiepileptics against pentylenetetrazol in mice. *J. Neural Transm Gen Sect.*, 103: 819-831.
12. Khosla, P. and P. Pandhi, 2000. Anticonvulsant effect of nimodipine alone and in combination with diazepam and phenytoin in a mouse model of status epilepticus. *Methods Find Exp Clin Pharmacol.*, 22: 731-736.
13. Goodman, L.S., L.E. Limbird, J.G. Hardman and A.G. Gilman, 2001. Insecticides. In: Goodman and Gilman's the pharmacological basis of therapeutics. 10th Edition, Mc Graw Hill, New York, USA.
14. Wurlpel, J.N. and S. Iyer, 1994. Calcium channel blockers verapamil and nimodipine inhibit kindling in adult and immature rats. *Epilepsia*, 35: 443-449.
15. De Falco, F.A., U. Bartiromo, L. Majello, G. Di Geronimo and P. Mundo, 1992. Calcium antagonist nimodipin in intractable epilepsy. *Epilepsia*, 33: 343-345.
16. Larkin, J.G., P.J. McKee, J. Blacklaw, G.G. Thompson, I.C. Morgan and M.J. Brodie, 1991. Nimodipine in refractory epilepsy: a placebo-controlled, add-on study. *Epilepsy Res.*, 9: 71-77.
17. Kriz, J., G. Upan and A. Simonia, 2003. Differential effects of dihydropyridine calcium channel blockers in kainic acid-induced experimental seizures in rats. *Epilepsy Res.*, 52: 215-225.
18. Palmer, G.C., M.L. Stagnitto, R.K. Ray, M.A. Knowels, R. Harvey and G.E. Garske, 1993. Anticonvulsant properties of calcium channel blockers in mice: NMDLA and BAYK 8644-induced convulsions are potently blocked by the dihydropyridines. *Epilepsia*, 34: 372-380.
19. McNamara, J.O., 1992. The neurobiological basis of epilepsy. *Trends Neurosci.*, 15: 357-359.