

Prevalence of Ovine Lungworms in and Around Wukro, Tigray Region, Ethiopia

¹Marshet Adugna, ²Berihun Afera and ²Netsanet Berhe

¹National Animal Health Diagnostic and Investigation Center,
²Mekelle University, College of Veterinary Medicine,

Abstract: Cross sectional study to determine prevalence of ovine lung worm from September 2007 to June 2008 was conducted at Wukro, Eastern part of Tigray, Northern part of Ethiopia. A total of 323 sheep fecal sample were collected and examined using the Baermann technique for the presence of larvae and eggs of the parasite. Total prevalence of the disease was 25.69% with 13.63% *Dictyocaulus filaria*, 5.57% *Mullerius capillaries* and *Protostrongylus rufescence* and 6.50% mixed infection. The prevalence in female and male were 25.85% and 25.64% respectively where there is statistically significant ($P > 0.05$). The prevalence in poor, medium and good body conditions were 48.14%, 25% and 21.6% respectively and there is statistical significance difference among the different body conditions ($P < 0.05$). Similarly, the prevalence was 31.58% on sheep of age less than 12 months and 20.46% on sheep greater than 12 months of age in which their difference is statistically significant ($P < 0.05$). The monthly occurrence of infection was higher in January and December with prevalence of 44.45% and 27.28% respectively and their difference was statistically significant ($P < 0.05$). In addition, the disease was also more prevalent in clinically sick animals (56.84%) than those that don't show any clinical signs (12.72%). The prevalence was statistically significant ($P < 0.05$). From this study age, poor body condition and season were the major risk factors for the infection. Depending on the findings strategic deforming of the flock, immunization of sheep and education of farmers to promote good husbandry practices were recommended.

Key words: Lungworm • Prevalence • Ovine • Wukro

INTRODUCTION

Ethiopia is found in the tropical latitude of Africa and has an extremely diverse topography, a wide range of climatic features and multitude of agro-ecological zones which makes the country suitable for different agricultural production system. This has contributed to the existence of a large diversity of farm animal genetic resources [1]. Livestock sector plans a vital role in the national economy of many developing countries including Ethiopia. The current livestock population of the country is estimated to be approximately 38 million, 23million, 18 million, 8 million, 1million and 58 million cattle, sheep, goats, equines, camels and chicken respectively [1]. But this resource has not been exploited as required mainly because of health and nutritional problems, limited genetic potential and other husbandry standards. Losses from inferior weight gains, inedible of organs and carcasses and lower milk yield are not negligible. These production losses to the

livestock industry are estimated at more than 900 million Birr annually. Different viral, bacterial, parasitic and nutritional deficiency diseases which affects production of the livestock were observed in the country such as pox disease, contagious ecthyma, pasturollosis and different internal and external parasite which include helminthes, nematodes and tick and lice infestation as well as mange mites are observed in the animals [2].

Among the disease constraint that causes loss of production is the lung worm infection in sheep. In the past few years many authors have devoted their attention to the prevalence of lung worm in the high lands of temperate areas [3]. The lung worm infection is known to be prevalent in many parts of Ethiopia especially the high land areas and they either cause verminous pneumonia or they predispose animals to other lung disease such as pneumonia [4]. A number of researchers and author reported the prevalence of lung worm in Ethiopia, but the loss it entails is still lacking.

Corresponding Author: Berihun Afera, Mekelle University, College of Veterinary Medicine, Tigray, Ethiopia, P.O.Box. 2084,

- Previously the rate of the diseases in the region was not known because there was no research conducted to determine the prevalence of the disease. In addition, most of the farmer's rare sheep as means of income generation and there are also many irrigation sites in the study site which might predispose the animals to lung worm and other parasitic and none parasitic infection. Therefore the objectives of the study were to determine the prevalence of ovine lung worm in and around Wukro.
- To assess other epidemiological factors associated with the disease

MATERIALS AND METHODS

Study Area: The study was conducted in Wukro, located 45 km from Mekelle, in this area there is marked variation in rainfall, with mean annual rainfall of 62mm. The minimum and maximum temperatures range from 11.8°C to 29.94°C respectively.

Study Design and Sample Size: Cross sectional study was deployed to determine prevalence of lung worms in the selected site. And the sample size was calculated based on the formula given by [5].

$$N = \frac{1.96^2 pq}{d^2}$$

Where N= total sample size, P =expected prevalence and q=1-p and d= absolute precision and in the study 95% level of confidence was used to determine the sample size and the for calculation of the sample size the following information were used and accordingly the sample size was calculated as 139 but to increase the precision of the result a total of 323 sheep were examined

- Total sheep population 26,359
- Prevalence rate of 10%
- Absolute precision of 5%

Study Animals and Characterization: For this study local sheep having different age, sex, clinical signs and body condition from the area were used. Age estimation was estimated using the procedure given by [6]. In addition, the study was carried out in different months of the year.

Study Protocol: Fecal sample was collected directly from rectum. The sample was then transported to wukroworeda veterinary clinics placed in refrigerator until all. Finally, the fecal sample was transported to Mekelle Regional laboratory. The collected specimens were examined with Baermann techniques and sedimentation technique [7].

Data Analysis: The data were entered in to Microsoft excel spread sheet and coded appropriately and for data analysis SPSS version 17 was used and to determine the association of the disease with the risk factors Chi-square (χ^2) were used and significance was considered when P-value is less than 0.05.

RESULTS

In this study a total of 323 sheep were examined for the presence of lung worms. Out of them, 83 were found to be positive with a total prevalence of 25.69% of which 13.62% (44) belongs to *Dictyocaulus filaria*, 5.57% (18), that of *Mullerius capillaries* and *Protostrongylus rufescence* and 6.50% (21) mixed infection as indicated in table 1.

Similarly the prevalence of lung worm was 31.58% (48) in sheep less than 12 months and 20.46% (35) in sheep greater than 12 months age. This indicated that it is statistically significant (P<0.05) as indicated in (Table 2.)

At the same time, the prevalence was 25.64% (30) in males and 25.85% (53) in females and this result where there is no statistically significant difference (P>0.05) as indicated in (Table 3).

At the same time the prevalence of the parasite during the four consecutive months were assessed and rated as 27.27% (21) in December, 44.45% (32) in January, 21.49% (23) in February and 10.44% (7) in March. This shows there is statistically significance (P <0.05) difference among the months (Table 4).

The prevalence rate were 21.6% (27), 25% (43) and 48.14% (13) in good, medium and poor body condition animals respectively as indicated in (Table 5).

As to the clinical signs, examination was made on all sheep where sample has been taken and were divided as with clinical sign and without clinical sign. The rate of the disease in animals with clinical manifestation was 54(56.84%), where as in sheep without clinical sign it was 29 (12.72%) as indicated in table 6.

Table 1: Prevalence of different species of lung worm parasite and their proportion

Species	Total no of positives	Prevalence of each parasites in %	Proportion of each out of positive
<i>D. filarial</i>	44	13.62%	53.01
<i>M. capillaries and P. rufescence</i>	18	5.57%	21.68
Mixed	21	6.50%	25.3
Total	83	25.69%	100

Table 2: Lung worms prevalence in different age groups

Age group	No of sheep in the age	No of sheep infected	Prevalence in %
<12months	152	48	31.58
>12months	171	35	20.46
Total	323	83	25.69

Table 3: Lung worm prevalence in male and female sheep

Sex	No examined	No affected	Prevalence in %
Male	118	30	25.64
Female	205	53	25.85
Total	323	83	25.69

Table 4: Monthly prevalence of lungworms

Months	No of sheep exam	No of sheep infected	Prevalence in %
December	77	21	27.27
January	72	32	44.45
February	107	23	21.49
March	67	7	10.44
Total	323	83	25.69

Table 5: Prevalence of lungworm in different body condition of sheep

Body condition	No of sheep examined	No of sheep infected	Prevalence in %
Good	125	27	21.6
Medium	172	43	25
Poor	27	13	48.14
Total	323	83	25.69

Table 6: Clinical sign versus prevalence

Clinical examination	No of sheep examined	No of sheep infected	Prevalence in %
With clinical signs	95	54	56.84
With out clinical signs	228	29	12.72
Total	323	83	25.69

DISCUSSION

The current finding indicated that the overall prevalence of lung worm parasites in sheep was 25.69% which is lower than that of Yohannes, [8], Wondwossen, [9] Jovanovic [10] and who reported prevalence's of 70.7%, 39.7% and 83-91.5%, in Debretabourawraja,

Assellaawraja and Shoa respectively. At the same time, the current finding is lower compared to [11] who reported prevalence of 37%, in and around Debrebirhan. However, it is higher than that of Frewengel [12] who reported prevalence of 13.24%, in and around Mekelle and that of Fesseha, G. and Gebrenegus T [13] with the rate of 15% in and around DebreZeit.

In the current finding the most prevalent species was *Dictyocaulus filaria* (13.64%) which is almost similar to the previous study conducted by Frewengel [12] who reported 13.24% in and around Mekelle. The prevalence of lung worm was 31.58% in sheep with age <12 months and 20.46% in sheep with age of > 12 months and it was found to be statistically significant ($P<0.05$). The result showed it affects mostly young sheep which might be due to the development of strong immunity in adult sheep following their previous exposure to the parasite [3].

The prevalence in female and male were 25.85% and 25.64% respectively where there is statistically significant ($P>0.05$). This agrees with the study conducted by [12] in which both sexes are equally susceptible to the lung parasites.

Monthly prevalence of lung worm showed that it was higher on December (44.45%) and lowest on March (10.45%) which agrees with Urquhart *et al.* [14] who stated that the disease is more prevalent in temperate areas with high rainfall.

The rate of the disease were 21.6%, 25% and 48.14% in sheep with good, medium and poor body condition scores respectively, where there is significance difference ($P<0.05$) among the different body conditions which might be due to immunosuppression in poor body condition animals and concurrent infection.

At the same time, the rates were 56.84% and 12.71% in those sheep with and without clinical manifestation. Here occurrence of disease without clinical signs appears due to the fact that those sheep which were infected earlier would shed larval without showing clinical signs [3] and those that show clinical signs but without larvae could be due to failure of procedures and existence of other disease conditions which are differential to verminous pneumonia. From the result, it seems that clinical sign would help to diagnose lung worm infection with higher rate.

CONCLUSION AND RECOMMENDATIONS

The current finding revealed that the disease is more prevalent in the study site characterized by high rate of lung worm parasites in ovine species which need particular attention by most of the professionals and the government as it affects both the production and productivity of small ruminants..

- Use of strategic deforming and prophylactic anthelmintic drugs should be monitored.

- Immunization of sheep with Gamma irradiated *Dictyocaulus filaria* larvae.
- Creation of public awareness to protect the disease as well as the production loss.
- Effective information exchange system should be instituted between field veterinarians and researchers for better understanding of the epidemiology of the disease

ACKNOWLEDGEMENTS

We wish to extend our sincere appreciation to Mekelle University, college of veterinary medicine and also the veterinarians of Wukro who have great contribution during sample collection and processing.

REFERENCES

1. Anon, 2005. State of Ethiopian animals, genetic resources. Country report. A contribution to the first report on the Worlds animal genetic resources, institute of Biodiversity Conservation, Addis Ababa, Ethiopia, pp: 74.
2. Getachew, T., A. Mulugeta, J. Carbal and R. Lalic, 1988. Studies on the effect of vaccination and anthelmintics treatment of ovine lung worms *D. filaria* infection in sheep.
3. Soulby, E.J.L., 1982. Helminthes, Arthropods and Protozoa of Domesticated Animals, 6th ed., Bailliare, Tindall, Londons, pp: 492-552.
4. Kinfe, G., 1980. Fenbendazol Anthelmintic efficacy in naturally infected local sheep with gastrointestinal nematodes and lung worm Ethiopian Bulletin, V1(2): 44-45.
5. Thrust field, 1995. Veterinary epidemiology 2nd ed. pp: 182-183.
6. Aiello, S.E. and A. Mays, 1998. The Merck Veterinary Manual, 8th ed. (Merck and Co Inc., White House station, NJ) pp: 131-140.
7. Charles, M., Hendrix and E.D. Robinson, 2006. Diagnostic parasitology for veterinary technician's, pp: 243.
8. Yohannes, G., 1989. Epidemiological study and anthelmintic treatment trial of ovine Dictyocauliasis in Debre-tabor Awaraja DVM Thesis, Addis Ababa University, Debre-zeit, Faculty of Veterinary Medicine, Ethiopia, pp: 48.
9. Wondwossen, T., 1992. Prevalence of ovine lungworms in and around Assela, DVM Thesis, Addis Ababa University, Faculty of Veterinary Medicine, Debre-Ziet, Ethiopia, pp: 47.

10. Jovanovic, M., 1992. Observations of *D. filaria* infection in and around Shoa, a Report to the Ministry of Agriculture, Addis Ababa, Ethiopia, pp: 12.
11. Natsanet, B., 1992. Study on prevalence and control of lungworm in local Ethiopian highland sheep in and around DebreBerhan, DVM Thesis, Addis Ababa University, Faculty of Veterinary Medicine, Debrezeit, Ethiopia, pp: 42.
12. Frewengel, S., 1995. Prevalence of ovine lung worm in and around Mekelle, DVM Thesis, Addis Ababa University, Faculty of Veterinary Medicine, Debrezeit, Ethiopia, pp: 24.
13. Fesseha, G. and T.M. Gebrenegus, 1977. Epidemiology of *Dictyocaulus filaria* in area around Debre-zeit, Modjo and note in its prevalence at Alemaya, Report to Ministry of Agriculture, Addis Ababa.
14. Urquhart, G.M., J. Armor, J.L. Duncan, A. Dunn and F.W. Mand Jening, 1987. Veterinary Parasitology 7th edition long man English language society, pp: 39-40, 57, 58.