

Evaluation of Haemolymph Components of Infected and Non Infected *Lymnaea acuminata* (Pulmonata: Lymnidae) with Xiphidiocercariae

Farindra Tiwari

Department of Zoology, M.G.P.G. College, Gorakhpur, 273001 UP, India

Abstract: In this study the haemolymph components of infected and none infected *Lymnaea acuminata* with xiphidiocercaria larvae was compared. Five hundred Fifty *L. acuminata* snails were collected from Gorakhpur region of eastern Uttar Pradesh. The snails were transported to the lab at MGPG college and cercarial shedding were studied. Haemolymph of snails were extracted and cells were counted using haemocytometer and cell-surface carbohydrate were recognized by conjugated lectin (Lentil). Haemolymph protein concentrations were measured by Bradford protein assay method and soluble protein compositions were determined on sodium dodecyl sulphate polyacrilamide gel electrophoresis (SDS-PAGE). From the 550 *L. acuminata* snails examined for cercariae, 27 snails were infected with xiphidiocercariae. Mean of haemolymph cells (haemocyte) number were obtained 93480 ± 2.43 (cells/ml) for none infected snails (25 snail) and 124560 ± 2800 (cells/ml) for infected snails (25 snail). Mannose carbohydrate was recognized on haemocyte of none infected and infected snails. Mean of protein concentration of haemolymph plasma was obtained as 1354 ± 160 $\mu\text{g/ml}$ (1.4 mg/ml) for none infected snails (25 snails) and 1802 ± 138 $\mu\text{g/ml}$ (1.8 mg/ml) for infected snail (25 snails). Comparing to none infected snails, the SDS-PAGE results of haemolymph plasma of infected snails, showed an extra protein band (70 kDa). The results showed a significant difference between the amounts and the kinds of proteins in haemolymph of infected and none infected snails. This information might be useful to understand of parasite detection, adhesion, engulfment and antigen agglutination by snail.

Key words: Cercaria • *Fasciola* • Snails

INTRODUCTION

Snail is one of the important components of an aquatic ecosystem, it acts as intermediate host of *Fasciola* species. Control of snail population below a certain threshold level is one of the important methods in the campaign to reduce the incidence of fascioliasis. Life cycle of the parasite can be interrupted by killing the snail or *Fasciola* larva redia and cercaria in the snail body. Fascioliasis is a worldwide zoonotic disease caused by *Fasciola hepatica* and *Fasciola gigantica* (Fasciolidae) [1]. *F. hepatica* has worldwide distribution but predominates in temperate zones while *F. gigantica* is found primarily in tropical regions [2-4]. The definite host is very broad and includes many herbivorous mammals including human being. Human fascioliasis has been reported in 51 different countries from five continents [4]. Fascioliasis is now recognized as an emerging human

disease. World health organization has estimated that 2.4 million people are infected with *Fasciola* and a further 180 million are at risk of infection [1]. Singh and Agarwal [5] reported that 94% of buffaloes slaughtered in local slaughter houses in Gorakhpur district are infected with *F. gigantica*. In northern India, *Lymnaea acuminata* is the intermediate host of the *Fasciola* species [2]. Although control of snail population below a threshold level is one of the important methods for effective control of fascioliasis [6-9], yet snails are one of the important components in the aquatic ecosystem. The *Fasciola* larval stages sporocyst, redia and cercaria in the snail body are in division phase of *F. gigantica*. Cercaria is a larval form of the parasite, developed within the snail. It finds and settles in a selective host and according to species, metacercaria, or adult form will be appeared. Xiphidiocercaria is cercariae with a stylet in the anterior rim of its oral sucker with which it actively penetrates its

host. Studies on cercariae infections in fresh water snails help us to understand snail-borne diseases presence and the location of potential transmission sites [10]. Several studies have examined the influence of parasites on the host organisms [2-4]. Cercariae identification of *Lymnaea acuminata* snails may help to explain the geographical distribution of diseases in India. Haemolymph analysis of infected snail to parasite also has potential to develop knowledge of parasites immunology. Therefore, the aim of this study was to compare haemolymph components of infected and none infected *L. acuminata* snails with xiphidiocercaria larvae.

MATERIALS AND METHODS

Cercariae Collection and Haemolymph Extraction from Snails:

Adult *Lymnaea acuminata* (2.6 ±0.20 cm in length) were collected locally and cercaria shedding infected and uninfected snails were separated in two groups. The snails were allowed to acclimatize for 24 hours in laboratory condition. Each infected snail was dissected in a glass petridish containing 10mL of dechlorinated water at 22-24°C. The pH of the water was 7.1-7.3 and dissolved oxygen, free carbon dioxide and bicarbonate alkalinity were 6.5-7.2mg/L, 5.2- 6.3mg/L and 102.0-105.0mg/L, respectively. Totally, 550 *L. acuminata* snails were collected from Gorakhpur region of eastern Uttar Pradesh by sweeping a net through the agricultures canals and drains. The snails were transported by wet cotton to the lab at M.G.P.G. College. In the lab, the snails were maintained in plastic containers. *L. acuminata* snails were fed by lettuce 1-2 times per week and the water in the containers was changed once 24-48 hrs. To collect cercariae, snails were examined by mean of the shedding method and collected cercariae were identified as fresh or stained samples by taxonomic key [6]. Haemolymphs of cleaned snails were extracted by capillary tube from the haemocoel of foot-and-mouth part. A haemocoel is a cavity or series of spaces between the organs of snail with open circulatory system [7].

Cell Counting and Cell-Surface Carbohydrate Recognition:

Extracted haemolymph was introduced into one of the V-shaped wells of Neubauer haemocytometer with a capillary tube and determined cell numbers [8]. Extracted haemolymphs were centrifuged at 3000 rpm for 5 minute at 4°C in microfuge and haemocytes were collected from the pellet. To recognize of cell-surface carbohydrate, FITC (fluorescein isothio cyanate) -

conjugated lectin was used. Then, FITC-lectin was added to the suspension of the cells in the test tubes and added FITC-lectin to the control tubes containing 100 mM inhibitory sugar (mannose). The tubes were incubated at 4°C for 60 min and the cells were washed three times by centrifugation (3000 rpm for 5 min) in PBS. Samples were mounted on slides and studied by fluorescence microscope [9].

Measurement of Protein Concentration and Detection of Protein Bands:

Extracted haemolymph samples were centrifuged at 8500 rpm for 30 min at 4°C by microfuge and plasma specimens were obtained. Concentrations of total proteins were measured by Bradford assay method. Sodium dodecyl sulphate Polyacrylamide geleelectrophoresis (7.5% gel and Coomassie blue staining) was used to separate the components of plasma protein [9].

Statistical Method: The independent *t*-test was used to distinguish the significant difference between two studied groups on the means of cell or protein concentrations in none infected and infected snails [10].

RESULTS

From the total number of snails, which examined for cercariae, 27 snails were found infecting with xiphidiocercaria (Fig. 1). The number of haemocytes was 93480±2043 (Cells/ml) for none infected snails and 124560 ±2800 (Cells/ml) for infected snail (Table 1). Mannose monosaccharide, CH₂OH (CHOH)₄CHO, was detected as surface carbohydrate on the cell at a 1/25 diluted FITC-lectin. Because of covering of FITC-lectin by inhibitory sugar on cells in the control samples, cells were not observable at above dilution under fluorescence microscope (Fig. 2). Total protein concentrations of haemolymph were estimated 1354 ±160µg/ml (1.4 mg/ml) for none infected snails and 1802 ± 138µg/ml (1.8mg/ml) for infected snails (Table 1). An extra-soluble protein (70 kDa) was detected on SDS-PAGE of haemolymph plasma of infected snails that is presented in Fig. 3. Independent samples *t*-test revealed there was significant difference in cell concentrations between infected snail (M = 124560, SD = 2800) and none infected (M = 93480, SD = 2043) conditions; *t* (48) = 2.01, *t*-value = 44.83, *P* < 0.05. There was also significant difference in protein concentration between infected snail (M= 1802, SD= 138) and none infected snail (M= 1354, SD=162) conditions; *t* (48) = 2.01, *t*-value = 10.66, *P* < 0.05.

Table 1: Number of haemocytes and protein concentration of haemolymph of none infected and infected snails *L. acuminata*.

Observations	None Infected Snails	Infected Snails
Number of Haemocytes (cells/ml)*	93480±2043	124560 ±2800
Protein concentration of haemolymph (µg/ml)*	1354±160µg/ml (1.4mg/ml)	1802±138µg/ml(1.8mg/ml)

*Significant difference was observed in between none infected and infected group of snails when independent *t*-test applied




Fig. 1: Xiphidiocercaria larvae from *L. acuminata* snail; A. Sporocyst with developing cercariae; B. Cercaria with stylet in oral sucker



Fig. 2: A. Haemocyte surface with mannose carbohydrate, which was conjugated by FITC-Lentil. B. Haemocyte control without surface mannose




Fig. 3: SDS-PAGE analysis of plasma proteins of extracted haemolymphs from infected (lane 1) and none infected (Lanes 2, 3, 4, 5, 6) *L. acuminata* snail and protein marker (lane 7) [S4]

DISCUSSION

In previous studies, it has been shown that xiphidiocercaria larvae belong to Plagiorchiidae Trematodes [5]. Recently, xiphidiocercariae has been reported from *Lymnaea natalensis* snail (10). There are few publications about xiphidiocercaria effects on the haemolymph compartments [11-12]; however, various

reports have been recorded on the effects of other cercariae on haemolymph of other snails including protein level. *Schistosoma mansoni* infection has increased the total protein concentration of haemolymph of *Biomphalaria alexandrina* [13]. In the present study, significant increase of protein concentration may be representing of humoral factors (opsonin and agglutinin) in haemolymph to parasite presence [14]. The use of

specific inhibition sugar has been indicated presence of glucose/ mannose, galactose, N-acetyl-galactosamine (GalNAc) and N-acetyl-glucosamine (GlcNAc) on the surface of the haemocytes [15]. Lectins are responsible for cell surface sugar recognition; therefore, based on the current study, existence of mannose carbohydrate on the cells surface helps them for detection and adhesion to parasite larva through crosslink glyco conjugates [16]. Effects of *Tricho bilharzias ocellata* haemocytes of *Lymnaea stagnalis* with respect to cell number and distribution of sub populations has been determined [17]. The present work reports significantly increase of cell number, which is very important for phagocytosis or engulfment of larval stages of parasite (miracidium, sporocyst and cercaria) [14]. Two snail populations with respect to haemolymph agglutinins (lectin) from *Lymnaea stagnalis* have been reported [18]. Lectins are considered to play main role in humoral immunity of molluscs serving as recognition factors for foreignness [19]. SDS-gel-separated protein band from this research might be a member of the lectinic protein family agglutinate, however, need to more characterizations by isolation, purification and identification.

In conclusion, the significant increase of the cells number and protein concentrations, as well as presence of cell surface mannose and extra soluble proteins in the haemolymph of infected snails with xiphidiocercaria larva might have an important role in parasite detection, adhesion, engulfment/antigen agglutination.

REFERENCES

1. Chingwena, G., S. Mukaratirwa, T.K. Kristensen and M. Chimbari, 2002. Larval trematode infections in freshwater snails from the Highveld and Lowveld areas of Zimbabwe. *J. Helminthology*, 76: 283-293.
2. Bezerra, J.C.B., W. Becker and U.E. Zelck, 1997. A comparative study of the organic acid content of the hemolymph of *Schistosoma mansoni*-resistant and susceptible strains of *Biomphalaria glabrata*. *Mem. Inst. Oswaldo. Cruz., Rio De Janeiro*, 92(3): 421-425.
3. Rupperecht, H., W. Becker and A. Schwanbek, 1989. Alterations in hemolymph components in *Biomphalaria glabrata* during long term infection with *Schistosoma mansoni*. *Parasitol. Res.*, 75(3): 233-7.
4. Loker, E.S., 1987. Alterations in *Biomphalaria glabrata* plasma induced by infection with the digenetic trematode *Echinostoma paraensei*. *J. Parasitol.*, 73(3): 503-13.
5. Farahnak, A. and N. Dabagh, 2008. Adhesion of Cercaria (Larva of Helminth Parasites) to Host by Lectins- carbohydrates bonds as a Model for Evaluation of *Schistosoma* Entrance Mechanisms in Cercarial Dermatitis. *Iranian J. Publ. Health.*, 37(2): 59-63.
6. Stewart, C.S., 1970. *The Trematodes*. M.C. Dubuque, Iowa: Brown Company Publisher.
7. Malek, E.T.A., 1962. *Laboratory guide and notes for medical malacology*. Minneapolis: Burgess Pub. Co.
8. Caprette, D.R., 2007. *Experimental bioscience; Laboratory methods: Using a Counting Chamber* [online]. Available from: <http://www.ruf.rice.edu/~bio-slabs/methods/microscopy/cellcount.html>.
9. Maizels, R.M., M.L. Blaxter, B.D. Robertson and M.E. Selkirk, 1991. *Parasite antigen and parasite genes: A laboratory manual for molecular parasitology*. 1st ed. Cambridge: Cambridge University Press, pp: 93-94.
10. Moema, E.B., P.H. King and C. Baker, 2008. Cercariae developing in *Lymnaea natalensis* Krauss, 1848 collected in the vicinity of Pretoria, Gauteng Province, South Africa. *J. Vet. Res.*, 75(3): 215-23.
11. Narayanan, R., 1980. Venkateswararao P. Effect of xiphidiocercarial infection on oxidation of glycolytic and Krebs cycle intermediates in *Lymnaea luteola* (Mollusca). *J. Invertebr Pathol.*, 36(1): 21-24.
12. Reddy, B.R. and P.V. Rao, 1992. Effect of xiphidiocercarial infection on the purine nucleotide cycle activity in the fresh water pulmonate snail, *Lymnaea luteola*. *J. Invertebr Pathol.*, 60(2): 117-120.
13. El-Emam, M.A. and F.A. Ebeid, 1989. Effect of *Schistosoma mansoni* infection, starvation and molluscicides on acid phosphate, transaminases and total protein in tissues and hemolymph of *Biomphalaria alexandrina*. *J. Egypt. Soc. Parasitol.*, 19(1): 139-47.
14. Sminia, T. and W. Knaap, 1981. The internal defence system of the freshwater snail. *Developmental and Comparative Immunology*, 5, Supplement 1: 87-97.
15. Mohamed, A.H., 2005. Characterization of surface lectins binding and SDS-PAGE protein patterns of *Biomphalaria alexandrina* haemocytes infected with *Schistosoma mansoni*. *J. Egypt Soc. Parasitol.*, 35(2): 615-30.
16. Weis, W.I. and K. Drickamer, 1996. Structural basis of lectin-carbohydrate recognition. *Annu. Rev. Biochem.*, 65: 441-73.

17. Amen, R.I., J.M. Tijnagel, W.P.W. Knaap, E.A. Meuleman, E.S. Klerk and T. Sminia, 1991. Effects of *Tricho bilharzia ocellata* on hemocytes of *Lymnaea stagnalis*. *Dev Comp Immunol.*, 15(3): 105-15.
18. Knaap, W.P.W., A. Doderer, H. Boerrigter-Barendsen and T. Sminia, 1982. Some Properties of an Agglutinin in the Haemolymph of the Pond snail *Lymnaea stagnalis*. *Biol. Bull.*, 162: 404-412.
19. Horák, P. and W.P.W. Knaap, 1997. Lectins in snail-trematode immune interactions: a review. *Folia Parasitologica*, 44: 161-172.