

Effects of Different Artificial Diets on Growth Performance and Body Composition on Rainbow Trout (*Onchorynchus mykiss*) (Walbaum, 1792) Fingerlings

¹M. Hosein. Jalae, ²Moein Faramarzi, ³Saeed Kiaalvandi and ⁴Farnaz Iranshahi

¹Department of Fisheries of Golestan province, Gorgan, Iran

²Department of Fishery, Gonbad University, Gonbad, Iran

³Department of Fishery, Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran

⁴Department of Fishery, Shahid Bahonar University, Kerman, Iran

Abstract: Rainbow trout (*Onchorynchus mykiss*) (Walbaum, 1792) is an important species in aquaculture. In this study, we evaluated effects of replacing fish meal with poultry by-product meal. Four isocaloric and isonitrogenous rations containing various levels (0% or control), 33%, 67% and 100% of fish meal) of poultry by-product meal (PBM) were fed to three replicate groups of Rainbow trout (*Onchorynchus mykiss*) fingerlings with a mean initial weight of 15.40±0.03 g. 50 fish per tank were tested for 10 weeks in 500 liter fibre glass tanks. Average weight gain of Rainbow trout fingerlings fed the control were significantly ($P<0.05$) higher (42.63±0.66) compared to fish fed 33%, 67% and 100% PBM of fish meal (30.14±0.06, 25.91±0.48 and 19.77±0.07, respectively). Significant variation in feed conversion ratios which varied between 1.71±0.02 and 2.81±0.03 for the control and 100% PBM, respectively, were obtained among the groups. Similarly, specific growth rate and protein efficiency ratio decreased significantly ($P<0.05$) as the level of PBM increased. However, condition factor, dress out percentage and the whole body composition didn't exhibit any significant variation among the test groups.

Key words: Diet • Rainbow trout • *Onchorynchus mykiss* • Poultry by-product meal • Fish meal

INTRODUCTION

Fish meal is the major protein source in aquaculture feeds. However, the supply of fish meal is not growing worldwide [1, 2] and it depends entirely on landings from the capture fisheries. Fish meal production from Peru and Chile, two countries producing about two-thirds of annual global production, fluctuates periodically by over 20% in El Nino years when ocean temperatures warm up and cause the fish stocks to move offshore, out of reach of the fishery [3].

Moreover, price of fish meal is often high. These necessitate replacing fish meal with cheaper protein sources [4]. One of the alternative ingredients to fish meal is poultry by-product meal (PBM). PBM is made of ground, rendered, or clean parts of the carcass of slaughtered poultry. PBM has been tested at varying success so far in coho salmon [5], chinook salmon [6], rainbow trout [7-10], tilapia [11, 12], sea bream [13], European eel [14], channel catfish [11], common carp, catla, rohu [15-17], sunshine bass [18] and Pacific white shrimp [19].

Fowler [6] and Sevgili [10] reported PBM could replace about 50% of fish meal in the diets for chinook salmon and rainbow trout. Hasan and Amin [20] found that processing techniques greatly affected the nutritional quality of PBM for *Cirrhinus mrigala* fry. They reported that autoclaved and boiled PBM showed better growth performances than sundried and/or oven dried PBM. Dong *et al.* [21] drew attention to the nutritional quality differences of PBM produced by different manufacturers. There is a lack of information on nutritional quality of PBM produced in Iran for fish diets. This study was planned to determine the level of PBM that could be used to replace fish meal in practical diets for Rainbow trout (*Onchorynchus mykiss*) fingerlings.

MATERIALS AND METHODS

Ingredients used in the study were purchased from local market. According to information provided by the manufacturer, PBM used consists of chicken slaughter wastes including viscera, heads, legs and feather and was produced by exposing to 150-200°C under a 2.5-atm

Table 1: Proximate composition of fish meal and PBM (as is basis) used in trial diets

	Fish meal	Poultry by-product meal
Dry matter (%)	93.16	94.23
Crude protein (%)	70.44	52.12
Crude fat (%)	7.36	23.47
Crude ash (%)	11.18	18.34

Table 2: Formulation and composition of the experimental diets (%)

Ingredients	Control	33%	67%	100%
Fish meal	30	20	10	0
Soy bean meal	15.5	15.5	15.5	15.5
PBM	0	12	24	36
Wheat middlings	31.5	30.35	29.2	28.1
Wheat	17.0	17.8	18.6	19.4
Choline chloride	0.1	0.1	0.1	0.1
Vitamin premix ¹	0.5	0.5	0.5	0.5
Mineral premix ²	0.1	0.1	0.1	0.1
Binder ³	0.3	0.3	0.3	0.3
Sunflower oil	5	3.35	1.7	0
Total	100	100	100	100
Nutrients levels determined by analysis(as is basis)				
Dry matter	90.99	91.31	90.47	91.14
Crude protein	36.49	36.37	35.0	33.83
Crude fat	7.72	8.7	9.23	9.78
Crude ash	6.48	6.92	7.63	8.26
Crude fiber	2.4	2.4	2.99	2.33
Nitrogen free extract	37.9	36.92	35.62	36.94
Gross energy(kcal/ 100 g) ⁴	444.7	451.1	443.7	445.1

¹Per kg premix: 4,000,000 IU vitamin A, 480,000 IU vitamin D3, 40,000 mg vitamin E, 2,400 mg vitamin K3, 4,000 mg vitamin B1, 6,000 mg vitamin B2, 40,000 mg niacin, 10,000 mg Ca-pantothenate, 4,000 mg vitamin B6, 10 mg vitamin B12, 100 mg D-biotin, 1,200 mg folic acid, 40,000 mg vitamin C and 60,000 mg inositol.

² Per kg premix: 23,750 mg Mn, 75,000 mg Zn, 5,000 mg Zn, 2,000 mg Co, 2,750 mg I, 100 mg Se, 200,000 mg Mg.

³ Lignosulfate

⁴ Gross energy based on 5.65, 4.1 and 9.5 kcal / g protein, carbohydrate and fat, respectively.

pressure for ten hours. Both fish meal and PBM were analyzed for proximate composition prior to the formulation of diets (Table 1). Four isonitrogenous and isocaloric diets were formulated to evaluate nutritional value of PBM for Rainbow trout fingerlings (Table 2). The control diet contained 30% of fish meal and 15.5% of soy bean meal as main protein sources.

PBM was tested at three inclusion levels (33%, 67% and 100% replacement of fish meal) by reducing fish meal levels. The diets were prepared by mixing the dry ingredients and oil, followed by the addition of water until stiff dough was obtained. The moist diet was extruded through a mincer with a 2 mm die. The resulting pellets were then dried on the shelves at the room temperature. The diets were stored in the plastic bags under ambient conditions over the experimental period.

The feeding trial was conducted in outdoor fibre glass tanks with holding capacity of 12,500 L. Each tank was supplied with a water flow of 10 l/min. Over the experimental period, water temperature (°C), dissolved oxygen (mg/l) and pH changed between, 26.1-27.5, 8.0-8.7 and 7.3-7.8, respectively. Fish fry (initial mean weight 15.40±0.03) were randomly allocated at a stocking rate of 50 fish per tank with three replicate tanks for each experimental diet.

All fish were fed two times daily at a fixed feeding rate of 4 % body weight per day for ten weeks. Total biomass of the fish from each tank was weighed at biweekly intervals and feeding rates adjusted accordingly. At the beginning of the trial, 25 fish and at end of the trial, 5 fish per tank were sampled to determine the whole body composition. 5 fish per tank were also picked at the end of

the study to determine condition factors and dress out percentage of the groups fed different PBM levels.

Fish performance, including average weight gain (AWG), feed conversion ratio (FCR), specific growth rate (SGR), protein efficiency ratio (PER), condition factor (CF) and dress out percentage (DOP) were determined as described by Metailler [22] and Goddard [23]. The protein content of the diets and the whole body was determined by kjeldahl, fat by solvent extraction, ash by placing the samples in a muffle furnace (550°C) for 12 h, fiber by placing the samples remaining in a muffle furnace (550°C) for 6 h after acid and alkali hydrolysis and moisture by drying (105°C) until constant weight has been attained. Nitrogen free extract was calculated by subtracting the protein, fat, fiber and ash from the dry matter [24]. Results were analysed by a one-way analysis of variance and the treatment means compared by Duncan's multiple range tests. Significance was tested at the P<0.05 level.

Results are presented as mean ± SE. Statistical analyses were performed with SPSS 16 for windows statistical package.

RESULTS

The growth response and performance data of Rainbow trout juveniles fed diets containing various inclusions of PBM are presented in Table 3. Weekly growth responses of Rainbow trout juveniles over the experimental period are shown in Figure 1.

The performance of Rainbow trout juveniles differed significantly (P<0.05) in terms of final weight, AWG, FCR, SGR and PER.

Growth responses were lower in groups fed diets with PBM compared to those fed the control diet. The growth performance worsened in fish fed diet containing even lowest PBM level (33%) and kept worsening as the level of PBM increased. However, there were no significant differences in CF and DOP among the groups (Table 3). Initial and final body compositions of fish were presented in Table 4. There were no significant differences in body composition among the treatments.




Fig. 1: Live weight variations in Rainbow trout juveniles over the experimental period

Table 3: Growth, FCR, SGR, PER, CF and DOP of Rainbow trout fry after 10 weeks

	Control	33%	67%	100%
Initial weight (g)	15.50±0.06	15.43±0.15	15.40±0.06	15.33±0.13
Final weight (g)	58.13±0.64 ^a	45.57±0.20 ^b	41.31±0.45 ^c	35.10±0.12 ^d
AWG (g)	42.63±0.66 ^a	30.14±0.06 ^b	25.91±0.48 ^c	19.77±0.07 ^d
FCR	1.71±0.02 ^d	2.24±0.01 ^c	2.42±0.06 ^b	2.81±0.03 ^a
SGR (%/day)	1.89±0.02 ^a	1.55±0.01 ^b	1.41±0.02 ^c	1.19±0.01 ^d
PER	1.46±0.02 ^a	1.23±0.07 ^b	1.07±0.03 ^b	0.96±0.01 ^c
CF	2.75±0.06	2.67±0.07	2.63±0.06	2.56±0.04
DOP (%)	85.83±1.62	85.65±1.26	84.54±0.46	85.55±0.49

Values are means± SE for three replications. Figures in the same row with different superscripts are significantly different (p<0.05).

AWG = Final weight (g)-Initial weight (g)

FCR = Total feed (g)/Total weight gain (g)

SGR = 100 (ln Wf - ln Wi)/time (days) where Wf is final weight, Wi is initial weight.

PER = Wet weight gain (g) / Protein fed (g)

CF = Final weight (g) / Fork length (cm) X 100

DOP = Viscera (g) / Whole weight (g) X 100

Table 4: The whole body composition analyses (on wet weight basis) of Rainbow trout fed test diets

Component (%)	Initial	Control	33%	67%	100%
Dry matter	26.64	28.96±1.40	29.50±0.30	29.38±0.20	29.49±0.16
Protein	16.35	15.66±0.86	15.55±0.12	15.40±0.16	15.88±0.51
Fat	7.50	11.51±0.72	11.97±0.35	12.29±0.33	11.49±0.58
Ash	2.67	1.56±0.10	2.00±0.10	1.72±0.22	1.87±0.16

Values are represented as means± SE for three replications.

DISCUSSION

PBM seems to be a good source of dietary protein for fish culture. Higgs *et al.* [5] found that defatted PBM and PBM mixed with hydrolysed feather meal could replace up to 33% and 75 % of fish meal, respectively, in coho salmon diets. About 50 % of fish meal was successfully replaced with PBM in chinook salmon and rainbow trout [6, 7, 9, 10]. Moreover, Gouveia [8] reported PBM mixed with hydrolysed feather meal could be used without growth retardation at a level of 80 % of total protein in trout diets. Nengas *et al.* [13] compared PBMs produced via old (OTPBM) or high technology (HTPBM) in sea bream diets and found that while OTPBM reduced growth performances at high inclusion levels, HTPBM did not effect negatively growth at a level of 100 % of fish meal.

In Pacific white shrimp and sunshine bass diets, 80 % and 100 % of fish meal replacement with PBM, respectively, did not effect negatively weight gain and feed conversion ratio. The results obtained from European eel [14], tilapia [12], catla [16], rohu [17] and carp [15] indicate that total replacement of fish meal with PBM could be possible. In the present study, the diets containing PBM even at lowest level significantly limited weight gain, FCR, SGR and PER. The values worsened as PBM level increased. These results are contradictory to ones mentioned in previous paragraph. Poor performance of this material may be due to; i) limiting amino acid (histidine, methionine+cystine, lysine and phenylalanine) content [10, 13, 25], feather, connective tissue and skin contents which are considered to be difficult for fish to digest [26-29], subjection of the product to high temperature (150-200°C) for a long time (10 hours) during the processing [13,10], or combination of all. High temperature during the raw material processing leads to lysine, cystine+cystine losses and thus, digestibility of protein and amino acids is reduced [30-31].

The reason of reduced growth performances of fingerlings fed diets with PBM was not their fat contents. Optimal dietary fat levels have been suggested to be below 12% in the practical diets of cyprinids [32]. In the present study, the highest fat level was 9.78% which is in the range of the optimal levels. Moreover, Gallagher and Degani [33] successfully used 10% poultry oil as a replacement of fish oil for the diets of European eels. Body compositions of Rainbow trout fry fed diets containing various levels PBM were not significantly differed in the present study. These findings are in agreement with the values reported by Hasan *et al.* [16],

Nengas *et al.* [13] and Sevgili [10]. The final body dry matter and lipid levels are higher [6, 16, 34] and ash is lower [20] than initial levels. However, final body protein level obtained in this trial was slightly lower compared to initial value. This is contradictory findings reported by Fowler [6], Gouveia [8] and Hasan and Amin [20]. Consequently, our results indicate that growth of Rainbow trout fingerlings was negatively effected with PBM levels. Dong *et al.* [21] found differences in proximate composition and protein digestibility among the samples of PBM from different manufacturers.

In our previous study [10], however, PBM from the same manufacturer was used to replace fish meal in practical rainbow trout diets. In this study, it was found that PBM could be used up to 20 % of diet as a protein source. Unlike nutritional disadvantages of PBM compared to fish meal, it is much cheaper and more easily available than fish meal. Thus, the level of PBM that can be used in diets of Rainbow trout fingerlings needs to be further evaluated.

REFERENCE

1. Rumsey, G., 1994. What is the future of fish meal use? Feed International., 15: 10-16.
2. Barlow, S., 1997. Fish meal-supply limits demand. Feed Tech., 1(1): 34-35.
3. Hardy, R.W., 1996. Alternate protein sources for salmon and trout diets. Animal Feed Science and Chnol., 59: 71-80.
4. Shepherd, T., 1998. Rendered products in aquaculture feeds. S. Frasen (Ed.), International Aqua Feed., 4: 13-17.
5. Higgs, D.A., J.R. Markert, D.W. Macourarie, J.R. McBride, B.S. Dosanjh, C. Nichols and G. Hoskins, 1979. Development of practical dry diets for coho salmon, *Oncorhynchus kisutch*, using poultry by-product meal, feather meal, soybean meal and rapeseed meal as major protein sources. K. Tiews and J.E. Halver (Eds.), Finfish Nutrition and Fish Feed Technology, Vol. II, Hieennemann GmbH, Berlin, pp; 191-218.
6. Fowler, L.G., 1991. Poultry by product meal as a dietary protein source in fall chinook salmon diets. Aquaculture, 99: 309-321.
7. Alexis, M.N., E.P. Papoutsoglou and V. Thecohari, 1985. Formulation of practical diets for rainbow trout (*Salmo gairdneri*) made by partial or complete substitution of fish meal by poultry by products and certain plant by-products. Aquaculture, 50: 61-73.

8. Gouveia, A.J.R., 1992. The use of poultry by-product and hydrolised feather meal as a feed for rainbow trout (*Oncorhynchus mykiss*). Publicacoes do Instituto de Zoologia., 227: 24.
9. Steffens, W., 1994. Replacing fish meal with poultry byproduct meal in diets for rainbow trout, *Oncorhynchus mykiss*. Aquaculture, 124: 27-34.
10. Sevgili, H., 2002. Gokkusagi Alabaligi (*Oncorhynchus mykiss*) rasyonlarında tavuk mezbaha artıkları ununun, balık unu yerine kullanılma olanakları. MSc thesis. Antalya: University of Mediterranean.
11. Sadiku, S.O.E. and K. Jauncey, 1995. Soybean flour-poultry meat meal blend as dietary protein source in practical diets of *Oreochromis niloticus* and *Clarias gariepinus*. Asian Fisheries Sci., 8: 159-167.
12. El-Sayed, A.F.M., 1998. Total replacement of fish meal with animal protein sources in Nile tilapia, *Oreochromis niloticus* (L.) feeds. Aquaculture Res., 29: 275- 280.
13. Nengas, I., M.N. Alexis and S.J. Davies, 1999. High inclusion levels of poultry meals and related by products in diets for gilthead sea bream, *Sparus aurata* L. Aquaculture., 179: 13-23.
14. Appelbaum, S., V. Birkan and A. Prilutzly, 1996. Use of chicken meal as a substitute for fish meal in the diet of young eels. Archives of Polish Fisheries, 4: 141-145.
15. Steffens, W., 1988. Utilization of poultry by-products meal for raising carp fingerlings (*Cyprinus carpio*). Archives of Animal Nutrition, 38: 147-152.
16. Hasan, M.R., A.M. Akand and A. Siddiqua, 1993a. Studies on poultry offal meal and silk worm pupae meal as dietary protein sources for Indian major carp, *Catla catla* (Hamilton). Bangladesh J. Training and Development, 6: 55-66.
17. Hasan, M.R. and P.M. Das, 1993b. A Preliminary study on the use of poultry offal meal as dietary protein source for the fingerlings of Indian major carp, *Labeo rohita* (Hamilton). S.J. Kaushik and P. Luquet (Eds.), Fish Nutrition Practise, Institut National de la Recherche Agronomique, Paris, pp: 793-801.
18. Webster, C.D., K.R. Thompson, A.M. Morgan, E.J. Grisby and A.L. Gannam, 2000. Use of hempseed meal, poultry by-product meal and canola meal in practical diets without fish meal for sunshine bass (*Morone chrysops* x *M. saxatilis*). Aquaculture, 188: 299-309.
19. Davis, D.A. and C.R. Arnold, 2000. Replacement of fishmeal in practical diets for the Pacific white shrimp, *Litopenaeus vannamei*. Aquaculture, 185: 291-298.
20. Hasan, M.R. and M.R. Amin, 1997a. Effect of processing techniques on the nutritional quality of poultry offal meal. Bangladesh J. Fisheries, 20: 139-144.
21. Dong, F.M., R.W. Hardy, N.F. Haard, F.I. Borrows, B.A. Rasco, W.T. Fairgrieve and I.P. Forster, 1993. Chemical composition and protein digestibility of poultry by-product meals for salmonid diets. Aquaculture, 116: 149-158.
22. Metailler, R., 1987. Experiments in nutrition. A. Bruno, (Ed.), Nutrition and Marine Aquaculture, Tunisia, Lisbon, pp: 304-320.
23. Goddard, S., 1996. Feed Management in Intensive Aquaculture. Chapman & Hall. Press, New York, pp: 194.
24. Akyıldız, R., 1984. Yemler Bilgisi Laboratuvar Kılavuzu. Ank. Üniv. Zir. Fak. Yay: 895, Uygulama Kılavuzu III, Ankara Üniv. Basımevi, Ankara, pp: 236.
25. Tacon, A.G.J. and A.J. Jackson, 1985. Utilization of conventional and unconventional protein sources in practical fish feeds. C.B. Cowey, A.M. Mackie and J.G. Bell (Eds.), Nutrition and Feeding in Fish, Academic Press, pp: 119-145.
26. Davies, S.J., J. Williamson, M. Robinson and I.B. Robert, 1989. Practical inclusion levels of common animal by-products in complete diets for tilapia (*Oreochromis mossambicus*, Peters). Proc. Third Int. Symp. on Feeding and Nutr. in Fish, Toba, Japan, pp: 326-332.
27. Fowler, L.G., 1990. Feather meal as a dietary protein source during parr-smolt transformation in fall Chinook salmon. Aquaculture, 89: 301-314.
28. Hasan, M.R., M.S. Haq, R.M. Das and G. Mowlah, 1997b. Evaluation of poultry feather meal as a dietary protein source for Indian major carp, *Labeo rohita* Fry. Aquaculture, 151: 47-54.
29. Hardy, R.W., 2000. New developments in aquatic feed ingredients and potential of enzyme supplements. L.E. Cruz-Surez, D. Ricque-Marie, M. Tapia-Salazar, M.A. Olvera-Novoa and R. Civera-Cerecedo, (Eds.), Avances en Nutricion Acuicola V. Memorias del V Simposium Internacional de Nutricion Acuicola, Mérida, Yucaton, Mexico, pp: 216- 226.

30. Opstvedt, J., R. Miller, R.W. Hardy and J. Spinelli, 1984. Heat-induced changes in sulfhydryl groups and disulfide bonds in fish protein and their effect on protein and amino acid digestibility in rainbow trout (*Salmo gairdneri*). *J. Agricultural Food Chemistry*, 32: 929-935.
31. McCallum, I.M. and D. Higgs, 1989. An assessment of processing effects on the nutritive value of marine protein sources for juvenile chinook salmon (*Oncorhynchus tshawytscha*). *Aquaculture*, 77: 181-200.
32. Kaushik, S.J., 1995. Nutrient requirements, supply and utilization in the context of carp culture. *Aquaculture*, 129: 225-241.
33. Gallagher, M.L. and G. Degani, 1988. Poultry meal and poultry oil as sources of protein and lipid in the diet of European Eels (*Anguilla anguilla*). *Aquaculture*, 73: 177-187.
34. Weatherup, R.N. and K.J. McCracken, 1999. Changes in rainbow trout, *Oncorhynchus mykiss* (Walbaum), body composition with weight. *Aquaculture Res.*, 30: 305-307.